



**Brooksville, Florida Plant Materials Center**  
**Year 2003 Annual Technical Report**



**Issued May 2004**

## TABLE OF CONTENTS

<b>PLANT MATERIALS CENTER STAFF .....</b>	<b>3</b>
<b>STATE PLANT MATERIALS TECHNICAL ADVISORY COMMITTEE .....</b>	<b>3</b>
<b>MISSION AND MAJOR OBJECTIVES .....</b>	<b>4</b>
<b>PLANT EVALUATION PROCESS .....</b>	<b>4</b>
<b>SOILS .....</b>	<b>5</b>
<b>CLIMATE.....</b>	<b>5</b>
<b>INITIAL EVALUATIONS .....</b>	<b>6</b>
Switchgrass ( <i>Panicum virgatum</i> ) (FLPMC-P-0001-RA) .....	6
Wiregrass ( <i>Aristida stricta</i> ) (FLPMC-P-0337-WO).....	7
Hairawn Muhly ( <i>Muhlenbergia capillaris</i> ) (FLPMC-P-9236-RA).....	7
Lopsided Indiangrass ( <i>Sorghastrum secundum</i> ) (FLPMC-P-9602-RA) .....	8
Chalky Bluestem ( <i>Andropogon glomeratus</i> var. <i>glaucopsis</i> ) (FLPMC-P-9601-RA).....	9
Hairawn Muhly ( <i>Muhlenbergia capillaris</i> ) (FLPMC-P-0201-UR).....	10
White Muhly ( <i>Muhlenbergia capillaris</i> ) (FLPMC-P-0102-U) Cultivar Development .....	10
Annual Phlox ( <i>Phlox drummondii</i> ) (FLPMC-P-0202-CR) .....	10
<b>ADVANCED EVALUATIONS .....</b>	<b>12</b>
Blue Maidencane ( <i>Amphicarpum muhlenbergianum</i> ) (FLPMC-P-9604-WE).....	12
Eastern Gamagrass ( <i>Tripsacum dactyloides</i> ) (FLPMC-P-9605-RA) .....	13
Intercenter Strain Trial Partridge Pea (FLPMC-0208-WL).....	13
Intercenter Strain Trial Florida Paspalum (FLPMC-P-0463-WL).....	14
<b>ACTIVE STUDIES .....</b>	<b>15</b>
The Influence of Fertility and pH on Leaf Color of Blue-Green Eastern Gamagrasses (FLPMC-T-0004-RA) ....	15
Seed Production of Upland Native Plants (FLPMC-T-9902-CR) .....	15
Eglin AFB Native Species Erosion Control Mats (FLPMC-T-0105-CR).....	16
Department of Transportation Cooperative Wildflower Seed Increase (FLPMC-S-0104-OT).....	17
Lopsided Indiangrass ( <i>Sorghastrum secundum</i> ) Residue Management Study (FLPMC-07RNGE-igS) .....	17
<b>PLANT MATERIALS RELEASED BY THE BROOKSVILLE, FL PMC.....</b>	<b>19</b>
<b>PUBLICATIONS AVAILABLE FROM THE BROOKSVILLE, FL PMC.....</b>	<b>19</b>

Cover Picture: *Muhlenbergia capillaris* (Hairawn Muhly)

---

United States Department of Agriculture, Natural Resources Conservation Service, Plant  
Materials Center, 14119 Broad Street, Brooksville, FL 34601; Tel: (352) 796-9600

---

## PLANT MATERIALS CENTER STAFF

*Clarence Maura, Jr.*..... *Manager*  
*Vacant*..... *Agronomist*  
*Mary Anne Gonter* ..... *Biological Technician (Plants)*  
*Beverly Gordon* ..... *Secretary*  
*Robert Santucci*..... *Biological Technician*  
*Edmond Black*..... *Biological Technician*

### PLANT MATERIALS SPECIALIST

*Samuel A. Sanders*

### NAT'L PLANT MATERIALS SPECIALIST

*Richard White*

## STATE PLANT MATERIALS TECHNICAL ADVISORY COMMITTEE

*Greg Hendricks, State Resource Conservationist, Chairman* ..... *Gainesville, FL*  
*Jill Dobbs, District Conservationist* ..... *Ocala, FL*  
*Gene Fults, Rangeland Management Specialist* ..... *Kissimmee, FL*  
*Brian McGraw, District Conservationist*..... *Blountstown, FL*  
*Maelo Reyes, Resource Conservationist*..... *LaBelle, FL*  
*Don Surrency, Plant Materials Specialist* ..... *Thomson, GA*  
*Otto Bundy, Horticultural Systems* ..... *Parrish, FL*  
*Mike Camizzi, Forestry Technician*..... *Eglin AFB, FL*  
*Rosemarie Garcia, Cargill Fertilizer, Inc.* ..... *Ft. Meade, FL*  
*Dr. John Hovanesian, Coastal & Native Plants*..... *Milton, FL*  
*Dr. Oghenekome Onokpise, FAMU*..... *Tallahassee, FL*

---

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14<sup>th</sup> and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

---

## MISSION AND MAJOR OBJECTIVES

The mission of the Plant Materials Program is to provide timely and effective vegetative solutions for identified resource needs. Superior accessions of adapted plants are developed, tested and released to commercial growers along with production and management methodology. Plant materials and state-of-the-art plant science technology are promoted to assist in conservation of natural resources and meeting the objectives of environmental programs. The use of native plant materials is emphasized. The following are major objectives of the Brooksville Plant Materials Center (PMC):

- Improve and maintain water quality
- Control erosion on cropland and stabilize critical areas
- Improve forage on pastures and rangeland
- Improve wildlife habitat

## PLANT EVALUATION PROCESS

**Assemble Plant Materials** - Assemblies are planned to satisfy a specific objective(s) indicated in a project plan. Collections are made from a wide area within the occurrence of the species to insure diversity of ecotypes and variability within a species.

**Initial Evaluation** - The process of recording performance of the plant under controlled conditions. It allows the observance of characteristics and performance of the various collected plants, in order to select the most promising for the proposed conservation use. These plantings are normally done at the PMC, but off-center initial evaluation plantings can be done if it suits the purpose.

**Advanced Evaluations** - Intensive testing of selected plants that were superior in one or more attributes during the initial evaluation process. Cooperating agencies or other PMC's are encouraged to participate in this process. Plantings in areas where climatic conditions are significantly different than the PMC aids in determining range of adaptation for the plant materials.

**Final Evaluations** - Selections that exhibit superior qualities for the intended use are placed in field plantings on sites away from the PMC, under actual growing conditions.

**Release of New Plant Materials** - This is one of the final step in the process. The plant's usefulness for meeting conservation needs is documented. Insofar as possible, materials are released in cooperation with, or with concurrence of, cooperating agencies. Source identified, selected, tested, cultivar, and germplasm releases require less stringent evaluation and speed the release process.

**Breeder and Foundation Increase** - PMC's maintain breeder and foundation seed or plants of cultivars that have been cooperatively released, or they arrange for maintenance with agencies and organizations participating in the release.


## SOILS

Soil at the Florida PMC is predominately Kendrick Loamy Fine Sand. Other types of soil at the Center consist of Arredondo Fine Sand, Blichton Loamy Fine Sand, Electra Variant Fine Sand, Fleminton Fine Sandy Loam, Floridana Variant Loamy Fine Sand, Kanapaha Fine Sand, Nobleton Fine Sand, Sparr Fine Sand, and Wauchula Fine Sand.

## CLIMATE


There were 38 inches of rain during the first six months of 2003. Total rainfall for the year was 65.85 inches. The 20-year average rainfall was 56.5 inches.

**Table 1. Year 2003 total monthly rainfall.**



The last frost occurred on April 1. The first frost occurred on January 5. There were 353 frost-free days in 2003. The lowest temperature recorded at the Florida PMC in 2003 was 23° F, which occurred on January 24.

**Table 2. Year 2003 average monthly high and low temperatures.**



## INITIAL EVALUATIONS

### Switchgrass (*Panicum virgatum*) (FLPMC-P-0001-RA)

**Project Stage:** Establishment year of 3 to 4 years of initial evaluation and genetic selection. Increased and advanced evaluation phases are expected to take 5 or more years.

**Cooperators:** K. Quesenberry, Department of Agronomy, University of Florida, Gainesville, and M. Williams, USDA, Agricultural Research Service, Subtropical Agricultural Research Station, Brooksville, FL.

**Introduction:** Switchgrass has excellent potential for use in revegetating reclaimed minedlands and native areas, providing high quality livestock forage and wildlife food, controlling erosion and improving water quality. There are a few commercially available cultivars of switchgrass, such as Alamo, that are adaptable to Florida conditions. However, in initial studies, Florida ecotypes appear to have better performance than those from other states. The primary problem hindering development of a Florida cultivar of switchgrass has been poor seed production.

**Objective:** Develop one or more strains of Florida switchgrass with high seed production capabilities, and whose primary use will be for reclaiming native areas, wildlife food and cover, and controlling erosion. If selections are identified that produce high amounts of forage, these will be released for range and pasture improvement.

**Progress:** During the fall of 2000, a total of 88 accessions were collected from 42 counties in Florida in the form of plants and also seed if it was available. Sixteen accessions had been collected previously, so that the final assembly totaled 104 accessions. Seed was available from 80 accessions, and was planted in 6" cone trays in the greenhouse in December of 2000. Transplants of 104 accessions (originating from seed or from original plants if seed was not available) were planted in a crossing block at the PMC on August 2, 2001. Five plants were planted from each accession in a randomized complete block design. In the fall of 2004, each plant is to be evaluated for growth characteristics, and seed samples will be taken. Germination will be run on each sample. Plants will be separated by bloom dates (from early, middle and late).

Growth characteristics in the assembly are highly diverse. Differences, especially in biomass production, are probably due to differences in ploidy level. Higher ploidy levels are known to have a more robust growth habit. An attempt was made, through the assistance of Dr. Quesenberry, to identify ploidy levels of several accessions. However, chromosomes were relatively small and exact counts could not be made. As time permits, more attempts will be made to determine if there are indeed differences in ploidy level between the various accessions.

**Future Research Needs:** Once the superior cultivars are developed, they will need to be tested for performance and adaptation in Florida, and surrounding states, prior to release.

### **Wiregrass (*Aristida stricta*) (FLPMC-P-0337-WO)**

**Project Stage:** Develop new project plan and assemble genetically diverse accessions.

**Introduction:** Wiregrass is a warm-season perennial bunchgrass distributed throughout Florida and the southern portions of Mississippi through North Carolina. It is adapted to a broad range of soil and moisture regimes, from wet flatwoods to longleaf pine-turkey oak sandhills. Once established, it is very drought resistant and hardy. Wiregrass is considered to be one of the most important grasses in a pineland habitat, because of its ability to carry fire. In native situations, wiregrass contributes a large percentage of the fuel for understory burn management programs. Livestock readily graze new growth after a burn. Wiregrass also provides cover and nesting sites for wildlife. It does produce fair quantities of seed if old residues are removed during the growing season.

**Objective:** To develop one or more varieties of wiregrass with good seedling establishment characteristics and high seed production.

**Progress and Future Research Needs:** Results from previous work with wiregrass and consultations with plant breeders suggest that viable seed production could be rapidly improved in this species using genetic selection. Large assemblies of wiregrass plants from several stable populations (e.g., parks or preserves) need to be gathered to form the base population from which these selections are to be made. Collections of these species will begin in the spring of 2004. Once an assembly is established, individual plants will be evaluated for seed production, seed quality, and growth characteristics.

### **Hairawn Muhly (*Muhlenbergia capillaris*) (FLPMC-P-9236-RA)**

**Project Stage:** The second and final year of initial evaluation. Accessions with ornamental potential were selected for vegetative increase. Accessions with seed production potential were selected for placement in polycross nurseries to develop varieties with high seed production. Variety development and advanced evaluation for seed producers are expected to take 7 or more years. Ornamental varieties increased vegetatively may be released within 1 to 2 years.

**Introduction:** Muhly is a hardy warm-season perennial bunchgrass distributed throughout Florida and several states in the southeastern U.S.A. It is adapted to a broad range of sites from seeps and marshes to longleaf pine-turkey oak sandhills. It is more common on wetter sites. In its vegetative state, muhly looks very similar to wiregrass, and fills the same roll. Livestock and wildlife graze early growth. In native communities, it provides fuel for understory burn management programs and cover for wildlife. Because of its attractive purple inflorescence, it is becoming very popular for use in buffers, highway beautification and as a xeriscape ornamental. It is known to produce viable seed, but more information is needed on pollination methods and seed production characteristics.

**Objective:** To evaluate, develop, and release Florida native varieties of hairawn muhly for conservation use, especially erosion control, native area restoration, xeriscapes, and wildlife cover.

**Progress:** Ten accessions with superior viable seed production and growth characteristics were selected for placement in a polycross nursery to develop a composite variety with high seed production. Two other accessions were also identified with consistently high seed viability in 2000 and 2001. These accessions are 9060304, collected from Jackson Co., and 9060428, collected from Bradford Co. They are much smaller and bloom a month earlier than most other collections in the assembly. These 2 accessions will be placed in isolation to increase the seed for advanced evaluation.

In addition to development of seed producing strains, 11 accessions were also selected for ornamental potential. These types had superior bloom and foliage attributes that would make them valuable for use in roadside, buffer strip, and xeriscape plantings. They are to be increased vegetatively for advanced evaluation.

**Future Research Needs:** Once superior accessions have been increased, they are to be placed in advanced evaluation studies throughout Florida, and possibly in other states. Advanced testing of ornamental selections could occur within 1 to 2 years if enough material is available. Development of seed producing cultivars could take between 3 and 5 years before material is available for advanced evaluations. Other studies need to include testing pesticides for use in controlling aphids on seedheads. Seeding studies will also need to be conducted in order to develop successful seed establishment technology for hairawn muhly.

#### **Lopsided Indiangrass (*Sorghastrum secundum*) (FLPMC-P-9602-RA)**

**Project Stage:** Development of composite varieties for foundation seed stock. The development and increase phases are expected to take 3 years of the remaining 6 to 7 years of this project.

**Introduction:** There is a growing demand for seed sources of native species that can be used to restore native habitats. Lopsided indiangrass is one of the dominant grass species on native uplands in Florida. It is a warm-season perennial bunch grass, adapted to a wide variety of soils and hydrology regimes. It produces good quality livestock forage and is important for erosion control and also wildlife habitat. Lopsided indiangrass has good seed production and seedling vigor compared to other Florida native species.

**Objective:** To evaluate, develop, and release native Florida variety of lopsided indiangrass for conservation uses, especially for erosion control, native area restoration, and wildlife cover.

**Progress:** In 2002, seed was collected from all plots and used to establish seedlings for a foundation field. These seedlings from the composite (9060565) are currently being raised in the greenhouse. They will be put into a field for foundation seed harvest in the spring of 2004. Any additional seed produced by these blocks in 2004 or 2005 should also be collected and held in long-term storage for reestablishment of foundation fields. It is expected that at least 3 years will be necessary to produce enough foundation seed for advanced evaluations and eventual release.

**Future Research Needs:** Once foundation seed is increased, advanced evaluations may need to be conducted around Florida to determine performance and adaptability. Further studies also need to be conducted to find ways to increase the longevity of production fields. Possible areas

of study may include further residue management trials and inoculation of seedlings with antagonistic organisms, such as mycorrhizae.

**Chalky Bluestem (*Andropogon glomeratus* var. *glaucopsis*) (FLPMC-P-9601-RA)**

**Project Stage:** Development of a composite variety. Development and increase of a composite is expected to take another 2 years. Advanced evaluation is expected to require another 2 to 4 years before the material is released.

**Introduction:** Chalky bluestem is a native warm season perennial bunchgrass distributed throughout Florida, southern North Carolina, South Carolina and Georgia, and west to East Texas. It is adapted to flatwoods, seeps, and the margins of freshwater marshes and ponds. It produces high quality livestock forage and is thought to be one of the most palatable native grasses on flatwoods sites. It is also an important plant for upland water quality and erosion control. Chalky bluestem is a prolific seed producer, and will readily colonize disturbed areas in wet flatwood sites.

**Objective:** To evaluate, develop and release a Florida native variety of chalky bluestem for conservation use, especially erosion control, wetland restoration and wildlife cover.

**Progress:** Ten superior accessions were selected for increased and advanced testing. Seed from original collections stored in the seed cooler was used to start seedlings in the greenhouse in April of 2000. In February of 2001, these ten accessions were randomly planted together on an irrigated, poorly drained site to form a composite (9060565). Seed was collected with the Flail Vac in November of 2002. Average germination was 52% with 13 lb/A being collected.

Seed needs to be collected again from this nursery for the next 1 to 2 years, to obtain enough material for planting foundation fields. Extra seed should be stored in long-term storage to maintain stocks for future generations. It is expected that it will take 3 years to obtain enough seed from the foundation field for advanced evaluations and eventual release.

**Future Research Needs:** Once seed is available, the new cultivar needs to be tested for adaptation and performance throughout Florida, and possibly other states. Seed conditioning and seeding methodology also need to be developed for this species to insure establishment success.

### **Hairawn Muhly (*Muhlenbergia capillaris*) (FLPMC-P-0201-UR)**

**Project Stage:** Plants of selected accessions were planted at 3 locations around the state in 2002. Cooperators have agreed to water and weed plants as necessary. Plots were evaluated in the fall of 2003 and in 2004. Top performers will be released onto the commercial market.

**Introduction:** Hairawn muhly is a perennial native Florida bunchgrass. It grows 1 to 4 feet in height, depending on ecotype, and produces very showy seedheads that range in color from pink to purple. A rare white bloomed type has also been found in some populations. Hairawn muhly is becoming very popular in xeriscape plantings along roadsides and urban areas. Eleven accessions with desirable ornamental qualities were selected in an initial evaluation study of an assembly of hairawn muhly at the Brooksville PMC.

**Objective:** Determine the range of adaptation of 11 accessions of hairawn muhly in Florida in low maintenance settings. Evaluation is to include performance, ornamental qualities, and plant persistence in 3 different climatic zones on different soil types.

### **White Muhly (*Muhlenbergia capillaris*) (FLPMC-P-0102-U) Cultivar Development**

**Project Stage:** To increase these plants in preparation for release.

**Introduction:** A white bloomed muhly was found in a native population of pink bloomed plants. This white variety has tremendous potential for use in the commercial landscaping industry, especially for xeriscapes. However, seedling vigor decreased when the white variety was grown in isolation, possibly because of inbred depression. Tests were conducted to determine pollination mechanisms. It appears that white blooms occur because of a recessive gene. White bloomed types also have lower vigor than pink bloomed types, and almost all seed is sterile when plants are forced to self-pollinate. Therefore, this accession needs to be released as a vegetatively propagated cultivar.

**Progress:** Adaptation studies were initiated in 2002, along with the ornamental muhly advanced evaluation project.

**Future Research Needs:** To increase this accession for commercial release.

### **Annual Phlox (*Phlox drummondii*) (FLPMC-P-0202-CR)**

**Project Stage:** To develop a variety of Florida annual phlox (*Phlox drummondii*) with unique flowers.

**Introduction:** Seed collected in 2001 was planted into trays in the PMC greenhouse in 2002. When the seedlings emerged, many of them had unique flower patterns, shapes, and colors, although there were also some common types. Seedlings were put in the shadehouse in the spring, so that natural pollinators could access them. All types were allowed to cross, and seed was hand collected as it ripened.

**Objective:** To develop a cultivar containing these unusual flower types.

**Progress and Future Needs:** Seed collected from the shadehouse in 2002 was planted into trays in the greenhouse in November of 2002. Once seedlings are large enough, they will be separated and planted into 6" pots. After the danger of frost is past, they will be placed in the shadehouse. Once plants bloom, common types will be pulled out so they will not cross with the unique types. Seed will be hand collected as it ripens. Collected seed will then be planted into trays in November of 2003 and the process outlined above will be repeated. These cycles will be continued until adequate seed is available for commercial release.

## ADVANCED EVALUATIONS

### Blue Maidencane (*Amphicarpum muhlenbergianum*) (FLPMC-P-9604-WE)

**Project Stage:** Increase of superior accessions and advanced evaluation. Advanced evaluations are expected to take 2 to 3 more years.

**Introduction:** Blue maidencane is a native, warm-season perennial rhizomatous grass distributed throughout Florida and coastal areas of Georgia and South Carolina. It is adapted to acid or neutral sandy soils that are wet for part of the year. It grows in sloughs and intermittently ponded areas in flatwoods range sites. Cattle preferentially graze this species, which produces high quality forage. Because it often forms solid stands, it is important for erosion control and maintaining water quality in fresh water systems.

**Objective:** To evaluate, develop, and release a Florida native variety of blue maidencane for conservation use, especially for erosion control and wetland restoration purposes.

**Progress:** From 1996 through 1998, a total of 157 accessions were assembled from throughout the state of Florida in the form of root and shoot stock. Initial evaluation plots were planted at the PMC in March of 1999. Because this species spreads aggressively by rhizomes, plots could only be evaluated for one year before accessions began growing together. Eleven superior accessions were selected for advanced evaluation (Table 3). In March of 2000, accessions were increased by planting rhizomes in tubs. Accessions 9059859, 9060309, and 9060311 were combined to form accession 9060489, since they have very similar performance and come from the same basic location. Accessions 9059866, 9060066, and 9060067 were also combined to form accession 9060490 for the same reasons.

**Table 3. Eleven superior blue maidencane accessions selected for increase and evaluation.**

Accession No.	County	Collector
9059859	Pasco	Deal/Pfaff
9060309	Pasco	Deal/Pfaff
9060311	Pasco	PMC
9059866	Charlotte	PMC
9060066	Sarasota (Myakka State Park)	Perry/Lackman
9060067	Sarasota (Myakka State Park)	Perry/Lackman
9059869	Palm Beach	PMC
9059956	Madison	PMC
9059971	Citrus	PMC
9060008	St. Johns	PMC
9060295	Polk	PMC

**Future Research Needs:** Three or more additional advanced evaluation sites are needed in Florida before superior accessions can be selected for release.

### **Eastern Gamagrass (*Tripsacum dactyloides*) (FLPMC-P-9605-RA)**

**Project Stage:** Increase of superior accessions and advanced evaluation. Advanced evaluations are expected to take another 1 to 2 years.

**Introduction:** Eastern gamagrass is a warm-season perennial bunchgrass with a broad area of distribution throughout the U.S.A., including all of the southern states. It has received a great deal of attention in recent years because of its tremendous forage production. It typically grows in moist fertile sites, and is often found lining the edges of canals and freshwater bodies in Florida. Florida ecotypes are markedly different than strains from other states, in terms of growth and winter dormancy characteristics. There is a demand in Florida for commercial seed sources of local ecotypes. In 1996 through 1998 an assembly of Florida eastern gamagrasses was evaluated for forage and seed production characteristics. Four accessions were selected with superior performance in these two categories; they were 9059213 (Clay Co.), 9059264 (Dixie Co.), 9059266 (Polk Co.), and 9059287 (Citrus Co.). All four accessions are apomictic and will not out-cross.

**Objective:** To evaluate, develop, and release one or more accessions of eastern gamagrass for conservation use including for buffer strips, pasture and rangeland improvement and wildlife food and cover.

**Progress:** Advanced evaluation plots of all four eastern gamagrass accessions were planted at the Collier, Polk, and Hamilton Co. sites. Evaluations will continue for the duration of the project.

**Future Research Needs:** Two or more additional advanced evaluation sites are needed in Florida before superior accessions can be selected for release. If seed production of this species is to become economical, production field management technology needs to be developed, especially in such areas as fertility and plant spacing.

### **Intercenter Strain Trial Partridge Pea (FLPMC-0208-WL)**

**Project Stage:** To determine adaptation of Partridge Pea cultivar and new releases to areas in the Southeast and southern Plains.

**Introduction:** The annual legume partridge pea is an excellent food source for wildlife and is also suitable for planting on many critical areas. The only commercially available cultivar of partridge pea is 'Comanche', released by the Texas PMC in Knox City. Lark selection, was released by the Kansas PMC in Manhattan, Kansas. The full range of adaptation of these two per-varietal releases is not known. This inter-center strain trial will determine the survival and growth potential of these releases at sites throughout the Southeast and southern Plains states, using Comanche as the standard of comparison.

**Objective:** The information obtained by this study will be used to support a cultivar release of Lark Selection. Technical notes, release documentation, revise FTOG standards.

**Progress:** Seed samples will be obtained from the relevant PMC's. Plots will be planted at MSPMC, ARPMC, ETPMC, TXPMC, STPMC, GAPMC, FLPMC, and other locations as available. Plot size to be planted will be 5' X 10' with 3 replications. Plots will be prepared and managed according to standard practices used at the testing sites. The planting rate will be 6 PLS pounds of seed/A, and seeds will be inoculated with proper inoculant prior to planting.

### **Intercenter Strain Trial Florida Paspalum (FLPMC-P-0463-WL)**

**Project Stage:** To determine area of adaptation for #9043874 ('Harrison' Florida paspalum select germplasm) in the southern United States.

**Introduction:** Florida paspalum is a native warm season perennial grass (Hatch, 1990). This grass produces a large seed that is eaten by gamebirds (Grelen, 1984). Florida paspalum is adapted from New Jersey, southeastern United States, Midwestern states, and east Texas (Hitchcock, 1950). However, the specific adaptation area of #9043874 has not been determined. This accession was collected from Harrison County, Texas by NRCS personnel, Ross Brown and Paul Leggett. This accession has been evaluated against other Florida paspalum accessions at the East Texas PMC.

**Objective:** The information gathered from this study will be used to support a cultivar release of Harrison select germplasm.

**Progress:** Rod rows will be planted in the spring at the participating PMC's: TXPMC, STPMC, MSPMC, FLPMC, GAPMC, ARPMC. Rod rows will be planted and managed according to practices used at the individual PMC's. PMC staff will collect the requested data as noted on the evaluation form.

## ACTIVE STUDIES

### The Influence of Fertility and pH on Leaf Color of Blue-Green Eastern Gamagrasses (FLPMC-T-0004-RA)

**Project Stage:** Third year of a three-year study.

**Introduction:** 'Martin Germplasm' and 'St. Lucie Germplasm' eastern gamagrass have potential for use in ornamental settings because of their blue-green colored foliage. However, at some sites, this attractive foliage color was not always noticeable. The factors that influence foliage color in these two accessions need to be determined.

**Objective:** To investigate the influence of different pH and fertility levels on the foliage color of Martin and St. Lucie eastern gamagrass.

**Progress:** On August 1, 2001, plots were established on 3 irrigated sites at the Brooksville PMC. Previous pH samples taken from these sites ranged from 4.9 to 6.0. Three fertility treatments will be applied in 2002, including 0, 50, 100 and 200 lbs/ac. of 10-10-10. Each plot consists of 2 or 3 plants with 1 border plant between treatments.

**Future Research Needs:** As mentioned under the Advanced Evaluation section of this report, it would be useful to make several demonstration plantings to determine how these two accessions perform under a variety of conditions.

### Seed Production of Upland Native Plants (FLPMC-T-9902-CR)

**Project Stage:** Increase of selected native accessions.

**Introduction:** Based on results of direct seeding studies on reclaimed mined lands and at the PMC, 2 species were selected for further increase and possible commercial release. These species are splitbeard bluestem (*Andropogon ternarius*) (formerly thought to be *Andropogon arctatus*) and pinewoods dropseed (*Sporobolus junceus*).

**Objective:** To increase, test and release native species for restoring dry sandy uplands.

**Progress:** In 2003, *Andropogon ternarius* plants were well established, and all bloomed in the fall. Seed was collected with the Flail Vac in November. Only 1 harvest was made to collect the maximum amount of seed. Timing was perfect for seed collection this year. Making the collections at lower brush speed significantly increased the amount of seed collected, and reduce the amount of inert matter that must be removed from the seed. A hammermill was very effective at dehulling seed. *Sporobolus junceus* plants are in the development stages right now.

**Future Research Needs:** Tests need to be conducted on both *Andropogon ternarius* and *Sporobolus junceus* to determine if debearding or dehulling damages seed and reduces viability.

Management of residue and top growth of both species also needs to be tested to find ways to maintain stands and increase seed production.

**Eglin AFB Native Species Erosion Control Mats (FLPMC-T-0105-CR)**

**Project Stage:** Third year of a four-year study.

**Introduction:** Natural resources personnel at Eglin Air Force Base wish to restore borrow pits and other critical areas with native species. Because many of these areas have very steep slopes, erosion is a serious problem, and native species often grow too slowly to stabilize critical areas. Biodegradable mats made out of coconut fiber (coir) are often used to stabilize severe slopes. Establishing grasses and forbs on these mats prior to placement on the site could potentially be used to revegetate critical areas with slow growing native species.

**Objective:** Develop methods of establishing native species on coir mats, and determine if these vegetated mats can successfully colonize and control erosion on critical areas with high slopes.

**Progress:** On December 4 and 5 of 2001, Eglin personnel supplied a seed stripper pulled with an ATV to collect seed from two native sites. In addition, PMC personnel made hand collections of selected grasses and forbs. The two sites ranged from xeric to mesic and were dominated by wiregrass. Growing season burns had effectively stimulated the wiregrass to flower and produce viable seed. Seed germination tests were conducted at the PMC laboratory (Table 4) in order to calculate seeding rates. Coir blanket mats (1/4" to 1/2" thick) measuring 3' x 40' were placed on asphalt and plastic at the PMC and were framed with 2" x 4" wood. Mats were seeded on February 14, 2002, and covered with approximately 1/2" of potting soil. Because of cooler weather, seed did not begin to germinate for 1 to 2 months. Once plants have developed a strong root system, they were transported to Eglin AFB and planted in June 2003.

**Table 4. Laboratory germination of various native species collected from Eglin AFB in 2002.**

Site	Species	Common Name	Seed Viability (%)
Indigo Pond	<i>Andropogon gyrans</i>	Elliot bluestem	46
	<i>Andropogon ternarius</i>	Splitbeard bluestem	32
	<i>Liatris spp.</i>	Gayfeather	48
	<i>Pityopsis</i>	Grass leaf golden aster	42
	<i>Schizachyrium stoloniferum</i>	Creeping bluestem	4
Range 78	<i>Aristida beyrichiana</i> (stripper mix)	Wiregrass	39
	<i>Aristida beyrichiana</i> (hand clipped)	Wiregrass	35
	<i>Schizachyrium stoloniferum</i>	Creeping bluestem	0

**Future Research Needs:** Once established, mats will need to be evaluated for 2 to 3 years to determine if this is a viable method of stabilizing critical areas with native species.

## **Department of Transportation Cooperative Wildflower Seed Increase (FLPMC-S-0104-OT)**

**Project Stage:** Third year of a three- to four-year study.

**Introduction:** The Florida Department of Transportation (DOT) would like to use Florida ecotypes of native wildflowers to beautify roadways and reduce mowing costs. However, there are very few Florida sources of wildflowers on the commercial market. Dr. Jeff Norcini of the University of Florida, IFAS, has been working under a grant from the DOT to develop Florida seed sources, and has asked the Brooksville PMC to help increase the seed for commercial growers.

**Objective:** To develop propagation and cultural management protocols, develop harvest methods, and increase seed stocks of two species of Florida wildflowers.

**Progress:** Seed of yellowtop (*Flaveria linearis*) collected from a south, central and north Florida location was brought to the PMC in the summer of 2001. Seed was started in flats in the greenhouse, and seedlings were transplanted to cone trays. Once seedlings develop a vigorous root system, they are being transplanted to the field. It has been noted that the south and north ecotypes of yellowtop have displayed a completely different growth habit. An accession of blackeyed susan (*Rudbeckia hirta*) is also being increased in a similar manner. A specialized vacuum harvester is being developed to collect the seed. Since blackeyed susan is a biennial, and yellowtop is a perennial, seed increase fields are expected to be productive for 2 or more years.

**Future Research Needs:** Herbicide tolerance information needs to be developed for these two wildflower species.

## **Lopsided Indiangrass (*Sorghastrum secundum*) Residue Management Study (FLPMC-07RNGE-igS)**

**Project Stage:** This is a residue management study that will continue for 1 to 2 years. Seed will be collected, depending on how long the majority of the plants survive.

**Introduction:** Lopsided indiangrass is an important component of native uplands in Florida. It has good seed production and has established well in critical area plantings on reclaimed minedlands. Unfortunately, seed production plots have been very short-lived at the Brooksville PMC, lasting only about 3 years on irrigated sites. Soil pathogens are most likely the cause of plant death, although this species may naturally be a short-lived perennial.

**Objective:** To determine how residue management method affects viable seed production and plant persistence.

**Progress:** Study plots were imposed on the composite Acc. 9060564 polycross nursery which was planted February 21, 2001, using seedlings from 6' containers. All plots are to be fertilized annually with approx. 50 lb/A 10-10-10 at the time of spring growth. Winter burn treatments are

to be applied approx. January before plants begin spring regrowth. Winter clip treatments are to be applied at the same time as burn treatments. Summer clip treatments are to be applied in late June prior to production of reproductive tillers.

**Future Research Needs:** Seed collected from this project will be used to establish the foundation field for Acc. 9060564.

## PLANT MATERIALS RELEASED BY THE BROOKSVILLE, FL PMC

<u>Year</u>	<u>Species</u>	<u>Cultivar</u>	<u>Cooperating Agency</u>
1944	<i>Paspalum notatum</i> (Bahagrass)	Pensacola	GA PMC
1960	<i>Panicum texanum</i> (Texas millet)	Artex	N/A
1962	<i>Lupinus elegans</i> (Mexican lupine)	Armex	N/A
1963	<i>Lupinus angustifolius</i> (Blue lupine)	Orlando	N/A
1969	<i>Aeschynomene americana</i> (American joint vetch)	F-149	N/A
1978	<i>Hemarthria altissima</i> (Limpograss)	Bigalta Greenalta Redalta	Univ.FL-I.F.A.S. “ “ “ “ “ “
1978	<i>Arachis glabrata</i> (Perennial peanut)	Florigraze	N/A
1985	<i>Arachis glabrata</i> (Perennial peanut)	Arbrook	FL Agri. Exp. Sta.
1990	<i>Spartina patens</i> (Marshhay cordgrass)	Flageo	GA PMC & Ft. Valley Agri. College
1991	<i>Helianthus debillis</i> (Beach sunflower)	Flora Sun	N/A
1992	<i>Panicum amarum</i> (Bitter panicum)	Northpa Southpa	N/A N/A
1994	<i>Spartina patens</i> (Marshhay cordgrass)	Sharp	GA PMC
1995	<i>Zea mexicana</i> (Mexican teosinte)	Chapingo	N/A
1996	<i>Panicum virgatum</i> (Switchgrass)	Miami Wabasso Stuart	N/A N/A N/A
1998	<i>Panicum hemitomom</i> (Maidencane)	Citrus	N/A
2000	<i>Tripsacum dactyloides</i> (Eastern gamagrass)	Martin St. Lucie	N/A N/A
2002	<i>Arachis glabrata</i> ( Perennial peanut)	Brooksville67	N/A
2002	<i>Arachis glabrata</i> (Perennial peanut)	Brooksville 68	N/A
2003	<i>Liatris elegans</i> ( Blazing star)	Floral Passion	N/A

## PUBLICATIONS AVAILABLE FROM THE BROOKSVILLE, FL PMC

	1997	Technical Note No. 35: Collecting Plant Materials
	1997	Plant Materials Program Fact Sheet
	1997	Florida Native Plant Collection, Production and Direct Seeding Techniques: Interim Report
June 1995 - Oct. 2003		Semi-Annual Newsletter: Sunshine State's PMC Impact
	1998	Forage Species on Sprayfields – Fact Sheet
	2000	Fact Sheet: Gully Stabilization in North Florida
	2003	Plant Materials Center, Brooksville, Florida - Visitor Information
	2002	Florida Native Seed Production Manual
Through 2003		Annual PMC Activity Reports
Through 2003		Progress Report of Activities