

Elsberry Plant Materials Center

2010 Annual Technical Report



Ozark Gerplasm Little Bluestem (*Schizachyrium scoparium*)

PLANT SOLUTIONS FOR CONSERVATION NEEDS

Visit us at: <http://plant-materials.nrcs.usda.gov> or <http://www.mo.nrcs.usda.gov>



United States Department of Agriculture
Natural Resources Conservation Service

‘Helping People Help the Land’

**ELSBERRY PLANT MATERIALS CENTER
2010**

Advisory Committee

**J. R. Flores, State Conservationist, Missouri, Chairman
Richard Sims, State Conservationist, Iowa
William J. Gradle, State Conservationist, Illinois**

Resource Personnel

**Dwaine Gelnar, State Resource Conservationist, Missouri
John D. Meyers, State Resource Conservationist, Iowa
Richard G. Hungerford, Jr., State Resource Conservationist, Illinois**

National Program Leader

John M. Englert

Central Region Plant Materials Specialist

Joel L. Douglas

Plant Materials Specialist

Jerry U. Kaiser

Plant Materials Center Personnel

**Ronald L. Cordsiemon II, Plant Materials Center Manager
Donald D. Tapley, Agricultural Science Research Technician
Nicholas S. Adams, Agricultural Science Research Technician
Garry Stewart, Agricultural Science Research Technician Aid**

Earth Team Volunteers

**Jimmy Henry
Bob Laird**

ELSBERRY PLANT MATERIALS CENTER

VISION – Excellence in developing plant science technology to help people help the land.

MISSION – To assist land users, federal, state and local partners, and industry growers in Illinois, Iowa and Missouri, in protecting, conserving, and improving natural resources by providing plant materials and plant related technology.

STRATEGY – The Elsberry Plant Materials Center vision and mission is advanced through evaluating and selecting superior plants, developing cultural and management technology, promoting the use of plants and related technology, through field and demonstration plantings, leading tours, and training NRCS employees and others on plant science technology. Plant Materials Committees in Illinois, Iowa and Missouri identify and prioritize plant materials technology needs and the three state conservationists establish direction and funding for the center.

GOALS – The Elsberry Plant Materials Center (PMC) provides plants for conservation, produces foundation seeds and plants or their equivalent, and promotes their use in solving natural resource problems on both private and public land. Beneficial uses for these plant materials include livestock forages, biomass and timber production, carbon sequestration, air quality, erosion reduction, wetland restoration, wildlife food and cover, water quality improvement, stream bank and riparian area protection, and other unique conservation needs. In addition to conservation plant release, the PMC also develops establishment and management technology for successful use of plants in resource conservation programs.

TACTICS – Specialists at the center identify plants that show promise for addressing a specific conservation need, develops related technology and test their performance in the field. After species are proven beneficial for solving the conservation problem, they are released to the private sector for commercial production for general public use. Opportunities for success are to continue working through NRCS field offices, soil and water conservation districts and other conservation partners who come in contact with clients in need of special plants and related technology for special situations. USDA programs that emphasize and utilize plant materials and plant science technology include Conservation Technical Assistance, Environmental Quality Incentive Program, Conservation Reserve Program, Wetlands Reserve Program, Grassland Reserve Program, and Wildlife Habitat Incentive Program with some opportunities in the Conservation Security Program, Resource Conservation and Development, and the Small Watershed and Flood Prevention program. The Elsberry Plant Materials Center also has the opportunity to support increasing interest in urban conservation by providing plants with unique landscape architecture value in addition to their erosion and sediment control, water and air quality benefits. Emerging opportunities in the agricultural sector lie in biomass for bioenergy production and carbon sequestration.

2010
Annual Technical Report
Elsberry Plant Materials Center
Elsberry, Missouri

<i>Introduction</i>	5
<i>Plant Materials Center Operations</i>	5
<i>Climatic Data</i>	7
<i>Study Activities:</i>	
<u>29I093R</u> – Miscellaneous Herbaceous Plant Evaluation	8
<u>29I097G</u> – Assembly and Evaluation of Big Bluestem, <i>Andropogon gerardii</i>	13
<u>29I108G</u> – Assembly and Evaluation of Low Growing Rhizomatous Switchgrass for Use in Waterways, Filter Strips and Other Conservation Uses	23
<u>29A116W</u> – Evaluation of Miscellaneous Trees and Shrubs	31
<u>29I124G</u> – Production of Native Iowa Ecotypes of Grasses and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Grasses and Forbs are Now Being Planted	35
<u>29A137O</u> – Wetland/Riparian Propagation	41
<u>29I141G</u> – Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i>, Michx.	54
<u>29I142G</u> – Production of Native Missouri Ecotypes of Grasses, Legumes, and Forbs for Roadsides, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted	74
<u>MOPMC-P-0001, WO, WL, WE</u> – Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i>, Michx.	80
<u>MOPMC-T-0105, PA</u> – Compatibility Study Using Native Warm Season and Cool Season Grasses with Native Legumes and Forbs	106
<u>MOPMC-T-0106, BU</u> – Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips	117
<u>MOPMC-P-0613, PA, WL</u> – Evaluation and Release of a Shade Tolerant Big Bluestem, <i>Andropogon gerardii</i>, L., for Silvopasture	131

Table of Contents - continued	
<u>MOPMC-T-0716</u> – In-Field Weathering Effects on Biomass of Biomass Yield and Biofuel Quality of Warm Season Grasses	136
<u>MOPMC-P-0717, PA, WL</u> – Evaluation and Release of a Shade Tolerant Little Bluestem for Silvopasture	145
<u>MOPMC-T-0718-WE, WL</u> – Evaluation of Flood Tolerance of Planted Oak Seedlings Derived from Different Seed Origins	148
<u>MOPMC-P-0820-UR</u> - Evaluation and Release of Native Plants for Urban Landscaping	150
<u>MOPMC-P-0821-BF</u>- Evaluation and Release of an Iowa Source, stiff stemmed Indiangrass, <i>Sorghastrum nutans</i>, for Biofuel.	152
<u>MOPMC-P-0822-PA,WL</u> - Inter Center Strain Trial – Yield and Persistence of 11 Big Bluestem Sources in Kansas, Missouri, Arkansas, and Mississippi	163
Herbaceous and Woody Seed and Plant Production 2009	168
2010 Plant Releases from the Elsberry PMC (Notice of Release)	
Ozark Little Bluestem	170
Releases From the Elsberry PMC to Date	177
Studies/Projects at the Elsberry PMC 1958 to Date	181

INTRODUCTION

The Elsberry Plant Materials Center (PMC) was established in June 1934 and is the oldest Center in the nation. It is one of 27 PMC's in the United States. The Center is located approximately 60 miles northwest of St. Louis, Missouri, on Highway 79. It includes 243 acres of various soil types. The Elsberry PMC primarily serves Illinois, Iowa and Missouri; however, it makes significant contributions to other states in the Midwest region.

Emphasis is focused on using native plants as a healthy way to solve conservation problems and protect ecosystems. The program seeks to address priority needs of field offices and land managers in both public and private sectors by working with a broad range of plant species, including grasses, forbs, legumes, trees, and shrubs.

The Elsberry PMC assembles tests, selects and develops improved plants and reliable techniques for successfully establishing and maintaining plants for conservation uses.

Of particular importance is finding suitable plants for wetland situations, high traffic areas, wildlife food and habitat, farmstead and field windbreaks, wind barriers, pastures, landscape and beautification, roadside restoration, riparian plantings, woodland, and erosion control on cropland. Each of the three states served by the Center has identified its plant materials problems, needs and priorities. PMC activities are directed toward meeting the needs and priorities set forth in the states' long-range plans. As early as 1939 the Center began searching for plants to respond to specific conservation problems. During the PMC's earlier existence it produced 10,000,000 seedlings for use in windbreaks during the dust bowl era. Today the Elsberry Plant Materials Center is still striving to solve a new realm of conservation problems in an ever changing world.

PLANT MATERIALS CENTER OPERATIONS

The Center's operations are carried out in accordance with policies set forth in the National Plant Materials Handbook.

Guided by the Center's Multi-Year Business Plan, plant species are collected (mainly local field collections [95%]). Other collections come from locations within the species range in the United States. Center personnel then prepare the seed/plant for planting. Each collection is given an identification number (accession) and planted in a uniform nursery. Initial evaluation data is recorded on such factors as seedling emergence and vigor, rate of growth, disease and insect resistance, and ability to spread. Also recorded are date and amount of bloom, seed production, winter hardiness, and foliage characteristics. Selections are made and seed increased for advanced evaluation plantings. Field plantings are then conducted to determine plant performance and soil and climatic adaptation throughout its intended area of use. Evaluations are made comparing selected candidate accessions with "standards of comparison" such as cultivars or varieties that are already in the commercial market, or other species used for the same purpose.

After several years (10-15) of evaluation, selected accessions are cooperatively released with the USDA-Agricultural Research Service (ARS), State Agricultural Experiment Stations, Conservation Commissions, Universities, Departments of Transportation, and/or other interested agencies. The Center releasing a named variety is responsible for maintaining the breeder and foundation seed. These fields undergo annual inspections by the Missouri Crop Improvement Association to insure that seed is available to commercial producers and ultimately to the public for solving conservation problems.

Additional avenues have been established and used by the Plant Materials discipline to release plants to the commercial market: Source Identified Releases, Selected, and Tested Releases. These three new avenues provide a quicker release of plants as compared to cultivar release (10-15 years).

The Elsberry Plant Materials Center has released over 90 plants during its 76-year history. In 2010, Ozark Germplasm little bluestem (*Schizachyrium scoparium*) was released as selected class release. Ozark Germplasm little bluestem is a selection of many collections from southern Illinois and southern Missouri. It was selected for its forage quality and production. Currently, the Elsberry Plant Materials Center has 86 active releases, 82 of them are native to the Elsberry PMC service area. For more information regarding Elsberry PMC releases, please contact the Elsberry Plant Materials Specialist.

**CLIMATIC DATA – CALENDAR YEAR 2010
TEMPERATURE (Fahrenheit)**

<u>Month</u>	<u>78 Year Monthly High Average</u>	<u>2010 Monthly High Average</u>	<u>2010 Monthly High Departure</u>	<u>78 Year Monthly Low Average</u>	<u>2010 Monthly Low Average</u>	<u>2010 Monthly Low Departure</u>
January	38.07	31.35	-6.72	18.46	16.39	-2.07
February	43.01	36.29	-6.72	22.43	19.86	-2.57
March	54.05	57.87	+3.82	36.55	36.10	-0.45
April	66.79	75.00	+8.21	42.21	48.17	+5.96
May	76.46	76.61	+0.15	56.60	55.48	-1.12
June	85.35	88.47	+3.12	70.76	67.67	-3.09
July	89.39	89.55	+0.16	64.70	69.45	+4.75
August	87.53	90.61	+3.08	62.76	68.06	+5.30
September	80.34	80.50	+0.16	49.80	58.20	+8.40
October	69.22	72.84	+3.62	43.30	43.42	+0.12
November	50.93	59.53	+8.60	32.55	34.03	+1.48
December	41.91	35.52	-6.39	22.90	20.00	-2.90
Total 2010	65.25	66.18	+0.93	43.59	44.74	+1.15

	2010	Typical
Last Killing Frost (26° and below)	<i>March 6</i>	<i>April 15</i>
First Killing Frost (26° and below)	November 3	October 15
Number of Frost-Free Days	242	184

**CLIMATIC DATA – CALENDAR YEAR 2010
Precipitation (Inches)**

<u>Month</u>	<u>80 Year Average</u>	<u>2010 Total</u>	<u>Departure</u>
January	1.91	1.96	+0.05
February	1.96	2.45	+0.49
March	3.15	3.07	-0.08
April	3.63	2.04	-1.59
May	4.11	4.15	+0.04
June	3.73	3.21	-0.52
July	3.54	6.78	+3.24
August	3.35	4.50	+1.15
September	3.37	5.91	+2.54
October	3.03	0.60	-2.43
November	2.90	3.37	+0.47
December	2.40	1.05	-1.35
Total	36.43	39.09	+2.66

Study: 29I093R

Study Title: Miscellaneous Herbaceous Plant Evaluation.

Study Leader: Bruckerhoff, S. B.

Introduction:

Plants arrive at the Plant Materials Center (PMC) from many sources and for many different purposes. Most of the plants are assigned to a specific study. Plants are also received that are not tied to a specific study. These can be from other PMC's for area of adaptation or plants in advanced stages of evaluation. Plants are received from individuals who are interested in an unfamiliar species or a plant with unusual characteristics. Many species existing on the center are not involved with an active study addressing a specific problem.

Problem:

Keeping track of numerous miscellaneous plants around the PMC without an organized evaluation system became inefficient. This study organizes miscellaneous plant material coming into the center for evaluation.

Objective:

Evaluate winter hardiness, insect and disease resistance, and vigor of plants for climatic adaptation. Plants brought in for other specific reasons like forage production, landscape beautification, shoreline stabilization, etc., will be evaluated accordingly.

Procedure:

As miscellaneous plants are received at the center, they are assigned an accession number and as much background information as available or necessary are documented. The accession is then assigned a location for planting that best suits its needs for evaluation. Plants are evaluated as necessary. Many plants are left for plant identification sessions or demonstrations for several years.

Discussion:

1984-1990

This study was initiated in April 1984 in the PMC pipeline area. There are approximately 150 different accessions of the following species of plants: Indiangrass, switchgrass, big bluestem, purpletop, little bluestem, buffalograss, wheatgrass, fescue, timothy, ryegrass, redtop, orchardgrass, kura clover, blackeyed susan, and lespedeza. Factors involved in evaluations dealt with area of adaptation.

1991-1994

Approximately 75 accessions were added during 1991. Forty of them were warm season grasses used in three FEP (Field Evaluation Planting) variety studies: 29A111G, 29A118G, and 29A127G. Twenty-six were accessions of common cool season grasses and legumes used for pasture and hay in the three-state area. These were commonly used for plant identification sessions.

1995-1998

The accessions added in 1997 are being looked at for forage. They include 'Steadfast' birdsfoot trefoil, 'Mandan' Canada wildrye, and several bermudagrasses including Hardy and OK-74-12-6. Also zoysia grass, centipedegrass, and buffalograss from the Fort Leonard Wood Wear Tolerance Study are being looked at for adaptation. Several big bluestem accessions from Study 29I097G are being evaluated as landscape plants.

1999

The accessions added in 1999 are a Lincoln County Missouri collection of Virginia wildrye and a Crawford County Missouri collection of Virginia wildrye variation geneses. These species are being looked at for shade tolerance for riparian areas and covercrop for tree plantings.

2000

No new accessions were added in 2000. Two species that are getting the most interest are the Lincoln County accessions of Virginia wildrye and 'Tufcote' bermudagrass.

The Lincoln County accession of Virginia wildrye is a shade tolerant cool season grass that has potential for a cover crop for woody plantings as well as a possible buffer species along riparian areas. This accession should be in commercial production and available soon.

The 'Tufcote' bermudagrass accession was tested at Fort Leonard Wood for wear tolerance and showed very good potential. It could be used on playgrounds, sports fields, lawns, as well as having potential for high livestock use areas. This species is not native and does show potential for spreading so it should not be planted in areas where it could escape and cause problems.

2001

Three new species of native legumes were added in 2001. Native legumes are seldom used in mixtures with warm season grasses planted for pastures primarily because of their cost, lack of availability, and lack of knowledge on which ones will perform best in a mixture.

The following species were planted for observational evaluation: goats rue, *Tephrosia virginiana*; sensitive brier, *Schrankin uncinata*; and Sampson's snakeroot, *Orbexilium pedunculatum*.

The Lincoln County Missouri collection of Virginia wildrye, accession 9083169, has shown excellent vigor and seed production. Forage quality is comparable to tall fescue, spring

green-up earlier than tall fescue and seedhead emergence is approximately two weeks later than tall fescue. This accession is scheduled for release in 2002.

2002

One new collection was planted in the miscellaneous block. Accession 9083240, western wheatgrass, *Pascopyrum smithii*, was planted as greenhouse plugs May 10, 2002. This material was collected in Audrain County, Missouri.

The Lincoln County Missouri collection of Virginia wildrye, accession 9083169, was released as a selected class and given the name Cuivre River. The Cuivre River selection has early vigorous growth that is earlier than tall fescue. Booting occurred at the end of May to the first week of June at Elsberry. This is approximately two weeks later than tall fescue.

Although Cuivre River was released as a selection and only limited testing has been done, its anticipated uses are wildlife food/cover, plant diversity in wetland and riparian plantings, covercrop for woody plantings, erosion control, and forage.

Cuivre River has not been tested for grazing but forage clippings were taken at different stages of growth and compared to tall fescue clippings from adjacent plots. Forage quality of the Cuivre River selection compared favorably to tall fescue as indicated by data below.

Clipping Date	Percent Protein		Percent ADF		Percent NDF	
	<u>TF</u>	<u>VWR</u>	<u>TF</u>	<u>VWR</u>	<u>TF</u>	<u>VWR</u>
4/24/02		27		26		47
5/30/01	9	12	40	34	61	60
10/11/01	15	15	31	34	52	55
11/15/01	20	17	22	24	37	44

TF = tall fescue; VWR = Cuivre River Virginia wildrye; ADF = acid detergent fiber; NDF = neutral detergent fiber.

2003

One new accession was added during 2003 and this was the medium height, forage type switchgrass that was selected and isolated from the low growing switchgrass assembly.

2004

Three accessions of cluster fescue, *Festuca paradoxa*, were added during 2004. The plants were germinated in the greenhouse from seed and transplanted April 7, 2004, to the initial evaluation area, tier F/a. The accessions established well and had excellent survival the first year. The plants will be evaluated on percent stand, vigor, height, and seed production next year. See collection information below.

Genus	Species	Common Name	Accession No.	Origin
Festuca	paradoxa	Cluster fescue	9083254	Tucker Prairie, MO
Festuca	paradoxa	Cluster fescue	9083255	Paintbrush Prairie, MO
Festuca	paradoxa	Cluster fescue	9083252	Harrison Co, MO

2005

No new accessions were added in 2005. The *Festuca paradoxa* was evaluated along with the other species in the forage quality study. The plants became very dormant by early summer and did not recover but made significant regrowth in the fall.

2006/2008

New accessions planted are as follows;

Observational Nursery

Genus	Species	Common Name	Accn No.	From	Date Pltd
Desmodium	glabellum	Dillenius Tick Trefoil	9055415	MIPMC	5/5/06
Desmodium	glabellum	Dillenius Tick Trefoil	9005087	MIPMC	5/5/06
Desmodium	paniculatum	Panicledleaf Tick Trefoil	9055428	MIPMC	5/5/06
Calamovilfa	longifolia	Prairie Sandreed	9086408	MIPMC	5/5/06
Elymus	riparius	Riverbank Wildrye	9086450	MIPMC	5/5/06
Elymus	canadensis	Icy Blue Canada Wildrye	9084347	MIPMC	5/5/06
Salix	sericea	Riverbend Silky Willow		MIPMC	6/15/06
Paspalum	floridam	Harrison Florida Paspalum	9043874	ETPMC	4/15/06
Bouteloua	gracilis	Blue grama	421782	KSPMC	4/20/07

2009

No new species were planted during 2009.

2010

There were no new plantings of any species' added to this study. A total overhaul of these plots are needed to effectively evaluate the survivability and vigor of each individual species. Many of the species have been sent from other PMC's to evaluate the "area of adaptation". Evaluations of each species' within this study will be taken late 2011.

Study: 29I097G

Study Title: Assembly and Evaluation of Big Bluestem, *Andropogon gerardii* Vitman.

Study Leader: Bruckerhoff, S. B.

Introduction:

Big bluestem is a tall, warm-season, perennial, native grass with stiff, erect culms; flattened and keeled sheaths; membranous ligules; and flat or folded leaf blades. Big bluestem has developed a very efficient spreading root system that may reach depths of 5-8 feet (150-200 cm). Big bluestem reaches a mature height of 3-4 feet (90-120 cm) in northern latitudes, and 6-8 feet (180-240 cm) or more in the southern part of its natural range. Although short rhizomes may be present, it usually makes a bunch type growth. Big bluestem is composed of many ecotypes with a wide range of adaptation to soil and climate. Big bluestem is one of the most widespread and important forage grasses of the North American tallgrass prairie region. It is usually associated with one or more of the other three dominant species, Indiangrass (*Sorghastrum nutans* (L) Nash.), switchgrass (*Panicum virgatum* L.), and little bluestem (*Schizachyrium scoparium* (Michx.) Nash.). Big bluestem occurs on subirrigated lowlands, nearly level to gently undulating glacial till plains, overflow sites, level swales and depressions, residual and glacial uplands, and stream terraces and bottomlands along rivers and tributaries. The abundant, leafy forage is palatable to all classes of livestock.

Problem:

There is a need for an adapted variety of big bluestem for pasture and range seedings, surface mine reclamation, critical area planting, recreational area development and other conservation uses in Arkansas and Southern Missouri.

Objective:

The objective is to assemble, evaluate, develop and cooperatively release an adapted variety and/or varieties of big bluestem for conservation use in the following Major Land Resource Areas: 116A, 116B, 117, 118, and 119.

Cooperators:

USDA-NRCS Plant Materials Center at Elsberry, Missouri and the USDA-NRCS Plant Materials Center at Booneville, Arkansas.

Assembly:

The assembly consists of vegetative materials from adapted ecotypes throughout Northwestern Arkansas and Southwestern Missouri Major Land Resource Areas: 116A, 116B, 117, 118, and 119. Collection dates were between November 9 and 13, 1987. Four collection sites per county within the geographic area of collection were made. The number of sites was determined by the size of the county. The study plan supplement lists the states and the number of sites per county.

Procedure:

Four collections per county in the targeted Major Land Resource Areas were requested. The intent was to get a broad genetic base of plant material; therefore, the site selection attempt was to get as diverse sampling as practical when selecting superior big bluestem plants in the field. If a county had more than one Major Land Resource Area, collections were made in each area. Collections were from typical locations, which included natural grasslands (range), relic areas, and road right-of-ways. Avoided areas were those that may have been artificially seeded. Where possible, collections came from diverse soil textural types, such as sandy and silty; or range site groupings such as: (1) Run-in sites represented by overflow, or subirrigated; (2) normal upland sites represented by sandy, silty or clayey. Six subsamples (6" x 6" x 8" deep) were collected vegetatively at each site.

The samples were transported in material provided by the Plant Materials Center that included cartons, plastic bags, accession data sheets, and instructions for handling.

Plant Materials Center personnel picked up the cartons containing the samples at designated central locations within each administrative area in November 1987.

Transplanting procedures included temporary storage and handling. The samples were first assigned accession numbers and placed in temporary storage. On February 15, 1988, each subsample was transplanted into separate containers and maintained under controlled greenhouse conditions. The plants were then divided between two locations, Elsberry, Missouri and Booneville, Arkansas Plant Materials Centers, and established in space plant initial evaluation nurseries.

Discussion:

1987-1989

A total of 370 accessions (collections) of big bluestem were initially collected during November, 1987, from the targeted areas: 194-Missouri; 85-Arkansas; 82-Oklahoma; and 8-Illinois. Individual plantlets were separated, transplanted into cone-tainers, and grown out in Forrest Keeling Nursery's greenhouse from February until May 1988. More than 4400 individual plantlets were transplanted into a space plant nursery with two replications and six plants per replication. The nursery is located in Field #14 at the PMC and was planted June 1988. The entire nursery was irrigated three times weekly in 1988 to insure good survival. Data collected in 1988 was mostly survival. Data collected in 1989 included survival, vigor, disease resistance, plant size, foliage size, and abundance and visual seed production. Accessions from each state were selected from the above criteria. The numbers selected from each state were as follows: Arkansas-14, Missouri-46, and Oklahoma-13. Table #1 shows the 73 accessions selected from the initial space plant nursery located in Field #14 on the PMC. These plants were vegetatively removed from the initial evaluation nursery in November 1989.

1990-1991

The plants selected in 1989 were transplanted into cone-tainers and grown out in the greenhouse that winter. These plants were planted in an isolated crossing block in Field #1 on May 23, 1990. Fifteen bulk pounds of clean seed were harvested in 1991.

1992-1993

The seed harvested in 1991 was sorted by weight and grown in cone-tainers in the greenhouse from January until April. Approximately 500 plants were planted in Field #7 in April and May 1992 for further evaluation.

Beginning in July 1993, the great flood began flooding approximately 86 acres on the PMC. The area where this planting was located was completely inundated with approximately eight feet of water. Just prior to the flooding of this site (July 8, 1993), the PMC staff uprooted 62 selections of big bluestem and re-established them to an upland site on the PMC (Field #8).

1994-1996

The nursery block established in Field #8 in July 1993 was evaluated for forage quality and quantity, seed production, plant maturity differences, and disease and insect resistance. Twenty-eight of the 62 plants were selected and allowed to cross. Seed from this crossing block is a composite of the original 73 accessions collected and is the breeders' block for the new accession 9078831. Seed was harvested in 1995 and 1996 and a seed increase plot will be established in 1997. The Booneville PMC also has made their selection and both will be included in the advanced evaluation.

1997-1998

The diversity in the original nursery block containing all 370 accessions is tremendous. There is a lot of variation within this species. The need for plant diversity for prairie restoration led to the release of the source-identified composite of all 370 accessions. This composite was given the accession number 9062323 and given the name OH-370 which stands for a composite of 370 collections made from the Ozark Highlands of Southern Missouri, Northern Arkansas, Eastern Oklahoma, and Southern Illinois. This plant was released in April 1997.

A 0.4-acre increase planting of 9078832 was planted May 22, 1997, in Field # 6. This planting was established in a conventional seedbed in 36" rows. The first year the planting produced 10 pounds bulk clean seed and in 1998 it produced 27 pounds bulk clean seed. The 1998 seed tested poorly but it is not known why. When seed becomes available from the Arkansas PMC the study will begin an advanced evaluation to compare the new accession, 9078831 with available varieties and also the accession Booneville has selected out of the original assembly of 370 collections.

The original planting was again evaluated the spring of 1997 looking for a tall, stiff stemmed, upright plant to use in wind barriers. Wind erosion is a problem in the flat and sandy crop fields in the bootheel area of Missouri. Switchgrass windbarriers are being tried in areas where field windbreaks using trees are not acceptable. Big bluestem was requested by the Missouri plant materials committee as an additional species to go along with switchgrass since the nursery is still intact. Five accessions (Table #2) were selected and increased vegetatively in the greenhouse and transplanted into an isolation block in Field #4. This block contained 126 plants and of those, 34 plants were selected to represent the crossing block that will serve as the breeders' block for a wind barrier selection. The final accessions represented in this block are 9056960, 9056913, and 9056914.

Selections were also made for landscape and beautification (Table # 3). These selections were transplanted into the rod row initial evaluation area for further evaluation.

1999

The increase plot of 9078831 was expanded in 1999 but did not develop as the 1997 original increase plot did. This accession is scheduled for release as a pre-varietal selection in 2000 if enough seed is available and field plantings are successful.

The wind barrier selection block was again evaluated in 1999 and narrowed down to a single accession, 9056960 (Table #2).

No additional selections were made for landscape plants in 1999 (Table #3).

2000

The increase plot of 9078831 was again expanded in 2000 but again was very slow to germinate. Seed was sent for testing and the sample contained a high percentage of dormant seed. This pre-varietal selection was scheduled to be released in 2000 and given the name OZ-70 that stands for Ozark Highland composite of 70 collections. The release has been delayed until a solution can be found for its high seed dormancy.

Seed was harvested from the wind barrier block and an increase planting will be made in 2001.

2001

The increase plot of 9078831 (OZ-70) was again expanded in 2001 but this year it was planted the first week of March to allow for stratification. Seed harvested in 2000 was used in the planting because seed less than one year old appears to have more dormancy than seed that has had time in storage. The portion of the plot that was planted in 2001 established well and even produced a small amount of seed the first year.

Seed harvested from the wind barrier accession was propagated in the greenhouse and transplanted into an evaluation nursery. The evaluation nursery has approximately 250 plants on a three-foot grid. These plants will be evaluated for two additional years for height, biomass production and lodging. This plant will be released as a tall, stiff stemmed selection.

2002

Field testing has shown possible problems with establishment of OZ-70 big bluestem. A trial was started using replicated plots to compare the establishment of OZ-70 with 'Rountree' big bluestem. First year data indicates that Rountree establishes quicker with higher stand density than OZ-70. It also indicated that the winter dormant plots (planted March 14, 2002) of OZ-70 were better than the spring planted plots (planted June 21, 2002). This was reversed with the Rountree. This information supports the high seed dormancy problem indicated in seed tests. These plots will be monitored one more year to see if the slow establishment has to do with the long-term density of the plots.

A comparison between new seed and one-year-old seed is planned for 2003. Seed tests indicate a problem with seed dormancy in new seed. Storage for one year could help rectify this problem.

2003

A trial comparing new (previous year's harvest) and older seed (one to five years old) was conducted in 2003. Establishment was quicker if new seed was winter dormant planted. This supports that newly harvested seed has higher seed dormancy but all lots of seed developed into successful stands the establishment year.

The technical review committee recommended proceeding with a Selected Release for this accession and OZ-70 Germplasm Big Bluestem was released December 2003.

Release Documentation

The OZ-70 selection has very good forage production and vigor that appears to be comparable or better than Rountree. OZ-70 is approximately two weeks later in booting than Rountree and forage quality is better when tested at Elsberry (see below). Rountree exhibits considerable more rust when compared to OZ-70 in Southern Missouri. OZ-70 also has very good seed production with a 2003 yield of 280 bulk pounds of clean seed per acre.

Forage clippings of OZ-70 Germplasm were compared with Rountree. These samples were replicated and taken at different stages of growth. Forage quality of the OZ-70 selection compared favorably to Rountree as indicated by following data.

Clipping Date	Percent Protein		Percent ADF		Percent NDF	
	OZ-70	Rountree	OZ-70	Rountree	OZ-70	Rountree
6/19/02	14.3	8	30.9	35.7	55.8	60.8
7/8/02	8.2	5.8	34.1	33.0	59.3	60.5
8/30/02*	11.4	11.9	34.3	34.7	54.6	56.6

*Regrowth material from 7/8/02 clipping.

ADF=acid detergent fiber; NDF=neutral detergent fiber.

OZ-70 Germplasm big bluestem was compared to 'Rountree' big bluestem for establishment and Rountree was quicker to establish indicating better seedling vigor when new (previous year's harvest) seed was planted. A seeding trial was conducted in 2003 and compared seed harvested in 2002, 2001, and a mixture of seed harvested in 1997 through 2000.

The results below indicate some seed dormancy in new crop seed but all plots developed very good to excellent stands and had seedhead production the first year.

	Stems Per Row Foot	Percent Cover
Winter dormant planting, 2002 seed	16	92
Winter dormant planting, 2001 seed	14	78
Winter dormant planting, 97-00 seed	8	65
Spring planting 2002 seed	10	60
Spring planting 2001 seed	14	87
Spring planting 97-00 seed	10	75

2004

The tall, erect, lodging resistant big bluestem currently being evaluated as a wind barrier selection, (accession 9083274) was increased for advanced testing. Seed was harvested in 2003 from the remaining plants in the final evaluation block. The increase block established well but no seed was harvested in 2004. Limited seed production is anticipated for 2005 and available for advanced testing in 2006.

Shorter growing collections were also isolated and evaluated. Six collections were narrowed to three (accessions 9056902, 9056905, and 9056906) and allowed to cross. This composite (accession 9078832) was harvested in 2003 and used to establish an increase block in 2004. Seed production is anticipated for 2005 and available for advanced testing in 2006. This selection will be evaluated for use in vegetative buffers and filters.

2005

The two increase blocks of big bluestem that were established in 2004 (tall, lodging resistant, - accession number 9083274 and shorter growing, - accession 9078832) both produced seed in 2005. These blocks were planted April 28, 2004. Accession 9083274 produced 65.7 bulk pounds on 0.183 acre for a yield of 359 bulk pounds per acre. Accession 9078832 produced 144.9 bulk pounds on 0.51 acre for a yield of 287 bulk pounds per acre.

2006

The two increase blocks of big bluestem planted April 28, 2004 were again managed for seed production. Neither plot was enlarged.

The tall, lodging resistant accession, 9083274, yielded 59.1 bulk pounds on 0.183 acre for a yield of 323 bulk pounds per acre. The year was very dry during parts of the growing season. Seed quality was poor resulting in a very poor percent PLS.

The shorter growing accession, 9078832, yielded 193.2 bulk pounds on 0.51 acre for a yield of 379 bulk pounds per acre. Again seed quality was not very good resulting in a poor percent PLS.

The shorter growing accession (9078832) also is very resistant to lodging and these characteristics look good for this accession's use in conservation programs. With adequate seed on hand for field plantings and grower interest, this accession was released as a selected class release in 2006 as Refuge Germplasm (see 2006 releases section for release notice).

2007

The tall, lodging resistant accession (9083274) increase plot yielded 50.3 bulk pounds on 0.183 acre. The plot was also expanded by a few rows but was slow to establish. Seed from the breeder's block was limited and possibly of poor quality due to a very dry summer. Seed harvested in 2005 from the increase plot was used in the biofuels study plots (MOPMC-T-0716).

The increase field of Refuge Germplasm big bluestem (9078832) has begun showing different heights that indicate contamination of the breeder plot. It was decided to move and isolate the breeder's block and reestablish a new foundation field.

2008

The increase plot of the tall, lodging resistant accession (9083274) of big bluestem was flooded for an extended period of time in 2008 and was destroyed. A new breeders block was established vegetatively and a new increase field is planned for 2010.

The increase plot of Refuge Germplasm big bluestem (9078832) was also destroyed in 2008.

2009

The new breeders block of the tall, lodging resistant accession (9083274) of big bluestem established in 2008 did very well in 2009. Approximately 5% of the 400 plants in the block were rouged before seed development due to lack of desired characteristics. Five pounds of bulk seed was harvested and used to reestablish an increase field. This accession is currently being evaluated in a biofuels study and also an inter center strain trial for big bluestem.

2010

A foundation plot of accession 9083274 was established in field 7A. The plot was harvested in the fall, though the tall stature that this accession was selected for was not evident. This plot will be looked at again in 2011 in order to evaluate this selection for its tall stature and whether or not this characteristic will breed true to the progeny. A possible reason for the 2010 plot not maintaining the tall height (9+ ft.) might be because the planting was seeded in late spring. Seed from this plot will be offered for field testing trials in 2011.

**Study 29I097G - Assembly and Evaluation of Big Bluestem,
Andropogon gerardii, Vitman.**

Table #1

Accessions Selected for Crossing Block

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
Levonna S. Vekman	Arkansas	Faulkner	9056956	118	Leadville
Mark L. Kennedy	Arkansas	Fulton	9056968	116A	Geesville
Luther O. Shaw	Arkansas	Izard	9056920	116A	Mako
NRCS-Field Office	Arkansas	Logan	9056964	118	Taff
NRCS-Field Office	Arkansas	Madison	9056962	118	Leadvale
Stephen T. Ford	Arkansas	Madison	9056945	117	Nixa-SL
John Y. Harrington	Arkansas	Madison	9056923	116A	Estate-SC
John Y. Harrington	Arkansas	Madison	9056952	116A	Estate-SC
Lane L. Gentry	Arkansas	Perry	9056922	119	Clebit
John D. Kopf	Arkansas	Scott	9056936	119	Carnasaw
Jeremy R. Funk	Arkansas	Sharp	9056914	116A	Gepp
NRCS-Field Office	Arkansas	White	9057058	118, 134	
NRCS-Field Office	Arkansas	White	9057060	118,134	
Robert S. Garner	Arkansas	Yell	9056908	119,118	Clebit-FSL
H. Dan Philbrick	Missouri	Barry	9056832	116B	
Dudley W. Kaiser	Missouri	Benton	9056840	116B	Bardley
NRCS-Field Office	Missouri	Camden	9056724	116A	Gatewood
William K. Quage	Missouri	Cedar	9056800	116B	Hector
Patricia A. Beneke	Missouri	Cole	9056821	115	Goutewood
Patricia A. Beneke	Missouri	Cole	9056806	115	Gatewood
Melodie Marshall	Missouri	Crawford	9056820	116B	
Melodie Marshall	Missouri	Crawford	9056886	116B	
Melodie Marshall	Missouri	Crawford	9056767	116B, 116A	Lebanon
Myron C. Hartzell	Missouri	Dent	9056773	116B	Coulstone
Myron C. Hartzell	Missouri	Dent	9056763	116B	Lebanon
John L. Lumb	Missouri	Douglas	9056833	116B	Doniphan
Art Kitchen	Missouri	Franklin	9056855	115	Crider
Art Kitchen	Missouri	Franklin	9065771	115	Union
NRCS-Field Office	Missouri	Gasconade	9056848	116B	Gladden
Clayton P. Robertson	Missouri	Gasconade	9056875	116B	
H. Lane Thurman	Missouri	Greene	9056716	116B	Chirty Silt Loam
NRCS-Field Office	Missouri	Hickory	9056839	116A	
Stanley Lamb	Missouri	Iron	9056774	116A	Midco
Howard Combes	Missouri	Howell	9056753	116A	Doniphan
Joe H. Everett	Missouri	Jefferson	9056842	115	GL
NRCS-Field Office	Missouri	LaClede	9056741	116A	Cherty Silt Loam
Kees VanderMer	Missouri	LaClede	9056791	116A	Union
Cecile Allen	Missouri	Lawrence	9056709	116B	Viraton
Ron R. McMurtrey	Missouri	McDonald	9056719	116A	
Larry E. Lewis	Missouri	Miller	9056732	116B	SIL
Larry E. Lewis	Missouri	Miller	9056868	116B	SIL
Henry E. Knipker	Missouri	Moniteau	9056890	116B	Glensted
Mary Beth Roth	Missouri	Morgan	9056831	116B	

Study 29I097G – Assembly and Evaluation of Big Bluestem, *Andropogon gerardii*, Vitman.

Table #1 - continued

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
Mary Beth Roth	Missouri	Morgan	9056837	116B	
Stephen E. Robbins	Missouri	Organ	9056770	116A	
William R. Dilbeck	Missouri	Polk	9056828	116B	
NRCS-Field Office	Missouri	Pulaski	9056746	116A	Wilderness
Clarence Wagy	Missouri	Reynolds	9056701	116A	
Charles E. Johnson	Missouri	Ripley	9056895	116A	
Charles E. Johnson	Missouri	Ripley	9056894	116A	
Steve Wall	Missouri	Shannon	9056762	116A	
Claude A. Peifer	Missouri	Ste. Genevieve	9056819	116B	Bloomsdale
Edward L. Templeton	Missouri	St. Francois	9056845	116A	Crider
Carl Wehrman and Dude Davidson	Missouri	Taney	9056712	116A	Clarksville
Jeff A. Lamb	Missouri	Texas	9056728	116A	Goss
NRCS-Field Office	Missouri	Wayne	9056854	116A	
Patrick L. Adams	Missouri	Washington	9056817	116A	Silty Clay Loam
Patrick L. Adams	Missouri	Washington	9056870	116A	Silty Clay Loam
John N. Emerson	Missouri	Webster	9056737	116B	
Dan D. Divine	Missouri	Wright	9056733	116B	
Andrew R. Inman	Oklahoma	Adair	9056996	117	Hector Complex
Billy D. Dudley	Oklahoma	Cherokee	9057010	116A, 117	Newtonia
Billy D. Dudley	Oklahoma	Cherokee	9057016	116A, 117	Talpa-Rock
Kenneth W. Swift	Oklahoma	Choctaw	9057025	112	Muskogee SL
Warren R. Sanders	Oklahoma	Coal	9057005	119	Boham
Steve D. Clark	Oklahoma	Latimer	9057014	118, 119	Stigler SL
Robert E. Blackman	Oklahoma	Mayes	9056995	112, 116A	Hector
Sam L. Viles	Oklahoma	McIntosh	9057035	118	Karma SL
Patrick I. Bogart	Oklahoma	Okmulgee	9057032	112, 118	Taloka SL
Patrick I. Bogart	Oklahoma	Okmulgee	9057037	112, 118	Taloka SL
NRCS-Field Office	Oklahoma	Ottawa	9057030	116A, 112	ETA-SL
William R. Bin	Oklahoma	Pushmataho	9957052	119	Bosville
William R. Bin	Oklahoma	Pushmataho	9057046	119	Bernow FSL

Wind Barrier Selection Isolation Block

Table #2

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
	Arkansas	Logan	9056960	118	Laedvale

Study 29I097G – Assembly and Evaluation of Big Bluestem, *Andropogon gerardii*, Vitman.

Landscape Selection Rod Row Area

Table #3

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
Clarence Wagy	Missouri	Carter	9056703	N116A	Opequon
Clarence Wagy	Missouri	Reynolds	9056708	N116A	Clarksville
Myron Hartzell	Missouri	Dent	9056812	116A	Elsah
Kenneth W. Swift	Oklahoma	Latimer	9057025	119	Freestone Variant - Bernow Variant Complex
	Oklahoma	McCurtain	9057049	1336	Kinta Clay Loam
Dennis W. Shirk	Missouri	Maries	9056877	116A	Lebanon
Larry B. Cash	Arkansas	Carroll	9056934	116A	Nixa

Study: **29I108G**

Study Title: Assembly and Evaluation of Low Growing, Rhizomatous Switchgrass, *Panicum virgatum* L. for Use in Waterways, Filter Strips and Other Conservation Uses.

Study Leader: Bruckerhoff, S. B.

Introduction:

Switchgrass is a warm-season, perennial, native grass. Plants are usually green or glaucous, with numerous scaly creeping rhizomes. Culms are erect, tough and hard, one to two meters rarely to three meters tall; sheaths glabrous; blades 10-60 centimeters long, three to 15 millimeters wide, flat glabrous, or sometimes pilose above or near the base, rarely pilose all over; panicle 15-50 centimeters long; acuminate; first glume clasping, two-thirds to three-fourths as long as the spikelet. Switchgrass frequents a wide variety of habitat, usually sunny including dry or moist prairies, moist seepage of rocky glades and buff escarpments, gravel bars of streams, open woods and along railroad tracks.

Problem:

There is a need for an adapted variety of a dense low growing, strongly rhizomatous switchgrass for use in waterways, filter strips, and for other conservation uses in Missouri, Illinois, Iowa, and adjacent states.

Objective:

The objective is to assemble, select, and develop a dense low growing strongly rhizomatous switchgrass, with good seedling vigor and seed characteristics, for use in waterways and streambank corridors.

Procedure:

The assembly consists of the collection of vegetative material from adapted ecotypes in Iowa, Illinois, and Missouri. The targeted collection area includes the following Major Land Resource Areas: 102b, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 131, and 134. Five collections from each NRCS administrative area were requested.

Vegetative collections were taken from natural prairie stands, prairie remnants or individual short growing plants growing in areas that are seasonally wet like a waterway. Total height of the plant was to be no more than three feet.

The samples were collected when the plant was dormant in the fall, divided into plantlets in the winter and placed into square open bottom containers and grown out in the greenhouse. Twelve plants per collection were grown out in the greenhouse.

The plants were planted into a randomized complete block with three replications. Each plot had three plants and all plants were planted on four-foot spacing. A border row was planted around the three replications. This study was planted into a clean tilled seedbed with recommended fertility and weed control. Plants were evaluated for survival, vigor, height, and spread that included rhizomatous characteristics, disease and insect resistance, lodging, and seed production.

Discussion:

1990-1991

The collections of *Panicum virgatum* L., low growing highly rhizomatous switchgrass was initiated in November 1990 and extended through 1991. One hundred eighteen collections were obtained from Major Land Resource Areas 102B-116, 131 and 134 in Missouri, Illinois and Iowa. The total number of collections received was 22-Illinois; 28-Iowa and 68-Missouri. All collections were assigned accession numbers and stored in a cool damp building.

1992-1993

The collections were vegetatively propagated in cone-tainers and placed in the greenhouse in January 1992. These plants were then transplanted in Field #7c on the PMC on June 9, 1992, in a randomized complete block with three replications. Baseline evaluations were taken this year; survival, spread, height, and number of panicles per plant. More detailed evaluations were scheduled for succeeding years.

Beginning in July 1993, the great flood began inundating the area where this project was located. Prior to the flooding of this site (July 2, 1993), additional evaluations were started and 67 accessions were vegetatively moved to an upland site on the PMC for continued evaluation. Table #1 lists the selected accessions, origins, and collectors.

1994-1995

Evaluations were continued on the 67 accessions during 1994 and 1995. The original planting in Field #7c that was flooded in 1993 was also checked for survivors. The planting was flooded by as much as eight feet of water for almost eight weeks. Nine plants were found that showed life and were dug up and moved to an upland site. These nine plants represented three accessions (Table #2).

Five accessions were selected out of the block of 67 for a short growing rhizomatous type. The five accessions (Table #3) were allowed to cross and seed was harvested and grown out in the greenhouse. The five accessions were also dug and increased in the greenhouse in containers.

1996

The five selected accessions (Table #3) were planted into a crossing block June 26, 1996. Half the block was from clonal material from each of the five accessions and the other half was from seed harvested from each of the five plants that were allowed to cross with each other. The accessions of each half of the planting were replicated five times with five plants per replication. Unwanted plants will be eliminated and the remainder of the block will be used for seed increase.

1997-1998

The three accessions (Table #2) of flood tolerant switchgrass were vegetatively increased in the greenhouse. Approximately 250 plants were transplanted April 1997 in Field #7. This is now the breeders' block for the accession 9083170 that is a composite of the three accessions listed in Table #2. Seed was harvested from this plot the first year and used to start a small increase plot in 1998. A small amount of seed was harvested from this increase plot the first year. It is also planned to increase the size of this plot in 1999.

The low growing switchgrass block containing five accessions (Table #3) was again evaluated in 1997. Thirty-five plants were selected from the block of 250. Selected plants were allowed to cross and produce seed. This seed was also used to start an increase field in 1998. This small increase plot produced minimal seed the first year. Seed was again harvested from the 35 plants in 1998 and will be used to make the increase plot size bigger in 1999. The 35 selected plants are the breeder's block for the new accession 9083172 that is a composite of the five accessions in Table #3.

1999

The increase plot of flood tolerant switchgrass, accession 9083170, was expanded in May 1999. This planting did not do well, possibly poor seed germination combined with a very dry summer. Weed control was also poor. Establishment of field plantings was also poor. Expanding the increase plot will again be planned for 2000. Seed was harvested from the breeder's block and the 1998-increase plot. This seed was small due to dry weather.

The increase plot of low growing switchgrass, accession 9083172, was also expanded in May 1999. This planting also did poorly, again possibly poor seed germination combined with a very dry summer. Weed control was also poor. Field testing will begin when seed becomes available. Expanding the increase plot will be planned for year 2000. Seed was harvested from the original 35-plant breeder's block and also the increase field. This seed was also small due to dry weather.

2000

Increase plots of the flood tolerant switchgrass, accession 9083170, and the low growing switchgrass, accession 9083172, were again planted in 2000. These plantings were very sparse and slow to establish. The plantings made in 1999 contained some plants with minimal seed produced. Plantings will again be tried in 2001 with more stratification.

2001

The increase plots of the low growing switchgrass, accession 9083172, that were planted in 1998 and 1999 have filled in and produced seed. The plots planted in 2000 and 2001 have failed. This accession appears to have high seed dormancy and combined with excessive weed competition caused poor establishment. An increase planting is planned for 2002 on an upland site with less weed problems.

The increase plots of the flood tolerant switchgrass, accession 9083170 that was planted in 1998 produced seed in 2001. The 1999 planting was very thin and the 2000 and 2001 plantings have failed. This accession appears to have high seed dormancy. Another increase planting is planned in 2002 with additional stratification.

Accession 9062244 was observed in the nursery block in field eight as having high forage production (very leafy), medium height, and late maturity. Protein analysis of a sample taken was 15.6%. This plant was increased in the greenhouse from vegetative material and planted into a 200-plant nursery in 2000. Unwanted plants were rogued out and seed was harvested in 2001. Plants that germinate quicker from the heaviest seed will be placed in an evaluation nursery in 2002.

2002

The low growing switchgrass, accession 9083172, increase plots had limited seed production in 2002. The 17.3-pound bulk seed produced will be used in the field-planting program for advanced testing. An additional 1.5 acres increase field was planted in 2002. No seed was harvested the establishment year from this plot.

The flood tolerant switchgrass, accession 9083170, increase plots also had limited seed production in 2002. The 32.5-pound bulk seed produced will be used in the field-planting program. Due to an extremely wet spring, no additional seed increase field was planted in 2002.

The medium height forage type switchgrass, accession 9062244, was propagated in the greenhouse and plants were selected for quick establishment and seedling vigor. These plants were transplanted into an evaluation nursery in Field #1 at the PMC.

2003-2004

The low growing switchgrass, accession 9083172, increase plots have been expanded but are slower than expected to develop and produce seed. Available seed is being used in the field planting program for advanced testing.

The flood tolerant switchgrass, accession 9083170, increase plots have been expanded but are also slower than expected to develop and produce seed. Available seed is being used in the field planting program for advanced testing.

The medium height forage type switchgrass, accession 9062244, was again propagated in the greenhouse and the evaluation nursery was expanded in 2003. The plants were allowed to develop and mature in 2004 with evaluations to begin in 2005.

2005

Seed was harvested from the low growing and flood tolerant increase plots. Both of these accessions are being evaluated in the field planting program with mixed results. Seed dormancy is a problem and results in poor and inconsistent establishment.

The medium height switchgrass accession will be placed into a study of its own and go through a recurrent selection process in the development of an improved forage type switchgrass.

2006

Seed was again harvested from the low growing and flood tolerant increase plots. Both are showing poor stand development on heavy soil types with moderate to heavy clay content. These two selections will undergo more testing and selection to improve seedling vigor.

2007

Seed from the low growing and flood tolerant selections were put in the germinator and selected for quick germination, five days or less. Two new evaluation plots were established from the plants selected out of the germinator.

2008

In 2008, plans were to compare germination of seed taken off the selected plots with that of seed taken from the original plots. The data shows the difference between seed that has been stratified and seed that had no stratification period. Within each treatment, low growing, flood tolerant and 'Cave-in-Rock' switchgrass were test for superior germination. The low growing and flood tolerant switchgrass had both SG0 (original) seed and the cycle 1 (selected) seed tested. In the case of the low growing switchgrass, the cycle 1 (selection) made a significant increase in germination. The flood tolerant switchgrass did not appear to have any statistical difference. 'Cave-in-Rock' switchgrass was tested as a comparison.

2009

The two varieties of switchgrass were tested in the fall using seed that was harvested during the 2009 growing season. Comparisons between seed produced in 2008 (Table 1) and seed produced in 2009 (Table 2) are similar in their results. In all cases, except the non-stratified flood tolerant switchgrass from the spring 2009 tests (Table 1), the cycle 1 selections improved in germination. It is worth noting, that the flood tolerant switchgrass, both SG0 and cycle 1 plots, were inundated for approximately 4 weeks just prior to flowering. Also seed from Cave-in-Rock switchgrass for the spring 2009 test (Table 1) was older seed and not from the current growing season. The fall 2009 tests (Table 2) were ran using current growing season seed and is a more accurate test. This test will be ran once more using 2009 growing season seed and if the results show that the cycle 1 selections are still superior, then plans will be to move forward with production plots and field testing.

2010

In 2010, the switchgrass plots on the Elsberry Plant Materials Center were very poor. Germination of seed from all plots were close to “zero” and it was decided not to harvest any switchgrass in 2010. Unfortunately, this holds up the process of proceeding with further evaluations by another year. Plans were to combine 2009 and 2010 seed in order to have an amount that could be used to start a SG1 plot. The plots of the “flood tolerant” and “low growing” selections will be planted in 2011 using 2009 seed and additional years of harvest will be added to these newly established plantings. These new plantings made at the PMC will be evaluated for seedling vigor and the ability to establish quickly. If these plantings do well, plans will be to offer seed from these plots in field plantings and make a comparison evaluation against another variety of switchgrass, such as Cave-in-Rock Switchgrass.

Table 1

Switchgrass Germination Test Spring 2009

	Non-Stratified				Stratified					
	Day 7	Day 14	Day 21	Day 28	% Germ	Day 7	Day 14	Day 21	Day 28	% Germ
Low Growing Switchgrass SG0	9	15	1	1	25	51	5	0	0	56
Low Growing Switchgrass Cycle 1	36	32	0	1	68	78	6	1	0	85
Flood Tolerant Switchgrass SG0	16	10	2	0	27	70	3	0	0	73
Flood Tolerant Switchgrass Cycle 1	15	5	2	1	22	70	5	1	0	76
Cave-in-Rock Switchgrass	37	44	0	0	81	80	5	0	0	85

There were 4 replications of 100 seeds per germination tray using seed harvested in 2008, with exception of Cave-in-Rock switchgrass. C-I-R seed came from a seed lot that was harvested from 1994 through 1997.

Stratified seed (cold/moist) was put in the cooler for 2 weeks.

Non-stratified seed was taken directly from the storage room and entered into the growth chamber.

Table 2

Switchgrass Germination Test Fall 2009

	Non-Stratified				Stratified					
	Day 7	Day 14	Day 21	Day 28	% Germ	Day 7	Day 14	Day 21	Day 28	% Germ
Low Growing Switchgrass SG0	1	6	2	1	10	38	4	0	1	43
Low Growing Switchgrass Cycle 1	1	6	3	2	12	41	7	1	1	49
Flood Tolerant Switchgrass SG0	0	3	0	0	3	48	3	0	0	51
Flood Tolerant Switchgrass Cycle 1	0	5	1	1	6	58	4	1	0	63
Cave-in-Rock Switchgrass	2	3	2	1	7	41	4	0	1	46

There were 4 replications of 100 seeds per germination tray using seed harvested in 2009.

Stratified seed (cold/moist) was put in the cooler for 2 weeks.

Non-stratified seed was taken directly from the storage room and entered into the growth chamber.

Study 29I108G-Selected Accessions of Low Growing Switchgrass

Table #1

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062155	Iowa	Louisa	108	Dean L. Pettit
9062157	Iowa	Cherokee	107	Lon Allan
9062158	Iowa	Clay	103	John P. Vogel
9062160	Iowa	Freemont	107	NRCS F. O.
9062163	Iowa	Hamilton	103	Dana C. Holland
9062165	Iowa	Woodbury	107	John P. Vogel
9062166	Iowa	Monona	107	Michael J. Kuera
9062178	Iowa	Muscatine	108	Douglas S. Johnson
9062181	Illinois	Champaign	108	Leon W. Wendt
9062188	Illinois	Macoupin	108	Ivan N. Dozier
9062189	Illinois	Macoupin	115	Ivan N. Dozier
9062190	Illinois	Macoupin	108	Ivan N. Dozier
9062195	Illinois	Carroll	105	Raymond J. Hudak
9062196	Illinois	Carroll	105	Raymond J. Hudak
9062205	Missouri	Barton	112	Jerry L. Cloyed
9062207	Missouri	Bates	112	Robert D. Bouland
9062208	Missouri	Pettis	116A	Thomas J. Hagedorn
9062209	Missouri	Christian	116A	C. Mark Green
9062211	Missouri	Ozark	116A	Carroll W. Foster
9062212	Missouri	Johnson	112	Robert T. Hagedorn
9062213	Missouri	Madison	116A	Sandra L. Lewis
9062214	Missouri	Ste. Genevieve	116B	Renee L. Phillips
9062215	Missouri	Oregon	116A	Stephen E. Robbins
9062216	Missouri	Shannon	116A	Steve Wall
9062217	Missouri	Reynolds	116A	Clarence W. Wagy
9062218	Missouri	Christian	116A	C. Mark Green
9062219	Missouri	Perry	116B	Claude E. Peifer
9062220	Missouri	Reynolds	116A	Clarence W. Wagy
9062221	Missouri	Dade	116B	Todd E. Mason
9062222	Missouri	Morgan	116B	James A. Maberry
9062223	Missouri	Franklin	116B	Arthur P. Kitchen
9062224	Missouri	Cedar	116B	Kim C. Ehlers
9062225	Missouri	Christian	116A	C. Mark Green
9062227	Missouri	Ozark	116	Carroll W. Foster
9062228	Missouri	Texas	116	Jeff A. Lamb
9062229	Missouri	Texas	116	Jeff A. Lamb
9062234	Missouri	Saline	107	Wayne E. McReynolds
9062237	Missouri	Ray	107	James M. Rehmsmeyer
9062238	Missouri	Worth	109	David A. Stevens
9062239	Missouri	Sullivan	109	Stuart A. Lawson
9062240	Missouri	DeKalb	109	Wm. A. Throckmorton

Table #1 - continued

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062242	Missouri	DeKalb	109	Wm. A. Throckmorton
9062243	Missouri	Buchanan	107	Rodney Saunders
9062244	Missouri	Dent	116	Myron C. Hartzell
9062246	Missouri	Sullivan	109	Stuart A. Lawson
9062247	Missouri	Buchanan	107	Rodney Saunders
9062248	Missouri	Sullivan	109	Stuart A. Lawson
9062250	Missouri	Nodaway	109	Kenton L. Macy
9062251	Missouri	Worth	109	David A. Stevens
9062252	Missouri	Daviess	109	James A. Sturm
9062253	Missouri	Daviess	109	James A. Sturm
9062254	Missouri	Maries	116A	Dennis W. Shirk
9062255	Missouri	Maries	116B	Dennis W. Shirk
9062256	Missouri	Maries	116A	Dennis W. Shirk
9062257	Missouri	Maries	116A	Dennis W. Shirk
9062259	Missouri	Shannon	116A	Steve Wall
9062261	Missouri	Shannon	116A	Steve Wall
9062265	Missouri	Sullivan	109	Stuart A. Lawson
9062267	Missouri	Gentry	109	Gary J. Barker
9062268	Missouri	Platte	107	Terry A. Breyfogle
9062269	Missouri	Sullivan	109	Stuart A. Lawson
9062270	Missouri	Platte	107	Terry D. Breyfogle
9062271	Iowa	Page	104	Kevin J. McCall
9062272	Illinois	Fayette	104	Brad S. Simcox
9062274	Iowa	Madison	108/109	Larry Beeler/Tom Oswald
9062193	Illinois	Fayette	113	Brad S. Simcox

Selected Accessions of Wet Tolerant Switchgrass

Table #2

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062193	Illinois	Fayette	113	Brad S. Simcox
9062213	Missouri	Madison		Sandra L. Lewis
9062235	Missouri	Miller	116	Matt L. Burcham

Final Accessions Selected for Low Growing Switchgrass

Table #3

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062205	Missouri	Barton	112	Jerry L. Cloyed
9062225	Missouri	Christian	116A	C. Mark Green
9062252	Missouri	Daviess	109	James A. Sturm
9062255	Missouri	Maries	116B	Dennis W. Shirk
9062257	Missouri	Maries	116A	Dennis W. Shirk

Study No. 29A116W

Study Title: Evaluation of Miscellaneous Trees and Shrubs.

Study Leader: Cordsiemon, R.

Introduction:

The evaluation of woody plant materials on the USDA-NRCS Elsberry Plant Materials Center began in 1989. Since that time plants have been added for multiple purposes. The evaluations of these plant materials have been in cooperation with the USDA-ARS, Plant Introduction Station, Ames, Iowa; Missouri Department of Conservation; and other plant materials centers.

Problem:

Trees and shrubs are needed to provide for windbreaks, recreation, and multipurpose use in the Midwest Region and provide multiple wildlife benefits throughout the three-state area. New selections, collections and public and private releases need to be evaluated as potential conservation species.

Objective:

The objectives of this study are to assemble and evaluate woody plant materials (both collections in the wild and also released cultivars) for conservation uses, area of adaptation, and to select and increase limited quantities of promising woody plants for advanced evaluation. Superior accessions or those exhibiting unique characteristics will be placed in field evaluations and field plantings in the three-state area being served by the PMC.

Assembly:

Plant materials of various woody species representing many species have been planted on the PMC. The sources include other PMC's, commercial nurseries, and other agencies.

Discussion:**1994-2004**

This study is a long-term ongoing evaluation of miscellaneous trees and shrubs that are not part of a collection made over several years. New species will be planted as they arrive at the Center. Although this study was started in 1989, it includes some species from past studies. Presently there are 29 different species included. Twenty-two are exhibiting 100 percent survival. Five species have failed to survive.

The trees and shrubs in this study are often utilized during plant identification courses held at the Center.

There were no evaluations conducted and no new species added in 2004. There are two new species planned for 2005 that will be received from the Plant Introduction Station in Ames, Iowa. The entire assembly is scheduled to be evaluated in 2005. Very little attention was given to this study in 2004 because the PMC was understaffed.

2005

An evaluation of survival was made in the summer of 2005. Trees and shrubs that had died were noted. The condition of the trees were also evaluated. Black chokeberry (*Aronia melanocarpa*) and common buttonbush (*Cephalanthus occidentalis*) were added to this study. These trees and shrubs will again be evaluated for their survivability and use in conservation.

2006

In April, three new species were added for evaluation, Musclewood (*Carpinus caroliniana*), Bur oak (*Quercus macrocarpa*), and Laurel willow (*Salix pentandra*). There were five trees planted of each species and evaluated for general conditions of the plants (bud break, plant injury, etc.) The buttonbush (*Cephalanthus occidentalis*) and black chokeberry (*Aronia melanocarpa*) were replanted in the fall, 11/14/2005, after dying from an earlier spring planting. They too were evaluated, but for survival, height, spread, injury, type of care given, plant performance, and variations among plants.

2007

Evaluations were taken on tree species sent from the ARS - Plant Introduction Station in Ames, Iowa. Data was sent back to ARS via their online evaluation forms. Also survival of the miscellaneous tree assembly was taken. In August the Three State Technical Review Committee recommended several species of trees to be eliminated from the assembly. The primary species that were recommended for removal were non-native species or species that were performing poorly.

2008

In 2008, the trees and shrubs present in the study were evaluated for survival. Trees from the Plant Introductory Station in Ames, IA were evaluated and data was sent via online evaluation forms. The Introductory Station also sent 2 new species for evaluation, Physocarpus opulifolius (Common Ninebark) and Quercus alba (White Oak). They were planted on 5/2/08. There were no trees eliminated from the evaluation block in 2008. The recommended trees for removal are still scheduled to be removed.

2009

The Miscellaneous Trees and Shrubs study had several older species removed from the plot. Data was taken and the extra space will be prepared for future evaluations and observational plantings.

2010

An area in field 9, north of the pipeline area, was cleared for future observational plantings of tree and shrub material. Three new species were added to the study for evaluation from the North Dakota PMC; Prairie Red Plum, 9047203; Ribes americanum (Black Currant), 9082687; and Photinia melanocarpa (Black Chokeberry), 323957. Eight plants of each were planted and in 2010 all plants were alive. In 2011, the miscellaneous tree study will have a full evaluation for height, spread, insect damage, disease resistance, fruit production, and overall plant vigor.

List of species included in study.

Table #1

Common Name	Genus	Species	Accession Number	Alternate No.	Source	Date Planted
'Densehead' mountain ash	<i>Sorbus</i>	<i>alnifolia</i>		7761	F.K. Nursery	11/65
'Ruby' redosier dogwood	<i>Cornus</i>	<i>stolonifera</i>	443229		Big Flats PMC	5/89
Late lilac	<i>Syringa</i>	<i>villosa</i>	9006228		Bismarck PMC	5/89
'Redstone' cornelian cherry dogwood	<i>Cornus</i>	<i>mas</i>	9055585		Elsberry PMC	5/89
'Roselow' sargent crabapple	<i>Malus</i>	<i>sargentii</i>	477986		Roselake PMC	5/89
'Elsmo' lacebark elm	<i>Ulmus</i>	<i>parvifolia</i>	9004438		Asia	5/89
Blueleaf honeysuckle	<i>Lonicera</i>	<i>korolkowi</i>	9062152		Nebraska	5/89
Birch	<i>Betula</i>	<i>species</i>	502295		Ames, IA	4/90
Willow oak	<i>Quercus</i>	<i>phellos</i>		4723	Ames, IA	4/90
Fragrant epaulettetree	<i>Pterostyrax</i>	<i>hispida</i>		A80779	Ames, IA	4/90
Bradford pear	<i>pyrus</i>	<i>calleryana</i>		19173	Ames, IA	4/69
Prairie rose	<i>Rosa</i>	<i>setigera</i>	495616		Ames, IA	4/90
Ural false spirea	<i>Sorbaria</i>	<i>sorbifolia</i>		7778	Ames, IA	4/90
Weeping lilac	<i>Syringa</i>	<i>pekinensis</i>	478008		Ames, IA	4/90
Flameleaf sumac	<i>Rhus</i>	<i>copallina</i>		7764	Ames, IA	4/90
Western paper birch	<i>Betula</i>	<i>occidentalis</i>	495882		Ames, IA	4/90
Amur honeysuckle	<i>Lonicera</i>	<i>mackii</i>	477998		Ames, IA	4/90
Mountain ash	<i>Sorbus</i>	<i>reducta</i>		A-8371	Ames, IA	4/90
Blackhaw	<i>Viburnum</i>	<i>prunifolium</i>		2813	Ames, IA	4/90
Largeleaf dogwood	<i>Cornus</i>	<i>macrophylla</i>		10178	Ames, IA	4/90
Border privet	<i>Ligustrum</i>	<i>obtusifolium</i>	477010		Ames, IA	4/90
Willow oak	<i>Quercus</i>	<i>phellos</i>		4724	Ames, IA	4/90
Arrowwood	<i>Viburnum</i>	<i>dentatum</i>			Elsberry, MO	4/90

<u>Common Name</u>	<u>Genus</u>	<u>Species</u>	<u>Accession Number</u>	<u>Alternate No.</u>	<u>Source</u>	<u>Date Planted</u>
Redbud	<i>Cercis</i>	<i>canadensis</i>	496399		Ames, IA	5/91
Birch	<i>Betula</i>	<i>species</i>	14942		Ames, IA	5/91
'Wichita' osage orange	<i>Maclura</i>	<i>pomifera</i>			Kansas	5/91
'Denmark' osage orange	<i>Maclura</i>	<i>pomifera</i>			Denmark, IA	6/92
Magenta	<i>Malus</i>	<i>species</i>	514275		Roselake PMC	4/93
Ocean view beach plum	<i>Prunus</i>	<i>maritima</i>	518824		Cape May PMC	5/93
'Sandy' rugosa rose	<i>Rosa</i>	<i>rugosa</i>			Cape May PMC	5/93
Wildwood bayberry	<i>Myrica</i>	<i>pennsylvanica</i>	548966		Cape May PMC	5/93
Wildwood bayberry	<i>Myrica</i>	<i>pennsylvanica</i>	434150		Cape May PMC	5/93
Wildwood bayberry	<i>Myrica</i>	<i>pennsylvanica</i>	548964		Cape May PMC	5/93
Ocean view beach plum	<i>Prunus</i>	<i>maritima</i>	518822		Cape May PMC	5/93
Ocean view beach plum	<i>Prunus</i>	<i>maritima</i>	518823		Cape May PMC	5/93
'Oahe' hackberry	<i>Celtis</i>	<i>occidentalis</i>	476982		Bismarck PMC	5/93
'King Red' Russian olive	<i>Elaeagnus</i>	<i>angustifolia</i>	434029		NPMC	5/93
Black Chokeberry	<i>Aronia</i>	<i>melanocarpa</i>	9083269	Ames 27371	Ames, IA	11/05
Common Buttonbush	<i>Cephalanthus</i>	<i>occidentalis</i>	9083270	Ames 27336	Ames, IA	11/05
Musclewood	<i>Carpinus</i>	<i>caroliniana</i>	9083283	Ames 27963	Ames, IA	04/06
Bur Oak	<i>Quercus</i>	<i>macrocarpa</i>	9004392	Ames 26202	Ames, IA	04/06
Laurel Willow	<i>Salix</i>	<i>pentandra</i>	9083284	Ames 27971	Ames, IA	04/06
Common Ninebark	<i>Physocarpus</i>	<i>opulifolius</i>	9083321	Ames 27970	Ames, IA	05/08
White Oak	<i>Quercus</i>	<i>alba</i>	9083322	Ames 27340	Ames, IA	05/08
Prairie Red Plum	<i>Prunus</i>	<i>spp.</i>	9047203		NDPMC	05/09
Black Currant	<i>Ribes</i>	<i>americum</i>	9082687		NDPMC	05/09
Black Chokeberry	<i>Photinic</i>	<i>melanocarpa</i>	323957		NDPMC	05/09

Study: 29I124G

Study Title: Production of Native Iowa Ecotypes of Grasses and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Grasses and Forbs are Now Being Planted.

Study Leader: Cordsiemon, R.

Introduction:

Well-adapted native grass, legume, and forb plantings offer many advantages as low cost sustainable vegetative cover for management of soil and water resources. Native plant communities resist noxious weed invasion, provide excellent erosion control, and generally require relatively low maintenance.

These characteristics make them an excellent selection for use in roadside plantings, critical areas, long term land retirement programs, and all other vegetative plantings where monocultures of native grasses are being planted. This is especially true along public transportation right-of-ways. These transportation corridors constitute a major land resource and management problem in the state of Iowa. Based on 1987 Natural Resources Inventory (NRI) data, over one million acres of Iowa land are devoted to rural transportation.

Proper vegetation management along these corridors is an important element in controlling soil loss and unwanted weedy plant species. Many of these acres are now seeded to introduced cool-season grass and legume species which are often invaded by noxious weeds requiring extensive mowing or herbicide treatment programs. These management techniques are expensive and can also result in additional water quality problems where herbicides are used extensively.

Managing or re-seeding these acres to promote native grasses, legumes, and forbs offers a low cost environmentally sound approach to roadside vegetation management. Herbicide use, soil erosion, and most mowing can be reduced significantly where a vigorous native grass, legume, and forb mixture dominates a roadside right-of-way. In addition, these goals are consistent with on-going NRCS programs designed to improve ground and surface water quality, reduce soil loss and increase wildlife habitat.

Problem:

Many adapted native species are either currently not commercially available or available only in very limited quantities. When native species are available, the origin is often from considerable distance away and adaptation can be a concern. The species that are available are often as a 'variety' that has been developed for pasture and hay. These are generally high forage producing and more vigorous than wild collections of seed that have not been through an evaluation and breeding program. Seed of local origin that have not been improved or selected for superior forage yield is more likely to remain in a prairie mixture without crowding out other species and becoming monoculture. There is a need for additional native grass, legume, and forb species for use in roadside and other types of conservation plantings.

Objective:

The objective of this study is to accelerate the collection and increase of selected native grass, legume, and forb species through a cooperative program between the University of Northern Iowa (UNI), USDA Natural Resources Conservation Service (NRCS), and the Iowa Roadside Integrated Vegetation Management Program (IRVM).

Cooperators:

The USDA Natural Resources Conservation Service, Plant Materials Center; the University of Northern Iowa; and the Integrated Roadside Vegetation Management Office.

Procedures:

The state of Iowa was divided into three zones: North, Central, and South (Table #1). Seed collected from within each zone was kept separate from the other zones. The IRVM office organized seed collections from each zone. Collections were made from native prairie remnants throughout each zone striving for a relatively equal and representative collection. Seed from each collection site was inventoried by location and a small portion was started in the greenhouse at UNI and transplanted into plots. The remainder of the seed was sent to the PMC, cleaned, and seeded for increase plots. Seed from the plots at UNI was hand harvested and also used to start increase plots or mixed with additional seed and became available to seed growers. When enough seed becomes available, the species is released as 'Source Identified' germplasm from the zone in which it was collected. Source identified seed has not been improved by evaluation and selection or plant breeding procedures.

Discussion:

The study officially started October 1, 1990, at the beginning of fiscal year 1991 with agreements signed. Seed collections had started earlier in the year and seed was available for increase plots the spring of 1991. Most of the plots started from 1991 to 1993 were destroyed in the flood the summer of 1993. Plant re-establishment started in 1994 and new plots have been started each year.

2000

New increase plots established in 2000 were *Liatris asper*, rough blazing star; *Monarda fistulosa*, horsemint; and *Lobelia siphilitica*, great blue lobelia. Surflan was used for weed control and the horsemint was not resistant.

New plant releases for 2000 were Northern Iowa Germplasm Big Bluestem, Northern Iowa Germplasm Tall Dropseed, Northern Iowa Germplasm Roundhead Lespedeza, and Southern Iowa Germplasm Prairie Blazing Star.

2001

There were no new plant releases through the plant materials program in 2001 but seed of previous releases was allocated to growers. Initial seed increase is now in production at the new UNI Native Roadside Vegetation Center at the University of Northern Iowa, Cedar Falls, Iowa. A new plot of Southern Iowa June grass was established at the PMC from plants started in the greenhouse. This species exhibits very slow growth and a serious problem is weed control.

2002

There were no new increase plots established in 2002. Seed production and allocation to growers continued on previously established plots.

New plant releases for 2002 were Northern, Central and Southern Iowa Germplasm New England Aster, Northern and Southern Iowa Germplasm Pale Purple Coneflower, Southern and Central Iowa Germplasm Rigid Goldenrod, and Southern Iowa Germplasm Tall Dropseed.

2003

In 2003 there were no new plantings or increases added. Production and allocations to growers continued from previously established plots. Weed control was maintained by using a non-selective herbicide in late winter/early spring on most plots, followed by a pre-emergent herbicide on all plots. Late spring and summer weed control was achieved by manual labor and selective herbicides.

There were eight new plant releases for 2003. They were Southern Iowa Germplasm Wild Burgamot (*Monarda fistulosa*), Northern, Central, and Southern Iowa Germplasm Rough Blazing Star (*Liatris aspera*), Northern Iowa Germplasm Purple Prairie Clover (*Dalea purpurea*), Central Iowa Germplasm Switchgrass (*Panicum virgatum*), Northern and Central Iowa Germplasm Junegrass (*Koeleria macanthra*). Refer to the table of contents for a complete list of PMC releases.

2004

The Iowa Ecotype Program continued to produce seed for the three different zones on the center in 2004. Although there were no new plots established and no plot increases, the PMC plans to introduce five new Iowa releases in 2005. Weed control was very similar to that of 2003, with the use of non-selective herbicide early and manual labor and selective herbicide later in the growing season. There were some plots taken out of production in 2004 because of consistently low seed production.

2005

The releases scheduled for 2005 were held off until 2006 because there was a lack of available seed. Plots were maintained the same as the past two years. Unproductive plots that had a supply of seed on inventory were mowed and not maintained or harvested.

2006

In fiscal year 2006 the PMC released Central Iowa Germplasm Pale Purple Coneflower (*Echinacea pallida*), 9068612. Future releases are still planned. Each species that the PMC is working with should have a release from each of the three zones. Those releases that are not represented will be a priority for the next few years. Fiscal year 2007 is scheduled to have two releases, northern and central zones of wild bergamot, (*Monarda fistulosa*). In fiscal year 2008 and 2009 the PMC will finish out the Iowa Ecotype Program with the releases of southern zone purple prairie clover, (*Dalea purpurea*), southern zone Junegrass, (*Koeleria macanthra*), and northern and southern zones of switchgrass, (*Panicum virgatum*). The Elsberry PMC and the University of Northern Iowa cooperatively grow and have seed on hand for commercial production.

2007

The Iowa Ecotype Program had two more releases in 2007 with the release of Northern and Central Iowa Germplasm horsemint (also known as wild bergamot), *Monarda fistulosa*. There are limited quantities of seed from both zones currently available and being produced at the Elsberry PMC. More production plots are being phased out as the Tallgrass Prairie Center in Iowa continues to grow and manage more production plots. Plots at the Elsberry PMC that have been taken out of production are being mowed until needed for other uses or the plot needs to be re-established.

2008

In 2008, Iowa ecotype plots, primarily from southern collections, were maintained and harvested for seed production.

2009

Few plots are left in production from the Iowa Ecotype Program. This past growing season only four plots were harvested for seed production. The southern zones for little bluestem and pale purple coneflower and central zones of little bluestem and prairie blazing star produced small amounts of seed and will be added to inventory. Emphasis on this program has fallen off and the PMC will continue to harvest readily available seed and hold on inventory, if there is ever a need to re-establish these stands. The Tallgrass Prairie Center, located on the campus of the University of Northern Iowa, is currently maintaining the releases from this program and has seed available to potential growers.

2010


In 2010, the Elsberry PMC harvested seed from Southern Iowa Germplasm big bluestem, Central Iowa Germplasm little bluestem, and Northern Iowa Germplasm and Southern Iowa Germplasm Indiangrass. This will likely be the final year that the Elsberry PMC harvest from the Iowa Ecotype Program plots, unless plots need to be re-established for future production. The Tallgrass Prairie Center will maintain the established releases and the Elsberry PMC will secure production information from them annually.

Ecotype Species		Iowa Geographic Zones - Year of Release		
Common Name	Scientific Name	Northern	Central	Southern
Pale Purple Coneflower	<i>Echinacea pallida</i>	2002	2006	2002
Purple Prairie Clover	<i>Dalea purpurea</i>	2003	1998	TBD
Switchgrass	<i>Panicum virgatum</i>	TBD	2003	TBD
Junegrass	<i>Koeleria macanthra</i>	2003	2003	TBD
Horsemint	<i>Monarda fistulosa</i>	2007	2007	2003
Rough Blazing Star	<i>Liatris aspera</i>	2003	2003	2003
New England Aster	<i>Aster novae-angliae</i>	2002	2002	2002
Tall Dropseed	<i>Sporobolus compositus</i>	2002	1996	2002
Stiff Goldenrod	<i>Oligoneuron rigidum</i>	1998	2002	2002
Big Bluestem	<i>Andropogon gerardii</i>	2000	1998	2000
Prairie Blazing Star	<i>Liatris pycnostachya</i>	1999	1999	2000
Bushclover	<i>Lespedeza capitata</i>	2000	1996	1997
Little Bluestem	<i>Schizachrium scoparium</i>	1999	1997	1999
Rattlesnake Master	<i>Eryngium yuccifolium</i>	1998	1999	1999
Indiangrass	<i>Sorghastrum nutans</i>	1996	1996	1998
Canada Wild Rye	<i>Elymus canadensis</i>	1995	1995	1997
Oxeye False Sunflower	<i>Heliopsis helianthoides</i>	1996	1995	1997
Sideoats gramma	<i>Bouteloua curtipendula</i>	1995	1995	1995

Study: 29I124G – Native Iowa Ecotypes

TABLE #1

IOWA ECOTYPE ZONE MAP



Study: 29A1370

Study Title: Wetland/Riparian Propagation, Establishment, and Demonstration

Study Leader: Cordsiemon, R.; J. Kaiser

Introduction:

There is a growing interest in wetland restoration throughout the conservation community. Government programs, such as USDA-Wetland Reserve Program, the USFWS Partners for Wildlife, Wetland Restoration Program, the Missouri Department of Conservation (MDC) Private Lands Wetland Program, and private programs sponsored by Ducks Unlimited and Waterfowl USA have all focused on the need for a suitable supply of plants in wetland restoration efforts.

The increasing use of wetlands as filters in agricultural waste management and the control of non-point source pollution also indicate the need for a greater knowledge base for proper plant selection.

Understanding wetland ecosystems will require improved and increased quality of information on wetland plants and ecosystems. Innovative approaches to field management and additional training of personnel in wetland conservation and management will also be needed. Intra- and interagency coordination and information exchange among state and federal agencies will help standardize monitoring and management strategies.

Problem:

Information is largely unavailable related to the propagation, adaptation, and potential use of many of the wetland species found in the Midwest. Wetland plants of interest often have multi-use potential providing wildlife benefits, shoreline stabilization, water quality improvement, and/or aesthetic benefits. They are also needed to fulfill conservation needs resulting from increased demands in wetland development and water treatment. The ability to document this information or to observe the interaction of selected species is restricted by the availability of plants and plant communities especially under controlled conditions. Proper use of species to address conservation problems is limited by specific knowledge and technology for using these plants.

Objectives:

The objectives of the Elsberry PMC wetland study are to:

1. Provide a demonstration of various plant materials for wetland conservation and aesthetic values.
2. Provide an area for interagency research on the biology of selected wetland plants.

Discussion:**1994 – 1999**

A large wetland was constructed in Field #4 on the Plant Materials Center in July 1994. Selected plant materials were planted with the intent of evaluating these plants for flood tolerance. The PMC has been working with a flood tolerant switchgrass since 1991. As a result, it was placed in this wetland for further testing along with six accessions of eastern gamagrass which were found

growing in wet conditions: accessions 9078842, 9078844 and 9078843 were collected in Atchison County, Missouri, 9078845 collected in Holt County, Missouri, 9078840 collected in Chariton County, Missouri and 9078846 was collected in Clinton County, Missouri. Local collections of bermudagrass and swamp milkweed were planted in the spring of 1998. Two collections of prairie cordgrass (Cuivre Island and Lost Creek) were also planted in this wetland. The switchgrass, eastern gamagrass and the prairie cordgrass were planted in 1997. All plants in this wetland were given time to establish prior to the beginning of the flooding operation which took place in October 1999. The wetland was flooded to a depth of 40 inches. This water remained in the wetland until early spring of 2000. Once the water is drained out of the wetland and enough time elapses for plant regrowth, evaluations on survival will take place.

The following Tables #1, #2, #3 and #4 reflect the plants' performance.

2000

Water was drained out of the wetland in segments because the drainpipe was not functioning properly. This operation started on March 21, 2000 and ended on March 30, 2000. The prairie cordgrass were the first plants to begin green up (March 30) followed by the bermudagrass planting. 'Cave-In-Rock' switchgrass sod (23 plugs) was planted on the west side of the flood tolerant switchgrass (sod) for comparison with other plant species in the wetland. On June 1, 2000, flood tolerant switchgrass was seeded in a plot 50 feet long and three feet wide. On August 9 an evaluation of the seeded flood tolerant switchgrass revealed no germination had taken place in the plot seeded on June 1. Poor germination has been experienced with this selection since 1998. There was no flooding of the wetland this fall to allow the Cave-In-Rock to get fully established. The following is a listing of percent survival of plants included in this study. The best performing plants in this study are Cuivre Island and Lost Creek collection of *Spartina pectinata*, *Tripsacum dactyloides* accessions 9078843, 9078845, and 'Pete'; and *Cynodon dactylon*. The following tables reflect the different plants' performance before and after a flooding event.

2001

The objective of the flooding was to parallel flood events that were occurring on the Mississippi River during that same time event. Began pumping turbid water into wetland on April 24, 2001 to flood the wetland to a depth of approximately 32 inches of water, which was achieved by April 27, 2001. The water was allowed to remain in the wetland for seven days. Water was then allowed to drain out of the wetland starting on April 30, 2001. All the water was drained out of the wetland by May 1, 2001. On May 8 evaluations were conducted to document re-growth after flooding. Again on June 11 a quick flooding scenario was conducted in the wetland to simulate a flash flooding event, similar to what was occurring on the Mississippi River. Thirty-four inches of turbid water was pumped into the wetland. The PMC began draining the water out of the wetland on June 15. The process of draining the water out of the wetland was completed on June 19.

The following is a listing of plant vigor ratings for each accession/variety included in this study. Plant evaluations for vigor were taken on June 21 and 26, 2001.

2002-2003

Plant performance evaluations were performed on April 24, 2002 and May 27, 2003. The wetland was not burned in 2002; however it was burned in 2003 and in previous years to remove accumulated vegetation. Flooding of the wetland began on April 29, 2002 and June 10, 2003. A total of 45 inches of water was pumped into the wetland (2002) and 42 inches in 2003 before the de-watering process began. All water was drained out of the wetland by May 17, 2002 and July 7, 2003. The plants were under water for 17 days in 2002 and 22 days in 2003. Once all the water was drained out of the wetland, follow-up evaluations took place on June 2002 and August 2003. The flood event in 2003 was to inundate the site for more than 20 days to test the switchgrass, *Panicum virgatum*. Table #4 reflects the plant performances during 2003 before and after the flood event. Previous years' plant performances can be found on Tables #1 - #3.

2003-2004

Switchgrass, *Panicum virgatum*, accessions 9062193, 9062235, 9083170 were compared to Cave-In-Rock. The percent was 76%, 77%, and 78% survival compared to Cave-In-Rock at 65%. The composite 9083170 Flood Tolerant switchgrass is the next generation of the three accessions 9062193, 9062235, and 9083170 which did perform from seed that was planted in 2000. Vigor was slow with only 20% stand the first growing season. Flood events occurred in 2001, 2002, and 2003 with the stand increasing in density to 85% by spring of 2004.

Prairie cordgrass, *Spartina pectina*, accessions 9083166 Cuivre Island and 9083167 Lost Creek planted on the 3'X 3' grid was a solid block in two growing seasons. The cordgrass planted on the 10'x 10' grid was a solid block in six growing seasons. The vegetative spread averaged 1.5 feet during a growing season. The flooding events did enhance the plants' ability to flourish and produce seed that spread seedlings in the wetland cell.

Virginia wildrye, *Elymus virginicus*, accession 9083169 Cuivre River was vegetatively transplanted in 2001. In the flood event of early spring 2002 there was 100% survival of the plants; however the flood event of 2003 late spring to early summer did result in a decline in the plants with 47% survival by spring 2004. Many seedlings were observed that came from seed in the soil that developed fall 2003 and spring 2004.

2004

The wetland cell was not burned and there was no flooding in 2004. A new block was added to the wetland for evaluation. The block contained 16 plants of low growing switchgrass, *Panicum virgatum*, erect big bluestem, and short growing big bluestem, *Andropogon gerardii*. 'Cave-In-Rock' switchgrass was added to the block as a check. The plants were transplanted from plugs grown in the greenhouse in order to get good established plants. They were evaluated for survival in October and only the low growing and 'Cave-in-Rock' switchgrasses were needed; three and four plants respectfully. The block will be flooded in late April to June of 2005 and the entire wetland planting will be evaluated.

2005

In 2005 there was no activity with this study. The warm season grasses, erect big bluestem, 9083274 and short-growing known as Refuge, 9078832, and low growing switchgrass, 9083172, and Cave-in-Rock, 469228, were allowed to establish. Then control of broadleaf weeds was addressed. The flooding sequences are again planned for 2006, but with lack of help may be put off indefinitely.

2006

A re-evaluation of this study was done and a determination was made to evaluate the warm season grasses and other plants in the wetland cell if time and labor is available. A survival evaluation was done on the warm season grasses (bluestems and switchgrasses). Those plants that have died were replaced.

2007

The wetland cell was mowed and the eastern side of the cell has been cleaned and made available for new specie evaluations. There were no flooding sequences in 2007 and plants that are in the wetland cell have been maintained. Plans are to flood the cell in 2008 and simulate the flooding sequence of the Mississippi River in 2007. Evaluations will be taken for survival and regrowth after the flooding event. Potentially new flood tolerant species may be added to the study in 2008.

2008

The warm season grasses, erect (commonly called Epic) big bluestem, 9083274 and Refuge, 9078832, and low growing switchgrass, 9083172, and Cave-in-Rock, 469228, were flooded in April as the plants began to emerge. The height of each plant was between 1/2 to 3 inches. The big bluestems did not progress and the switchgrasses were slow, showing very little growth. After 14 days of inundation the water was dropped and the switchgrasses flourished. Both big bluestems were very slow to recover and did not become full plants as they had in 2007. They were very spindly. The switchgrass plants performed very well after the water was drained. Based on observations from this study, the selected big bluestem accessions would not be a beneficial plant for use in frequently flooded or wetland situations. Refer to Table #5 for plant performance.

2009

There were no evaluations performed in this study in 2009. Information for several different sedges in frequently flooded areas is of growing interest. Different sedge species from the ongoing sedge study could be a possibility for evaluation. This study will stay active on an “as needed” basis.

2010

There were no evaluations performed for this study in 2010. This study will stay active on an “as needed” basis.

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #1
Plugs Planted 5-2-97 (Eastern Gamagrass)							
2002 Data	Began Flooding on 4/24/02						
2003 Data	Began Flooding on 6-10-03						
	Total #	Active	Weed	Disease/	Developed	Vigor	Ave. Ht.
	Planted	Growing	Comp.	Insect	Seed Head		
Eastern Gamagrass 9078840 Chariton, Missouri. 5' spacing, planted 5/2/97.							
							25 plants planted
Dates Evaluated							
7/9/1998	20	20	severe	moderate	yes	good	2'5"
9/29/1999	20	20	moderate	light rust	yes	good/exc	3'5"
5/11/2000	19	17	moderate	moderate	none	poor	6"
9/19/2000	13	13	mod/sev	light rust	none	good	2'5"
6/26/2001	20	20	light	none	yes	good	3'4"
4/24/02 (BFE)	18	18	light	none	none	good	8"
6/17/02 (AFE)	15	15	light	none	yes	exc	2'
5/27/03 (BFE)	15	15	light	none	yes	exc.	2'
8/5/03 (AFE)	*						
Percent surviving as of 6/17/02 was 75%							
Eastern Gamagrass 9078844 Atchison, Missouri. 7' spacing, planted 5/2/97.							
							18 plants planted
Dates Evaluated							
7/9/1998	12	12	severe	moderate rust	yes	poor	2'5"
9/29/1999	12	12	moderate	moderate rust	yes	fair	2'5"
5/11/2000	12	10	moderate	moderate	none	poor	6"
9/19/2000	12	13	severe	light rust	Yes	fair	2'
6/26/2001	12	9	light	light rust	yes	fair	2'10"
4/24/02 (BFE)	9	9	light	none	none	fair	7"
6/17/02 (AFE)	9	9	light	none	none	exc.	2'
5/27/03 (BFE)	*						
8/5/03 (AFE)	*						
Percent surviving as of 6/17/02 was 75%							
Eastern Gamagrass 9078842 Atchison, Missouri. 15' spacing, planted 5/2/97.							
							9 plants planted
Dates Evaluated							
7/9/1998	5	5	severe	none	yes	fair	2'
9/29/1999	5	5	severe	none	yes	fair	2'5"
5/11/2000	5	3		none		poor	6"
9/19/2000	5	4	severe	none	none	fair	1'8"
6/26/2001	3	3	light	none	yes	fair	2'2"
4/24/02 (BFE)	4	4	light	none	none	fair	7"
6/17/02 (AFE)	4	4	light	none	none	exc.	2'
5/27/03 (BFE)	*						
8/5/03 (AFE)	*						
Percent surviving as of 6/17/02 was 44%							
Rating for Vigor: 1=Excellent; 9=Poor							
Rating for Weed Competition and Dis/Insect: 1=Excellent; 9=Severe							
* = Cannot determine rows of plants							

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #1-continued	
	Total # Planted	Active Growing	Weed Comp.	Disease/ Insect	Developed Seed Head	Vigor	Ave. Ht.	
Eastern Gamagrass 9078846 Clinton, Missouri. 8' spacing, total planted 5/2/97.								
							16 plants planted	
Dates Evaluated								
7/9/1998	11	11	severe	none	yes	good	2'	
9/29/1999	11	11	moderate	none	yes	good	2'5"	
5/11/2000	8	8	moderate	none	none	poor	7"	
9/19/2000	10	10	severe	light rust	none	fair	2'	
6/26/2001	8	8	light	light rust	yes	good	3'2"	
4/24/02 (BFE)	10	10	light	none	none	good	8"	
6/17/02 (AFE)	10	10	light	none	yes	exc.	2'6"	
5/27/03 (BFE)	*							
8/5/03 (AFE)	*							
Percent surviving as of 6/17/02 was 63%								
Eastern Gamagrass 9078843 Atchison, Missouri. 15' spacing, planted 5/2/97.								
							9 plants planted	
Dates Evaluated								
7/9/1998	13	13	severe	none	yes	poor	2'5"	
9/29/1999	13	13	moderate	none	yes	moderate	3'	
5/11/2000	5	5		none	none	poor	7"	
9/19/2000	10	10	severe	slight rust	none	fair	2'	
6/26/2001	4	4	light	light	none	fair	2'6"	
4/24/02 (BFE)	4	4	light	light	none	fair	8"	
6/17/02 (AFE)	4	4	light	light	none	good	2'	
5/27/03 (BFE)	*							
8/5/03 (AFE)	*							
Percent surviving as of 6/17/02 was 44%								
Eastern Gamagrass 9078845 Holt, Missouri. 8' spacing, planted 5/2/97.								
							16 plants planted	
Dates Evaluated								
7/9/1998	12	12	severe	none	yes	good	3'5"	
9/29/1999	12	12	severe	none	yes	good	3'	
5/22/2000	12	9	severe	none	none		8"	
9/19/2000	16	16	severe	slight rust	yes	good	2'5"	
6/26/2001	10	10	light	none	yes	good	3'2"	
4/24/02 (BFE)	10	10	light	none	none	good	8"	
6/17/02 (AFE)	10	10	light	none	none	exc.	2'6"	
5/27/03 (BFE)	*							
8/5/03 (AFE)	*							
Percent surviving as of 6/17/02 was 63%								
Rating for Vigor: 1=Excellent; 9=Poor								
Rating for Weed Competition and Dis/Insect: 1=Excellent; 9=Severe								
BFE - Before Flooding Event								
AFE - After Flooding Event								
* = Cannot determine rows of plants								

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #2
Plugs Planted 6-24-97 (Flood Tolerant Switchgrass)							
2002 Data: Flood Event from 4/29/02 to 5/17/02							
2003 Data: Flooding began 6/10/03							
	% Cover/ Plant #	Active Growing	Weed Comp.	Disease/ Insect	Developed Seed Head	Vigor	Ave. Ht.
Switchgrass 9062213 3' spacing, 41 total planted (plugs) 6/24/97.							
Dates Evaluated							
7/9/1998		35 plants	moderate	none	all plants	poor/fair	2'
9/29/1999		35 plants	moderate	none	all plants	fair	2' 5"
4/26/2000		35 plants	moderate	none	none	exc.	5" regrowth
9/19/2000	85% row	35 plants	moderate	none	all plants	exc.	4'5"
6/26/2001		33 plants	light	none	none	exc.	3' 4"
4/24/02 (BFE)		31 plants	light	none	none	good	6"
6/17/02 (AFE)		31 plants	light	none	none	exc.	2' 6"
5/27/03 (BFE)		32 plants	light	none	none	exc.	1' 7"
8/5/03 (AFE)		32 plants	light	none	none	good	2' 5"
Percent surviving as of 6/17/02 was 76%							
Switchgrass 9062235 4' spacing, 31 total planted (plugs) 6/24/97.							
Dates Evaluated							
7/9/1998		22 plants	moderate	none	all plants	poor/fair	5' 5"
9/29/1999		22 plants	moderate	none	all plants	fair	5'
4/26/2000		26 plants	moderate	none	none	exc.	6' 5"
9/19/2000		26 plants	moderate	none	All plants	exc.	4' 5"
6/26/2001		24 plants	light	none	none	exc.	2' 9"
4/24/02 (BFE)		20 plants	light	none	none	good	6"
6/17/02 (AFE)		20 plants	light	none	none	good	2'
5/27/03 (BFE)		23 plants	light	none	none	exc.	1' 8"
8/5/03 (AFE)		23 plants	light	none	none	good	2' 9"
Percent surviving as of 6/17/02 was 65%							
Switchgrass 9062193 5' spacing; 25 total planted (plugs) 6/24/97.							
Dates Evaluated							
7/9/1998		17 plants	moderate	none	all plants	fair	3' 5"
9/29/1999		17 plants	moderate	none	all plants	good	4' 5"
4/26/2000		21 plants	moderate	none	all plants	exc.	6' 5"
9/19/2000		21 plants	moderate	none	all plants	exc.	5'
6/26/2001		20 plants	light	none	none	exc.	3' 6"
4/24/02 (BFE)		16 plants	light	none	none	good	5"
6/17/02 (AFE)		14 plants	light	none	none	exc.	2' 6"
5/27/03 (BFE)		19 plants	light	none	none	exc.	1' 5"
8/5/03 (AFE)		19 plants	light	none	none	good	2' 8"
Percent surviving as of 6/17/02 was 56%							
BFE - Before Flooding Event							
AFE - After Flooding Event							
* = Cannot determine rows of plants							

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC						Table #2 - continued		
	% Cover/ Plant #	Active Growing	Weed Comp.	Disease/ Insect	Developed Seed Head	Vigor	Ave. Ht.	
Evaluation Dates:		4/24/02 & 6/17/02						
Cave-In-Rock Switchgrass 23 plants planted.								
Dates Evaluation								
4/18/2000	23	23	severe	none	none	good	5"	
9/19/2000	9	growing weak	severe	none	yes	poor	2'	
6/21/2001	21	21	light	light	none	good	2' 6"	
4/24/02 (BFE)	10	10	light	none	none	good	8"	
6/17/02 (AFE)	12	12	light	light	none	good	2' 6"	
5/27/03 (BFE)	16	16	light	light	none	good	1' 5"	
8/5/03 (AFE)	16	16	light	light	none	fair	2' 6"	
Percent surviving as of 6/17/02 was 52%								
Flood Tolerant Switchgrass, seeded 50' row plus 3' wide.								
Dates Evaluated								
Seeded 6/1/00 50' x 40" plot - .0038 ac. Rate 6# PLS/ac.								
9/19/2000	15%- 20% of 50' row	fair	moderate	none	6/5 5%	good	8"	
6/21/2001	22	22	light	none	none	exc.	3'	
4/24/02 (BFE)	16	16	light	none	none	good	5"	
6/17/02 (AFE)	33	33	light	none	none	good	1' 6"	
5/27/03 (BFE)	45%	45%	light	none	none	good	1' 6"	
8/5/03 (AFE)	45%	45%	light	none	none	good	2' 5"	
Flood tolerant switchgrass plugs block, 63 plants planted 5/25/99.								
Dates Evaluated								
4/26/2000	92%	58 plants	none	none	6/5 100%	exc.	6' 5"	
9/19/2000	95%	95%	none	none	6/5 100%	exc.	4' 5"	
6/21/2001	80%	66 plants	light	none	none	exc.	3'	
4/24/02 (BFE)	85%	66 plants	light	none	none	good	6"	
6/17/02 (AFE)	85%	66 plants	light	none	none	good	2'	
5/27/03 (BFE)	85%	66 plants	light	none	none	good	1' 3"	
8/5/03 (AFE)	85%	66 plants	light	none	none	fair	2' 2"	
Bermudagrass block plugs, planted 5/25/99.								
Dates Evaluated								
9/28/1999	35%	100%	light	none	50%	exc.	3"	
4/26/2000		100%	light	none	none	exc.	3-5"	
9/19/2000	100%	100%	light	none	100%	exc.	9"	
6/21/2001	100%	100%	none	none	none	exc.	6"	
4/24/02 (BFE)	100%	50%	none	none	none	good	2"	
6/17/02 (AFE)	90%	90%	none	none	none	good	3"	
5/27/03 (BFE)	100%	100%	none	none	none	fair	1"	
8/5/03 (AFE)	100%	100%	none	none	none	fair	1"	
BFE = Before Flood Event								
AFE = After Flood Event								

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #3		
Prairie Cordgrass									
2002 Data: Flood Event from 4/29/02 to 5/17/02									
2003 Data: Flooding Began 6/10/03									
		Active					Ave. Ht.	Average	
	Total #	Growing	Weed	Disease/	Developed		Seed	Forage	
	Planted	Spreading	Comp.	Insect	Seed Head	Vigor	Head	Height	
							10' x 10'		
Prairie Cordgrass Collection, planted 9/29/97							3 2 1		
East →							6 5 4		
							9 8 7		
7/9/1998	9	6" average	severe	none		NA exc.	-	-	
8/1/1999	9	30" average	moderate	none		9/9 good	-	-	
9/19/2000	9	4.5" ave.	none	none		9/9 exc.	6'.5"	5'.0 forage	
6/21/2001	9	6'	light	none	none	exc.	6'	45"	
4/24/02 (BFE)	9	7.5'	light	none	none	exc.	none	17"	
6/17/02 (AFE)	9	8'	light	none	none	exc.	none	36"	
5/27/03 (BFE)	9	8.5'	light	none	none	exc.	none	30"	
8/5/03 (AFE)	9	8.5'	light	none	none	exc.	6.5'	40"	
Percent surviving as of 6/17/02 was 100%									
Cuivre Island Prairie Cordgrass Collection, planted 5/15/98							3' x 3'		
					North ↑		4 3 2 1		
							8 7 6 5		
7/9/1998	8	5.5"	severe	none	6 plants	good/exc.	4'.0"	4'.0"	
5/25/1999	8	1'.5" each	moderate	none	none	exc.	none		
		direction							
Lost Creek Prairie Cordgrass Collection, planted 5/15/98							3'x3'		
							12 11 10 9		
							16 15 14 13		
7/9/1998	8	6"	severe	none	4 plants	good/exc.	4'.0"	4'.0"	
5/25/1999	8	1'.5" each	moderate	none	none	exc.	none		
		direction							
9/19/2000									
Total block for both collections			none	none	35%	exc.	6' 0"	5' 0"	
									More lodging Cuivre Island collection
9/19/2000									
14' x 13'5" total spread of blocks			none	none	35%	exc.	6'.0"	More lodging Cuivre Island collection	
9/19/2000									
3' x 3' block is filled in total prairie cordgrass			none	none	35%	exc.	6'.0"	More lodging Cuivre Island collection	
6/26/2001		solid	none	none	none	exc.	6'.0"	50"	
4/24/02 (BFE)		80%	none	none	none	exc,	15"		
6/17/02 (AFE)		solid block	none	none	none	exc.	48"		
5/27/03 (BFE)		solid block	none	none	none	exc.	none	29"	
8/5/03 (AFE)		solid block	none	none	none	exc.	6.5'	42"	
BFE - Before Flooding Event									
AFE - After Flooding Event									

Study: 29A1370 - Wetland/Riparian Propagation, Establishment, and Demonstration

Table #4

Genus/Species	Common Name	Accession No.	Vigor Rating		Date of Rating	
			BFE	AFE	BFE	AFE
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9098840	*	*	5/27/03	8/5/03
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078844	*	*	5/27/03	8/5/03
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078842	*	*	5/27/03	8/5/03
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078846	*	*	5/27/03	8/5/03
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078843	*	*	5/27/03	8/5/03
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078845	*	*	5/27/03	8/5/03
<i>Tripsacum dactyloides</i>	Eastern gamagrass	Pete	*	*	5/27/03	8/5/03
<i>Panicum virgatum</i>	Switchgrass	9062193	Exc.	Good	5/27/03	8/5/03
<i>Panicum virgatum</i>	Switchgrass	9062235	Exc.	Good	5/27/03	8/5/03
<i>Panicum virgatum</i>	Switchgrass	9062213	Exc.	Good	5/27/03	8/5/03
<i>Panicum virgatum</i>	Switchgrass	C-I-R	Good	Fair	5/27/03	8/5/03
<i>Panicum virgatum</i>	Switchgrass Direct Seeded 2001	9083170 Flood-Tolerant	Exc.	Good	5/27/03	8/5/03
<i>Spartina pectinata</i>	Prairie cordgrass	Cuivre Island	Exc.	Exc.	5/27/03	7/5/03
<i>Spartina pectinata</i>	Prairie cordgrass	Lost Creek	Exc.	Exc.	5/27/03	7/5/03
<i>Cynodon dactylon</i>	Bermuda grass	Elsberry	Fair	Fair	5/27/03	7/5/03
<i>Asclepias incarnata</i>	Swamp milkweed	Iowa	Good	Fair	5/27/03	7/5/03
<i>Lobelia cardinalis</i>	Cardinal flower	Forrest Keeling	Good	Poor		7/5/03
<i>Carex scoparia</i>	Broomsedge	MDC	Died			7/5/03
<i>Elymus virginicus</i>	Virginia Wildrye	Cuivre River	Fair	Top Growth Died	5/27/03	7/5/03
<i>Spartina pectinata</i>	Prairie cordgrass Seedlings		Exc.	Exc.	5/27/03	7/5/03
<i>Panicum virgatum</i>	Plugs of switchgrass	9062213 9062235 9062193	Good	Fair	5/27/03	7/5/03

BFE = Before Flood Event

AFE = After Flood Event

- = Cannot determine rows from plants/seed that germinated

Study: 29A1370 - Wetland/Riparian Propagation, Establishment, and Demonstration

Table #5

Genus/Species	Common Name	Accession No.	Vigor Rating		Date of Rating	
			BFE	AFE	BFE	AFE
<i>Andropogon gerardii</i>	Epic Big Bluestem	9083274	Exc.	Fair	4/14/08	5/5/08
<i>Andropogon gerardii</i>	Refuge Big Bluestem	9078832	Exc.	Fair	4/14/08	5/5/08
<i>Panicum virgatum</i>	Low Growing Switchgrass	9083172	Exc.	Exc.	4/14/08	5/5/08
<i>Panicum virgatum</i>	Cave-in-Rock Switchgrass	469228	Exc.	Exc.	4/14/08	5/5/08

BFE = Before Flood Event

AFE = After Flood Event

- = Cannot determine rows from plants/seed that germinated

Study: 29I141G

Study Title: Assembly and Evaluation of Little Bluestem, *Schizachyrium scoparium*, Nichx.

Study Leader: Cordsiemon, R.L.

Introduction:

Little bluestem is a native warm season prairie grass. It was a major component making up as much as 50 percent of the tall grass prairie that was native to much of the Elsberry PMC service area. It can also be a major component of glade areas and mixed grass prairies. Little bluestem can be found in prairies, open woods, dry hills, and fields, from Quebec and Maine to Alberta and Idaho, south to Florida and Arizona.

Problem:

There are no current varieties of little bluestem on the market that have an origin within the three-state service area. Available varieties do not always perform as well as expected. There is a need for an adapted and improved variety of little bluestem for pasture and range seedings, surface mine reclamation, critical area planting, wildlife plantings, recreational area development and other conservation uses in Missouri, Iowa, and Illinois.

Objective:

The objective is to assemble, evaluate, develop and cooperatively release an adapted variety and/or varieties of tested class of little bluestem for conservation use in Missouri, Iowa, and Illinois.

Procedure:

Vegetative material from native ecotypes was collected throughout the states of Missouri, Iowa, and Illinois. A minimum of three collections per Major Land Resource Area/state was requested. (Approximately 60 collections total.) Field selection of collected plant material was based on forage quantity and plant vigor.

Each collection (accession) was one individual plant. A collection was made up of more than one plant if they were in the same immediate area (within five feet) and appeared to be clones of each other.

Discussion:**1996**

The study was approved in July 1996. Collection instructions were sent out and plants were dug in October and November. The samples were picked up shortly after collection and stored in the packing shed at the Plant Materials Center. At this time we received 113 collections from the three-state area. There are a few additional collections expected.

1997 - 1998

The collections were vegetatively propagated in containers in January and grown out in the greenhouse until April. These plants were then transplanted in Field #1 on the PMC April 22-24, 1997 in a randomized complete block with four replications (see Table #2 for map of plot layout). Thirteen additional collections were made in the summer of 1997 and planted into the replications August 14-15, 1997. This brought the total accessions represented to 130: 79 from Missouri, 20 from Illinois, 27 from Iowa, and four standards of comparison. A list of collectors

can be seen in Table #1. First year evaluation consisted of survival. The second year evaluations consisted of survival, height, late dormancy, and form.

1999

The assembly was evaluated in 1999 for forage amount and vigor (Tables #3 and #4). The higher rated plants will have forage quality samples taken in 2000.

2000

The assembly was evaluated for mid season forage production, quality and vigor on June 27, 2000. The entire planting was then clipped to a height of six inches on June 28, 2000. The assembly was evaluated for amount of regrowth and vigor on July 25, 2000 and forage quality samples were taken on August 1, 2000. The assembly was clipped the second time on August 2, 2000 and evaluations for regrowth amount and vigor were taken October 24, 2000.

2001

Evaluations from previous years were correlated and the best plants from the top 10-20 percent of the total accessions were propagated in the greenhouse from clonal material from each individual plant. Plants were then isolated in two locations. A northern region was established containing plants from Iowa, northern Missouri, and northern Illinois. A southern region was established containing plants from southern Missouri and central and southern Illinois. These isolation blocks will receive additional evaluation to remove unwanted plants and the remaining plants will be allowed to produce seed. Plants from this seed will be selected for the next evaluation nursery. After further evaluation, plants from the nursery planted in 2003 will be used as a breeder's block for improved selections. Plants selected for each region can be found in Table #5.

2002

The south region crossing block did very well in 2002. Very few plants were rouged out and seed was harvested from each accession in the block. This seed will be used to establish the next evaluation nursery scheduled for 2003.

The north region crossing block did not do well in 2002. Weed control became a problem and many of the plants were reestablished and did not make seed. Filling in additional plants is scheduled for 2002 and also seed production from this crossing block.

2003

Seed from the south region crossing block was evaluated for quick establishment and plants were grown in the greenhouse for establishment of the recurrent selection evaluation nursery. Approximately 500 plants were transplanted on three foot centers in this evaluation block.

The plants will be allowed to develop and be evaluated for forage. Plants in the north region crossing block were not all equally matured and no seed was harvested from this block.

2004

The plants in the southern region evaluation block were given 2004 to develop and mature. Evaluation of this block will begin in 2005.

Seed was harvested from the northern region crossing block, cleaned, and planted in the greenhouse. These plants were evaluated for quick establishment and seedling vigor. Selected plants will be transplanted into an evaluation nursery.

2005

The plants in the southern region nursery were evaluated based on vigor, amount of forage production, leafiness, drought resistance, disease and insect resistance, and late maturity. Of

312 plants 195 plants were selected (62.5%) and allowed to cross pollinate. Seed was collected from the selected plants to establish a foundation field next year. This south region selection was given the accession number 9083271.

Greenhouse plants selected for seedling vigor from seed harvested from the northern crossing block were transplanted into an evaluation nursery.

2006

A .75 acre foundation field (G1) of southern region selection (accession 9083271) was planted in field #12 on the PMC. Establishment was good but no seed was harvested the first year. Some plants did produce seed but there was not enough to justify a harvest.

The northern region crossing block was evaluated for survival and missing plants were reestablished with greenhouse plants selected for seedling vigor. This evaluation nursery had no further evaluation or selection.

2007

The southern region selection (accession 9083271) G0 block (field 11) was harvested in 2007 but seed was very limited. This seed was used to expand the foundation (G1) field (Field 12). The small expanded portion of the field established poorly. The larger part of the G1 field produced 58.9 pounds bulk seed from approximately 0.60 acre. This seed will be used for field plantings for additional testing.

The northern region crossing block was evaluated for forage production, seed production and late maturity. This block started with 506 plants with a survival of over 95%. There were 159 of these selected and allowed to cross pollinate. Seed was harvested individually from the 159 and will be placed in the germinator and plants germinating the quickest will go to the next evaluation block.

2008

The southern region selection (accession 9083271) of little bluestem, G0 block (field 11) was harvested in 2008 and produced 11.1# bulk seed. Seed was very limited. This seed will be used to establish a new foundation (G1) plot in field 7 because the original increase plot in field 12 has too much indiangrass contamination from the adjoining plot. The G1 plot in field 12 produced 107.6 pounds bulk seed from approximately 0.60 acre. This seed will be used for field plantings for additional testing.

The northern region crossing block (Cycle 2) of little bluestem was again evaluated for forage production, seed production and late maturity. The top 20 individual plants were selected and allowed to cross pollinate. Seed was harvested individually from the 20 plants and placed in the germinator and the plants that germinated the quickest were added to the next block (Cycle 3, Breeders Block at FKN).

2009

The southern region selection (accession 9083271) of little bluestem, G0 block (field 11) was harvested in 2009 and produced 16.4# bulk seed. This seed will be used to enlarge the foundation (G1) plot in field 7 that was planted in 2009 but did not produce seed the first year. The G1 plot in field 12 produced 58.1# bulk seed from approximately 0.60 acre. This seed will be allocated to seed growers interested in producing the new release in 2010.

The northern region crossing block (Cycle 2) of little bluestem was again narrowed to the top 20 plants again and allowed to cross pollinate. Seed was harvested individually from the 20 selections and placed in the germinator and the plants that germinated the quickest were added to the Cycle 3, Breeders Block at FKN. Undesirable plants in this block were rouged out.

2010

Southern region selection (9083271) was released as a selected class release in 2010. The new release name is Ozark Germplasm. A few growers have already picked up the release and it should be available on the commercial market in 2011. The Elsberry PMC harvested over 300 bulk pounds of seed from just over an acre. Seed for this release is currently available to other interested growers.

The northern region (cycle 3 – breeders block) SG0 plot was completed on the FKN property and will be harvested in 2011. Seed from this plot will be used to establish an SG1 (foundation) field. Field testing will begin once seed has been harvested from the SG1 field.

Study 29I141G - Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i>, Nichx.					
Little Bluestem					Table #1
REFERENCE					
ACCESSION	NUMBER	COLLECTOR	MLRA	COUNTY	STATE
9078894	MO-1	Robert S. Crowder	M115	Chariton	Missouri
9078951	MO-2	Robert J. Crowder/ George L. Pollard	109	Chariton	Missouri
9078895	MO-3	Joe Tousignant	N116B	Cape Girardeau	Missouri
9078896	MO-4	Douglas Rainey	M115	Clark	Missouri
9078897	MO-5	David S. Mackey	113	Knox	Missouri
9078898	MO-6	Larry R. Brewer	M109	Putnam	Missouri
9078899	MO-7	Tommy Robins/ Jim Hoefler	116	Ripley	Missouri
9078900	MO-8	Grant P. Butler	N116B	Jefferson	Missouri
9078901	MO-9			Iron	Missouri
9078902	MO-10	Tommy Robins/ Jim Hoefler	116	Carter	Missouri
9078903	MO-11	Arch J. Mueller	M115	Ste. Genevieve	Missouri
9078904	MO-12			St. Francois	Missouri
9078905	MO-13	J. Mark Mitchell		Butler	Missouri
9078906	MO-14	Randy C. Miller	N116A	Shannon	Missouri
9078907	MO-15	Tom Johnson	N116B	Bollinger	Missouri
9078908	MO-16	Tom Johnson	N116A	Bollinger	Missouri
9078909	MO-17	Randy C. Miller	N116B	Reynolds	Missouri
9078910	MO-18			Franklin	Missouri
9078911	MO-19	Tom Johnson	N116A	Wayne	Missouri
9078912	MO-20	Mark E. Nussbaum	N116B	Cape Girardeau	Missouri
9078913	MO-21	Frank Oberle	115	Adair	Missouri
9078914	MO-22	David S. Mackey	113	Knox	Missouri
9078915	MO-23	Claude F. Peifer	116B	Perry	Missouri
9078916	MO-24	Grant P. Butler/ Bryan L. Westfall	N116A	Washington	Missouri
9078917	MO-25	John E. Turner	113/115	Monroe	Missouri
9078918	MO-26	David S. Mackey	113	Knox	Missouri
9078919	MO-27	Douglas Rainey	M115	Clark	Missouri
9078920	MO-28	Frank Oberle	115	Adair	Missouri
9078921	MO-29		M115	Montgomery	Missouri
9078922	MO-30	David S. Mackey	113	Knox	Missouri
9078923	MO-31	Curtis W. Walker	109	Clinton	Missouri
9078924	MO-32	James A. Mayberry	109	Carroll	Missouri
9078925	MO-33	Gary J. Barker	M109	Gentry	Missouri
9078926	MO-34			Vernon	Missouri
9078927	MO-35	Louis Byford		Atchison	Missouri
9078928	MO-36	Todd E. Mason	M109	Worth	Missouri
9078929	MO-37	Louis Byford		Atchison	Missouri
9078930	MO-38	Louis Byford		Atchison	Missouri
9078931	MO-39	Ronald L. Musick	M109	Harrison	Missouri

Study 29I141G - Little Bluestem			Table #1 - continued		
REFERENCE					
ACCESSION	NUMBER	COLLECTOR	MLRA	COUNTY	STATE
9078932	MO-40	Gary J. Barker	M109	Gentry	Missouri
9078933	MO-41	Curtis Walker	109	Gentry	Missouri
9078934	MO-42	Curtis Walker	107	Buchanan	Missouri
9078935	MO-43	Louis Byford		Atchison	Missouri
9078936	MO-44	Ronald L. Musick	M109	Harrison	Missouri
9078937	MO-45	Louis Byford		Atchison	Missouri
9078938	MO-46	Louis Byford		Atchison	Missouri
9078939	MO-47	Bob Sipec		Holt	Missouri
9078940	MO-48	Bib Sipec		Holt	Missouri
9078941	MO-49	Bob Sipec		Holt	Missouri
9078942	MO-50	Ian S. Kurtz	116A	Taney	Missouri
9078943	MO-52	Dennis Shirk/ Ed Gillmore	115	Gasconade	Missouri
9078944	MO-53	Dennis Shirk/ Ed Gillmore	116	Osage	Missouri
9078945	MO-54	Raleigh Redman	112	Henry	Missouri
9078946	MO-55	Dennis Shirk/ Ed Gillmore	116	Maries	Missouri
9078947	MO-56	Jerry Cloyed	M112	Barton	Missouri
9078948	MO-57	Ian S. Kurtz	116A	Taney	Missouri
9078949	MO-58	Ben A. Reed	M112	Barton	Missouri
9078950	MO-59	Jerry Cloyed	M112	Barton	Missouri
9078952	MO-60	M. Denise Brown	N116A	Miller	Missouri
9078953	MO-61	M. Denise Brown	N116B	Miller	Missouri
9078954	MO-62	Howard L. Coambes	N116B	Cedar	Missouri
9078955	MO-63	Howard L. Coambes	N116B	Cedar	Missouri
9078956	MO-64	Douglas G. Newman		Shannon	Missouri
9078957	MO-65	Tom E. Toney		Wayne	Missouri
9078958	MO-66	Rod Doolen		Wayne	Missouri
9078959	MO-67	Rod Doolen		Wayne	Missouri
9078960	MO-68	Kenneth L. Dalrymple		Pike	Missouri
9078963	MO-69	Maurice Davis/ Steve Clubine		Pettis	Missouri
	MO-70	Maurice Davis/ Steve Clubine		Benton	Missouri
	MO-71	Maurice Davis/ Steve Clubine		St. Clair	Missouri
	MO-72	Maurice Davis/ Steve Clubine		Benton	Missouri
9078964	MO-73	Maurice Davis/ Steve Clubine		Pettis	Missouri
9078965	MO-74	Maurice Davis/ Steve Clubine		Pettis	Missouri

Study 29I141G - Little Bluestem				Table #1 - continued	
REFERENCE					
ACCESSION	NUMBER	COLLECTOR	MLRA	COUNTY	STATE
	MO-75	Maurice Davis/ Steve Clubine		Pettis	Missouri
	MO-76	Maurice Davis/ Steve Clubine		Benton	Missouri
9078966	MO-77	Maurice Davis/ Steve Clubine		Maries	Missouri
9078967	MO-78	Dennis Shirk		Maries	Missouri
9078968	MO-79	Steve Clubine		Maries	Missouri
9078969	MO-80	Maurice Davis		Maries	Missouri
9078970	MO-81			Lawrence	Missouri
9078961	IA-27	Robert R. Bryant/ Shawn Dettman	108	Scott	Iowa
9078847	IA-1	Curt Donohue	109	Clarke	Iowa
9078848	IA-2	Curt Donohue	109	Clarke	Iowa
9078849	IA-3	Janet M. Thomas/ John P. Vogel	107	Cherokee	Iowa
9078850	IA-4	John P. Vogel	107	Woodbury	Iowa
9078851	IA-5	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078852	IA-6	Henry D. Tordoff/ Galen Barrett	107	West Pottawattamie	Iowa Iowa
9078853	IA-7	John P. Vogel	107	Woodbury	Iowa
9078854	IA-8	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078855	IA-9	John P. Vogel	107	Plymouth	Iowa
9078856	IA-10	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078857	IA-11	Julie K. Watkins/ Charlie E. Kiepe	108	Franklin	Iowa
9078858	IA-12	Brad Harrison	103	Dallas	Iowa
9078859	IA-13	Shawn A. Dettman	108	Muscatine	Iowa
9078860	IA-14	Jim Ranum	105	Allamakee	Iowa
9078861	IA-15	Rick Cordes	104	Howard	Iowa
9078862	IA-16	James Ranum	105	Allamakee	Iowa
9078863	IA-17	Jay E. Ford	107	Crawford	Iowa
9078864	IA-18	Steve Maternack	103	Polk	Iowa
9078865	IA-19	Jay E. Ford	107	Crawford	Iowa
9078866	IA-20	Jay E. Ford	107	Crawford	Iowa
9078867	IA-21	Al Ehley	104	Cerro Gordo	Iowa
9078868	IA-22	Al Ehley	104	Cerro Gordo	Iowa
9078869	IA-23	John P. Vogel	102	Lyon	Iowa
9078870	IA-24	Jay E. Ford	107	Crawford	Iowa

Study 291141G		Plot Layout Map												Table #2																				
Little Bluestem		Randomized Complete Block																																
		Four Replications																																
		Field #1																																
		North ↑																																
PLT #		1	2	3	4	5 - 28	29	30	31	32	33	34	35 - 58	59	60	61	62	63	64	65 - 76	77	78	79 - 90	91	92	93	94	95	96	97 - 120	121	122	123	124
TIER #		ROADWAY																																
I		REP 1																																
II		REP 2																																
III		REP 3																																
IV		REP 3																																
V		REP 3																																
VI		REP 3																																
VII		REP 3																																
VIII		REP 3																																
IX		REP 3																																
X		REP 3																																
XI		REP 3																																
XII		REP 3																																
XIII		REP 3																																
XIV		REP 3																																
XV		REP 3																																
		Highway JJ																																

Study 291141G		Little Bluestem										Table #2 - continued										Rep #3
PLT #	TIER #	62 63 64	65 66 67	68 69 70	71 72 73	74 75 76	77	78	79 80 81	82 83 84	85 86 87	88 89 90	91 92 93	North								
I		R R R	R R R	R c c	c c c	c c c	c	a	a a a	a b b	j b b	j j j	R R R									
II		MO-45	IL-6	MO-71	IA-13	MO-31	B	a	IL-4	MO-63	MO-11	IL-8	IL-11									
III		MO-61	MO-19	MO-43	MO-50	MO-40	B	R	IA-21	IL-13	IL-17	MO-68	MO-29									
IV		IA-9	MO-51	MO-58	IA-17	MO-55	E	O	MO-47	MO-56	MO-2	MO-13	IL-11									
V		MO-35	MO-1	MO-23	IA-24	MO-24	E	A	IL-5	CAMPER	MO-69	IL-12	MO-25									
VI		MO-39	MO-28	MO-36	MO-42	MO-53	E	D	MO-54	IA-26	IA-14	IA-5	IA-15									
VII		MO-77	IA-19	CIMMERON	IA-18	MO-64	C	W	MO-6	MO-33	MO-73	MO-16	IL-3	IL-8 only								
VIII		MO-9	MO-7	IA-23	IL-20	IA-4	C	A	MO-32	IA-26	MO-52	MO-22	MO-44	one planted								
IX		IA-6	MO-80	IL-2	IA-10	MO-5	G	Y	IA-7	MO-20	IL-16	MO-48	IA-16									
X		MO-8	IA-12	MO-78	MO-30	IA-25	G		MO-79	MO-17	MO-59	MO-14	IL-7									
XI		MO-34	MO-12	MO-46	IA-8	MO-18	I	a	IA-11	IL-21	MO-72	IA-22	PASTURA									
XII		IL-14	MO-26	MO-4	IL-19	MO-38	I	a	MO-74	MO-33	MO-21	MO-65	IL-9									
XIII		IL-18	IA-27	MO-66	ALDOUS	MO-67	O	a	IA-3	MO-27	MO-81	MO-41	IA-20									
XIV		MO-60	MO-10	MO-37	MO-15	MO-62	O	a	MO-49	IL-15	MO-57	IA-1	IL-10									
XV		H H A	A K K	F F D	D L L	M M N	N	a	a c c	c c c	c c c	c c c	h c c									
PLT #		94 95 96	97 98 99	100 101 102	103 104 105	106 107 108	109 110 111	112 113 114	115 116 117	118 119 120	121 122 123	124		Rep #4								
I		R i R	a a a	X X X	X X U	i U U	U U U	U U U	U U U	W W W	W W W	W W W	d d d									
II		IA-9	IL-18	MO-8	MO-74	MO-40	IA-25	MO-5	MO-5	MO-42	IA-4	IA-20	d d d									
III		MO-58	IA-19	MO-28	IL-17	MO-53	IL-8	PASTUR	PASTUR	MO-37	IL-10	MO-77	d d d									
IV		ALDOUS	MO-80	IA-21	MO-2	IA-8	MO-26	IA-26	IA-26	MO-68	MO-14	MO-52	d d d									
V		MO-51	IA-18	MO-20	MO-46	IL-1	MO-1	MO-62	MO-62	MO-44	MO-9	MO-34	d d d									
VI		IA-17	IA-10	MO-33	IA-24	MO-43	IL-12	IA-5	IA-5	MO-81	CIMMERON	MO-19	d d d									
VII		MO-64	IA-10	CAMPER	MO-3	MO-69	MO-61	IA-16	IA-16	IL-4	MO-35	MO-21	d d d									
VIII		IA-27	MO-39	IL-19	MO-57	IL-6	MO-38	MO-67	MO-67	MO-25	MO-48	IL-14	e e e									
IX		MO-60	MO-15	MO-63	IA-7	MO-36	IL-15	MO-49	MO-49	IA-13	MO-29	MO-30	e e e									
X		MO-12	MO-41	MO-32	MO-55	IA-12	MO-47	IA-26	IA-26	IL-21	MO-65	IL-9	e e e									
XI		IL-20	IA-23	IA-11	MO-46	MO-17	IL-2	IL-13	IL-13	MO-45	IL-11	IA-22	f f f									
XII		MO-50	MO-6	MO-59	IA-14	MO-31	MO-54	MO-79	MO-79	IA-3	MO-16	IL-7	f f f									
XIII		MO-71	MO-78	MO-27	MO-73	MO-18	IA-15	MO-66	MO-66	MO-72	MO-22	MO-10	f f f									
XIV		MO-7	MO-11	IL-16	MO-23	IA-1	IL-5	IA-6	IA-6	MO-13	IL-3	MO-56	f f f									
XV		c R R	MO-24	R h R	R S h	h S S	S S T	h h h	h h h	T V V	V h g	g g g	g									

Study 29I141G			Forage Rating: 8/9/99										Table #3		
Little Bluestem															
			1 = High			9 = Low									
Local Number	Rep 1			Rep 2			Rep 3			Rep 4			Percent Survival	Ave. Living Plants	Best Plant Location/s
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12			
MO-7	2	3	1	3	3	2	2	1	2	5	1	3	100	2.33	1 P 1, 8, 11
MO-12	1	2	1	3	2	2	3	2	2	1	1	1	100	1.75	1 P 1, 3, 12, 11, 12
MO-21	1	2	2	6	2	3	4	3	3	4	4	5	100	3.25	1 P 1
MO-74	3	3	5	4	4	4	5	5	4	1	2	1	100	3.42	1 P 10, 12
MO-80	3	3	x	4	5	5	4	4	2	1	4	3	92	3.45	1 P 10
MO-4	x	5	5	4	8	2	3	4	4	6	x	x	83	4.10	2 P 6
MO-9	4	4	4	3	4	4	3	4	3	2	3	3	100	3.42	2 P 10
MO-14	4	4	3	4	4	4	5	2	2	4	4	3	100	3.58	2 P 8, 9
MO-15	3	2	3	5	4	3	6	4	5	4	3	5	100	3.92	2 P 2
MO-22	4	5	5	3	4	2	5	5	6	x	8	x	83	4.70	2 P 6
MO-23	3	5	6	2	6	8	5	4	5	8	8	3	100	5.73	2 P 4
MO-24	3	x	2	x	4	4	3	4	3	3	4	5	83	3.18	2 P 3
MO-32	4	x	8	6	7	3	3	4	5	2	5	6	92	4.82	2 P 10
MO-34	4	4	4	3	4	3	x	x	4	2	x	5	75	3.00	2 P 10
MO-37	2	4	3	7	5	4	x	5	4	3	4	3	92	3.67	2 P 1
MO-42	5	5	6	4	5	2	4	4	4	5	5	7	100	4.67	2 P 6
MO-50	3	3	4	2	2	2	3	4	6	2	3	4	100	3.17	2 P 4, 5, 6, 10
MO-51	3	3	3	3	4	4	4	6	3	4	3	2	100	3.50	2 P 12
MO-53	4	4	5	5	5	5	2	4	5	5	6	7	100	4.75	2 P 7
MO-56	3	3	2	2	5	4	5	3	3	3	3	3	100	3.25	2 P 3, 4
MO-58	3	3	3	5	4	5	5	5	5	2	2	4	100	3.83	2 P 10, 11
MO-59	2	3	4	4	4	5	3	3	3	3	4	4	100	3.50	2 P 1
MO-66	3	3	x	3	3	3	3	2	4	4	5	5	92	3.45	2 P 8
MO-73	7	4	4	3	3	2	4	5	5	7	8	6	100	4.83	2 P 6
MO-79	2	3	2	5	3	5	3	8	5	4	4	3	100	3.92	2 P 1, 3
MO-2	4	5	3	5	5	5	5	3	3	3	4	3	100	4.00	3 P 3, 8, 9, 10, 12
MO-5	7	3	3	5	5	5	6	8	4	4	5	4	100	4.92	3 P 2, 3
MO-8	6	x	5	5	4	5	7	4	8	3	3	4	92	4.91	3 P 10, 11
MO-10	4	5	5	3	3	5	5	5	5	7	5	4	100	4.67	3 P 4, 12
MO-11	x	7	x	4	5	6	6	6	5	3	3	6	83	4.25	3 P 10, 11
MO-13	5	8	5	5	x	5	4	4	3	6	4	6	100	4.58	3 P 9
MO-16	4	3	8	6	6	54	5	6	4	4	5	100	75	3.00	3 P 2
MO-17	4	4	3	4	3	7	8	6	5	4	5	5	100	4.83	3 P 3, 5
MO-18	3	4	3	7	7	8	x	x	x	5	5	5	75	3.92	3 P 1, 3
MO-19	3	5	5	3	4	3	4	6	5	3	5	4	100	4.17	3 P 1, 4, 6, 10
MO-20	8	7	6	7	6	5	3	4	5	4	8	3	100	6.60	3 P 7, 12
MO-25	3	3	x	5	5	5	5	4	6	5	5	6	92	4.33	3 P 1, 2
MO-26	3	4	4	5	x	4	3	4	4	3	4	5	92	4.30	3 P 1, 7, 10
MO-27	5	6	3	4	5	4	6	5	4	5	5	7	100	5.36	3 P 3
MO-29	4	3	x	4	5	4	4	6	3	3	5	8	92	4.45	3 P 2, 9, 10
MO-30	3	4	5	7	7	x	4	4	7	4	3	4	92	4.73	3 P 1, 11
MO-31	7	3	4	4	4	6	7	8	x	5	5	5	92	5.27	3 P 2

Study 29I141G			Forage Rating: 8/9/99										Table #3 - continued		
Little Bluestem															
			1 = High					9 = Low							
Local Number	Rep 1			Rep 2			Rep 3			Rep 4			Percent Survival	Ave. Living Plants	Best Plant Location/s
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12			
MO-33	3	x	3	5	5	3	4	5	5	8	8	4	92	5.89	3 P 1, 3, 6
MO-35	4	7	8	5	6	7	5	3	6	5	4	x	92	5.45	3 P 8
MO-38	6	6	5	3	3	4	4	6	7	3	3	4	100	5.40	3 P 4, 5, 10, 11
MO-41	5	6	5	4	4	7	6	x	4	3	x	5	83	4.90	3 P 10
MO-43	4	4	x	5	5	5	5	6	5	4	3	4	92	4.55	3 P 11
MO-46	4	x	4	4	3	3	3	5	5	4	4	4	92	3.91	3 P 5, 6, 7
MO-47	5	6	6	6	5	4	3	4	5	5	8	4	100	5.08	3 P 7
MO-48	3	7	8	5	5	6	4	4	6	4	5	5	100	5.17	3 P 1
MO-52	3	3	3	4	3	3	4	5	4	4	3	4	100	3.58	3 P 1, 2, 3, 5, 6, 11
MO-54	x	x	x	5	5	5	4	5	5	6	4	3	75	4.67	3 P 12
MO-57	4	4	x	3	5	x	4	4	x	5	4	3	92	3.27	3 P 4, 12
MO-60	7	4	6	4	6	3	6	4	6	5	5	4	100	5.00	3 P 6
MO-61	5	8	6	x	4	5	x	8	8	3	7	5	83	5.90	3 P 10
MO-65	4	5	6	7	x	x	4	5	3	4	6	6	83	5.00	3 P 9
MO-67	3	3	3	3	3	3	6	5	x	3	3	3	92	3.45	3 P 1, 2, 3, 4, 5, 6, 10, 11, 12
MO-69	4	5	4	3	3	5	4	5	4	7	4	5	100	4.42	3 P 3, 4
MO-71	x	5	5	4	3	5	4	4	5	4	5	3	92	4.27	3 P 5, 12
MO-77	6	x	6	4	6	4	3	4	5	6	6	5	92	5.00	3 P 7
MO-78	5	6	5	5	3	5	3	5	6	4	3	3	100	4.42	3 P 5, 7, 11, 12
MO-1	4	5	4	4	4	6	4	7	5	4	5	5	100	4.75	4
MO-3	4	7	4	5	4	4	4	4	4	5	4	5	100	4.50	4
MO-6	7	7	7	7	7	5	x	8	7	4	4	4	92	6.09	4
MO-28	6	5	6	6	7	5	4	7	7	4	x	x	83	4.75	4
MO-36	4	4	5	6	6	6	x	5	5	5	6	5	92	5.18	4
MO-39	4	6	7	4	6	4	6	5	x	6	5	x	83	5.89	4
MO-40	7	6	7	5	4	4	x	6	5	5	5	5	92	5.36	4
MO-44	7	4	5	5	6	7	7	x	6	5	4	6	92	5.64	4
MO-45	4	4	4	5	6	6	5	6	5	4	4	4	100	4.75	4
MO-49	6	5	6	6	5	x	5	5	4	7	5	6	92	5.45	4
MO-55	x	6	x	4	4	5	4	5	x	8	x	5	67	5.13	4
MO-62	4	4	5	5	4	5	5	7	6	5	5	6	100	5.08	4
MO-63	5	6	5	5	4	4	8	4	6	4	5	5	100	5.08	4
MO-68	7	6	6	6	8	4	5	6	5	4	4	4	100	5.42	4
MO-72	5	6	5	5	6	5	4	6	6	5	4	4	100	5.08	4
MO-81	x	4	5	5	4	6	x	x	x	6	x	8	58	5.43	4
MO-64	x	7	6	7	6	6	6	5	8	x	7	5	92	5.73	5
MO-70															
MO-75															
MO-76															

Study 29I141G				Forage Rating: 8/9/99												Table #3 - continued			
Little Bluestem																			
				1 = High						9 = Low									
Local Number	Rep 1			Rep 2			Rep 3			Rep 4			Ave. Percent Survival	Living Plants	Best Plant	Location/s			
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12							
IA-16	x	x	4	3	6	5	3	x	1	x	5	5	75	3.56	1	P 9			
IA-27	1	1	3	3	4	5	5	5	4	5	4	2	100	3.50	1	P 1, 2			
IA-6	4	5	6	5	2	4	3	4	3	7	4	5	100	4.33	2	P 5, 6			
IA-8	5	6	3	5	3	5	5	5	5	5	3	2	100	4.33	2	P 12			
IA-12	7	5	7	x	4	5	4	3	2	4	5	5	92	4.64	2	P 9			
IA-15	5	4	5	x	x	x	2	x	5	5	5	6	67	4.63	2	P 7			
IA-23	6	5	5	8	8	6	5	4	x	2	4	6	92	5.36	2	P 10			
IA-1	8	5	5	5	4	4	4	5	x	3	7	3	92	4.82	3	P 10, 12			
IA-2	4	4	4	3	4	4	6	5	5	4	x	6	92	4.45	3	P 4			
IA-3	x	x	8	x	3	3	4	5	4	4	5	4	75	4.44	3	P 5, 6			
IA-4	5	8	4	3	x	3	4	7	5	4	7	5	92	5.00	3	P 4, 6			
IA-5	4	5	4	3	6	8	6	4	4	3	5	x	92	4.73	3	P 4, 10			
IA-7	5	3	3	5	5	5	4	4	6	5	5	5	100	4.58	3	P 2, 3			
IA-9	4	6	7	6	6	6	8	6	6	4	3	4	100	5.50	3	P 11			
IA-11	6	5	6	5	7	3	5	5	6	4	x	5	92	5.18	3	P 6			
IA-13	4	4	6	4	7	x	5	4	x	3	4	3	83	4.40	3	P 10, 12			
IA-17	3	7	4	5	x	4	6	x	6	4	6	5	83	5.00	3	P 1			
IA-19	6	x	x	6	3	3	x	4	4	x	x	x	50	4.33	3	P 5, 6			
IA-20	x	4	x	7	5	5	4	x	4	6	7	3	75	5.00	3	P 12			
IA-24	4	5	3	5	4	4	4	4	5	5	5	4	100	4.33	3	P 3			
IA-25	4	5	6	6	5	6	6	4	5	3	5	3	100	4.83	3	P 10, 12			
IA-26	x	3	4	3	3	6	x	x	4	5	6	x	67	4.25	3	P 2, 4, 5			
IA-10	6	7	7	4	5	5	5	6	7	6	4	x	92	5.64	4				
IA-14	4	6	4	5	5	6	4	5	5	5	7	5	100	5.08	4				
IA-18	5	6	5	6	5	6	5	4	5	4	5	5	100	5.08	4				
IA-21	4	5	4	4	x	6	x	x	6	-	4	5	67	4.75	4				
IA-22	x	x	x	7	x	x	7	6	6	5	8	8	58	6.71	5				
IL-12	8	7	5	3	8	4	5	5	4	4	2	x	92	5.00	2	P 11			
IL-17	3	4	3	2	3	5	3	4	2	2	3	3	100	3.08	2	P 4, 9, 10			
IL-18	5	4	6	3	3	3	5	6	4	3	2	4	100	4.00	2	P 11			
IL-2	6	6	6	4	5	6	5	3	5	4	5	3	100		3	P 8			
IL-5	6	5	7	4	8	3	4	5	5	5	4	5	100	5.08	3	P 6			
IL-7	4	4	3	4	7	6	8	6	8	6	8	8	100	6.00	3	P 3			
IL-8	x	x	5	4	x	8	x	6	4	x	4	3	58	4.86	3	P 12			
IL-11	x	x	3	x	4	x	5	x	6	x	x	x	33	4.50	3	P 3			
IL-14	4	5	x	3	5	x	6	4	7	6	5	6	83	5.10	3	P 4			
IL-16	5	5	4	4	3	3	4	x	3	7	6	4	92	4.36	3	P 5, 6, 9			
IL-19	5	6	7	3	3	3	4	3	4	3	4	3	100	4.00	3	P 4, 5, 6, 8, 12			

Study 29I141G			Vigor Rating: 8/9/99											Table #4		
Little Bluestem			1 = High			9 = Low										
Local Number	Rep 1			Rep 2			Rep 3			Rep 4			Percent Survival	Living Plants Ave.	Best Plant Location/s	
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12				
MO-4	x	3	4	4	6	2	4	5	5	3	x	x	75	4.00	2	P 6
MO-7	2	3	2	3	3	3	5	2	2	5	2	2	100	2.83	2	P 1, 3, 8, 9, 11, 12
MO-12	3	3	3	4	2	2	4	4	3	3	3	3	100	3.08	2	P 5, 6
MO-16	3	2	6	6	4	3	4	5	6	4	5	3	100	4.25	2	P 2
MO-24	5	x	5	x	5	3	5	5	5	2	4	6	83	4.50	2	P 10
MO-25	2	3	x	5	4	3	5	4	4	5	6	6	92	4.27	2	P 1
MO-32	3	x	6	5	5	3	4	6	4	2	4	6	92	4.36	2	P 10
MO-35	2	6	7	2	4	5	6	6	3	5	4	x	92	4.55	2	P 1, 4
MO-42	5	4	5	3	4	2	4	4	6	4	5	6	100	4.33	2	P 6
MO-47	4	5	6	4	5	4	2	2	3	4	6	3	100	4.00	2	P 7, 8
MO-56	3	4	3	3	3	2	4	4	4	4	3	3	100	3.33	2	P 6
MO-61	5	5	4	x	3	4	x	7	7	2	5	4	83	4.60	2	P 10
MO-67	3	3	3	2	3	3	5	4	x	4	5	5	92	3.64	2	P 4
MO-69	4	5	6	3	3	4	2	3	5	8	4	5	100	4.33	2	P 7
MO-79	2	3	3	3	3	4	5	6	4	5	4	3	100	3.75	2	P 1
MO-1	3	4	3	4	3	5	5	5	5	3	5	4	100	4.08	3	P 1, 3, 5, 10
MO-3	3	4	4	5	4	3	4	5	5	4	3	4	100	4.00	3	P 1, 6, 11
MO-5	5	3	3	5	4	6	5	7	4	5	6	4	100	4.75	3	P 2, 3
MO-6	3	7	6	6	5	5	x	5	5	5	5	3	92	5.00	3	P 1, 12
MO-8	5	x	4	6	3	3	6	6	5	6	5	7	92	5.09	3	P 5, 6
MO-9	5	5	6	3	3	3	4	4	4	5	5	5	100	4.33	3	P 3, 4, 5
MO-11	x	5	x	5	6	6	7	5	3	5	4	6	83	5.20	3	P 9
MO-13	5	7	6	6	x	5	5	6	3	6	5	7	92	5.55	3	P 9
MO-14	4	4	3	5	5	5	4	6	6	4	5	4	100	4.58	3	P 3
MO-15	3	3	3	4	3	3	5	4	4	4	4	4	100	3.67	3	P 1, 2, 3, 5, 6
MO-17	5	5	5	4	4	7	7	5	4	3	4	5	100	4.83	3	P 10
MO-19	3	3	3	4	4	4	4	5	5	4	4	4	100	3.92	3	P 1, 2, 3, 5, 6
MO-21	3	3	3	6	4	4	5	4	4	6	6	6	100	4.50	3	P 1, 2, 3
MO-22	4	3	3	3	3	3	5	5	4	x	7	x	83	4.00	3	P 2, 3, 4, 5, 6
MO-23	5	5	3	4	5	7	5	5	6	5	7	5	100	5.17	3	P 3
MO-26	4	4	4	3	x	3	6	5	5	4	5	5	92	4.36	3	P 4, 6
MO-27	3	5	3	3	4	3	6	6	5	3	4	6	100	4.25	3	P 1, 3, 4, 6, 10
MO-29	4	3	x	6	5	5	5	5	3	5	5	6	92	4.73	3	P 2, 9
MO-31	6	3	3	3	3	4	5	5	x	6	4	5	92	4.27	3	P 2, 3, 4, 5
MO-33	4	x	6	4	4	3	4	4	4	6	6	5	92	4.55	3	P 6
MO-34	4	3	3	3	3	4	x	x	4	4	x	3	75	3.44	3	P 2, 3, 4, 5, 12
MO-36	4	3	3	6	5	5	x	4	5	3	5	5	92	4.36	3	P 2, 3, 10
MO-37	3	3	3	4	3	4	x	5	5	4	4	4	92	3.82	3	P 1, 2, 3, 5
MO-38	4	4	3	5	4	4	3	5	5	3	3	4	100	3.92	3	P 7, 10, 11
MO-39	5	6	7	4	3	3	5	3	x	5	7	x	83	4.80	3	P 5, 6, 8
MO-40	3	8	8	4	5	3	x	5	4	8	8	7	100	5.25	3	P 1, 6, 11

Study 29I141G			Vigor Rating: 8/9/99											Table #4 - continued			
Little Bluestem			1 = High			9 = Low											
Local Number	Rep 1			Rep 2			Rep 3			Rep 4			Percent Survival	Living Plants	Best Plant	Location/s	
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12					
MO-43	6	3	4	4	4	4	5	6	5	4	5	3	100	4.42	3	P 2, 12	
MO-45	4	4	3	3	4	3	4	5	3	3	4	4	100	3.67	3	P 3, 4, 6, 9, 10	
MO-46	3	x	3	3	3	4	5	5	3	5	3	4	92	3.73	3	P 1, 3, 4, 5, 9, 11	
MO-48	4	5	5	3	4	4	5	3	5	4	6	6	100	4.50	3	P 4, 8	
MO-51	4	5	4	3	3	3	4	5	4	4	4	4	100	3.92	3	P 3, 4, 5	
MO-52	5	4	5	5	3	4	5	6	5	5	4	5	100	4.67	3	P 5	
MO-53	5	5	6	4	5	6	3	4	4	5	5	6	100	4.83	3	P 7	
MO-54	x	x	x	5	7	3	6	7	7	6	3	4	75	5.33	3	P 11	
MO-60	4	4	4	3	4	3	5	3	5	5	6	6	100	4.33	3	P 4, 6, 8	
MO-62	4	4	4	3	4	5	4	4	4	5	6	7	100	4.50	3	P 4	
MO-63	4	4	4	3	3	3	5	5	4	4	6	4	100	4.08	3	P 4, 5, 6	
MO-65	3	4	4	6	x	x	5	6	5	5	7	6	83	5.10	3	P 1	
MO-66	5	5	x	4	3	3	6	6	5	6	7	7	92	5.18	3	P 5, 6	
MO-71	x	3	5	5	3	4	5	4	5	3	4	4	92	4.09	3	P 2, 5, 10	
MO-72	3	3	3	3	5	4	3	4	5	5	4	3	100	3.75	3	P 1, 2, 3, 4, 7, 12	
MO-73	6	5	3	3	3	3	5	7	4	6	7	6	100	4.83	3	P 3, 4, 5, 6	
MO-77	6	x	6	5	3	5	3	4	5	6	6	6	92	5.00	3	P 5, 7	
MO-78	6	4	4	4	6	4	4	5	3	4	4	3	100	4.25	3	P 9, 12	
MO-80	4	3	x	3	3	3	6	6	5	3	6	6	92	4.36	3	P 2, 4, 5, 6, 10	
MO-81	x	3	5	5	4	4	x	x	x	6	x	5	58	4.57	3	P 2	
MO-2	4	5	5	4	5	6	4	4	5	4	4	4	100	4.50	4		
MO-18	4	6	4	4	5	7	x	x	x	6	4	6	75	5.11	4	P 1, 3, 4, 11	
MO-20	4	6	6	6	5	5	6	5	5	4	6	4	100	5.17	4		
MO-28	6	4	5	4	6	5	5	6	5	4	x	x	83	5.00	4		
MO-30	4	5	5	4	4	x	5	5	6	5	4	4	92	4.64	4		
MO-41	4	7	4	5	5	4	6	x	5	4	x	4	83	4.80	4		
MO-44	6	4	4	5	5	5	7	x	6	5	4	6	92	5.18	4		
MO-49	8	8	8	8	8	x	7	7	6	6	4	4	92	6.73	4		
MO-50	5	5	5	4	4	4	6	6	4	5	5	5	100	4.83	4		
MO-55	x	5	x	4	6	5	5	4	x	6	x	4	67	4.88	4		
MO-57	4	5	x	5	4	x	6	5	x	5	6	5	75	3.75	4		
MO-58	6	5	4	6	5	6	7	7	7	4	4	5	100	5.50	4		
MO-59	7	6	5	5	4	4	7	6	7	6	6	5	100	5.67	4		
MO-68	5	5	5	4	5	5	5	4	4	6	4	5	100	4.75	4		
MO-74	5	6	6	4	4	5	5	5	5	5	5	4	100	4.92	4		
MO-10	6	7	7	5	5	5	5	6	6	7	6	4	100	5.75	5		
MO-64	x	7	7	5	7	7	6	6	6	x	7	5	83	6.30	5		
MO-70																	
MO-75																	
MO-76																	

Study 29I141G			Vigor Rating: 8/9/99												Table #4 - continued		
Little Bluestem																	
			1 = High			9 = Low											
Local Number	Rep 1			Rep 2			Rep 3			Rep 4			Percent Survival	Living Plants	Best Plant	Location/s	
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12					
IA-3	x	x	5	x	3	2	6	6	7	7	5	5	75	5.11	2	P 6	
IA-4	4	5	3	4	x	3	4	6	4	2	5	5	92	4.09	2	P 10	
IA-5	6	6	6	4	5	6	6	5	6	2	5	x	92	5.18	2	P 10	
IA-9	4	4	4	4	4	5	6	5	5	3	2	5	100	4.25	2	P 11	
IA-10	3	4	5	3	4	4	5	5	5	6	2	x	92	4.18	2	P 11	
IA-13	2	3	4	3	5	x	5	4	x	4	5	3	92	3.45	2	P 1	
IA-15	5	4	4	x	x	x	2	x	6	4	4	5	67	4.25	2	P 7	
IA-27	2	2	2	2	3	3	5	6	5	4	3	3	100	3.33	2	P 1, 2, 3, 4	
IA-1	6	3	3	5	5	4	4	4	x	4	7	4	92	4.45	3	P 2, 3	
IA-2	3	3	3	4	5	5	6	5	5	5	x	6	92	4.55	3	P 1, 2, 3	
IA-6	6	4	4	4	3	3	5	4	4	7	3	5	100	4.33	3	P 5, 6, 11	
IA-7	3	3	4	3	3	3	3	4	6	4	4	4	100	3.67	3	P 1, 2, 4, 5, 6, 7	
IA-8	5	6	3	3	3	4	5	6	5	4	3	4	100	4.25	3	P 3, 4, 5, 11	
IA-12	4	5	6	x	5	4	3	5	4	3	3	3	92	4.09	3	P 7, 10, 11, 12	
IA-14	6	5	5	3	3	3	5	7	7	4	6	5	100	4.92	3	P 4, 5, 6	
IA-16	x	x	4	3	5	4	3	x	5	x	5	6	67	4.38	3	P 4, 7	
IA-17	4	6	5	4	x	4	5	x	4	3	5	3	83	4.30	3	P 10, 12	
IA-18	5	6	5	5	4	5	4	4	5	3	3	4	100	4.42	3	P 10, 11	
IA-23	4	4	4	5	6	6	5	5	x	3	3	4	100	4.08	3	P 10	
IA-25	5	5	5	5	4	4	4	5	5	4	4	3	100	4.42	3	P 12	
IA-26	x	6	4	3	4	5	x	x	4	4	6	x	67	4.50	3	P 4	
IA-11	7	6	7	4	5	4	6	6	7	5	x	5	92	5.64	4		
IA-19	6	x	x	5	4	4	x	4	4	x	x	x	50	4.50	4		
IA-20	x	4	x	7	5	5	5	x	6	5	6	5	75	5.33	4		
IA-21	4	4	5	4	x	5	x	x	4	x	5	4	67	4.38	4		
IA-22	x	x	x	5	x	x	5	4	4	6	8	8	58	5.71	4		
IA-24	5	5	4	6	6	6	7	7	7	6	5	5	100	5.75	5		
IL-8	x	x	6	4	x	5	x	2	3	x	5	3	58	4.00	2	P 8	
IL-12	6	6	2	3	5	3	4	4	3	3	2	x	92	3.73	2	P 3, 11	
IL-1	7	x	3	5	7	6	5	6	8	6	5	5	92	5.73	3	P 3	
IL-2	3	3	4	4	5	3	4	5	5	5	4	4	100	4.08	3	P 1, 2, 6	
IL-3	3	7	3	5	x	x	6	7	6	5	x	x	67	5.25	3	P 1, 3	
IL-5	5	5	6	5	3	4	5	6	5	5	4	5	100	4.83	3	P 5	
IL-6	7	5	4	8	3	5	x	x	x	5	4	7	75	5.33	3	P 5	
IL-9	5	x	3	x	4	5	5	3	3	5	4	6	92	3.91	3	P 3, 8, 9	
IL-10	4	4	5	5	4	3	x	x	8	x	6	6	100	3.75	3	P 6	
IL-11	x	x	3	x	4	x	3	x	5	x	x	x	33	3.75	3	P 3, 7	
IL-13	x	5	x	4	5	5	6	6	7	x	6	3	75	5.22	3	P 12	
IL-14	5	4	x	3	4	x	5	3	5	5	4	5	83	4.30	3	P 4, 8	
IL-15	5	7	x	x	5	4	6	6	5	4	4	3	83	4.90	3	P 12	

Study 29I141G - Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i> Michx.					
Selected accessions for each region					Table #5
Northern Region					
IA - All					
MO - North of Missouri River					
IL - Northern 2/3rds of the state					
REFERENCE					
ACCESSION	NUMBER	COLLECTOR	MLRA	COUNTY	STATE
9078896	MO-4	Douglas Rainey	115	Clark	Missouri
9078913	MO-21	Frank Oberle	115	Adair	Missouri
9078914	MO-22	David S. Mackey	113	Knox	Missouri
9078924	MO-32	James A. Mayberry	109	Carroll	Missouri
9078934	MO-42	Curtis Walker	107	Buchanan	Missouri
9078849	IA-3	Janet M. Thomas/ John P. Vogel	107	Cherokee	Iowa
9078854	IA-8	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078861	IA-15	Rick Cordes	104	Howard	Iowa
9078862	IA-16	James Ranum	105	Allamakee	Iowa
9078884	IL-12	Bill Kleiman		Lee	Illinois
9078891	IL-19	Martha E. Sheppard	115	Calhoun	Illinois
Southern Region					
MO - South of Missouri River					
IL - Southern 1/3 of state					
REFERENCE					
ACCESSION	NUMBER	COLLECTOR	MLRA	COUNTY	STATE
9078895	MO-3	Joe Tousignant	N116B	Cape Girardeau	Missouri
9078899	MO-7	Tommy Robins/ Jim Hoefer	116	Ripley	Missouri
9078915	MO-23	Claude F. Peifer	116B	Perry	Missouri
9078942	MO-51	Ian S. Kurtz	116A	Taney	Missouri
9078950	MO-59	Jerry Cloyd	M112	Barton	Missouri
9078952	MO-60	M. Denise Brown	N116A	Miller	Missouri
9078964	MO-73	Maurice Davis/ Steve Clubine		Pettis	Missouri
9078965	MO-74	Maurice Davis/ Steve Clubine		Pettis	Missouri
9078968	MO-79	Steve Clubine		Marries	Missouri
9078969	MO-80	Maurice Davis/			Missouri
9078893	IL-21	Remington T. Irwin	114	Wayne	Illinois

Study No. 29I142G

Study Title: Production of Native Missouri Ecotypes of Grasses, Legumes and Forbs for Roadsides, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted.

Study Leader: Cordsiemon, R.C.

Introduction:

Well-adapted native grass, legume and forb plantings offer many advantages as a low cost sustainable vegetative cover for management of soil and water resources. Native plant communities resist noxious weed invasion, provide excellent erosion control, and generally require relatively low maintenance.

These characteristics make native plants an excellent selection for use in roadside plantings, wildlife habitat enhancement, long-term land retirement programs, public land and all other vegetative plantings where mono-cultures of grasses are presently being planted. This is especially true along public transportation corridors that constitute a major land resource and management problem in the state of Missouri. Based on 1987 National Resource Inventory (NRI) data, over one million acres of Missouri land are devoted to rural transportation. Other federal and state agencies also own a significant land base in Missouri.

Proper vegetation management along these corridors is an important element in controlling soil loss and unwanted weedy plant species. Many of these acres are now seeded to introduce cool-season grass and legume species which are often invaded by noxious weeds requiring extensive mowing or herbicide treatment programs. These management techniques are expensive and can also result in additional water quality problems where herbicides are used extensively.

Managing or reseeding these acres to promote native grasses and forbs offers a low cost environmentally sound approach to roadside vegetation management. Herbicide use, soil erosion, and most mowing can be reduced significantly where a vigorous native grass and forb mixture dominates a roadside right-of-way. In addition, these goals are consistent with on-going NRCS programs designed to improve ground and surface water quality, reduce soil loss and increase wildlife habitat.

Problem:

Many adapted forb, legume and grass species of native origin are either currently not commercially available or available only in very limited quantities, which make them very expensive. Species that are available are often varietal releases that have undergone an evaluation and selection process or a plant-breeding program. Most varieties are designed for high forage production and are highly vigorous plants. They are generally excellent for pasture and hay production but can be too domineering for diversified mixtures. Their origins are often not from within the state in which they are being planted. There is a need for additional native species for use on public lands and other types of conservation plantings with origins close to where they are being planted.

Objective:

The objective of this study is to accelerate the availability of selected native grass, legume and forb species.


Cooperators:

The Missouri Department of Conservation (MDC), USDA Natural Resources Conservation Service (NRCS), Plant Materials Center (PMC), the University of Missouri at Columbia, Missouri (UMC), and the National Audubon Society-Audubon Missouri (NAS).

Procedures:

The state of Missouri was divided into four zones: Northern Glaciated Plains, Zone #1; Western Prairie, Zone #2; Ozarks, Zone #3; and the Bootheel Region, Zone #4 (See Table #1). Plant materials were collected as seed by the study coordinator, selected personnel from USDA-NRCS, MDC, UMC, and other knowledgeable interested persons. Collections were made from prairie remnants throughout each zone striving for a relatively equal and representative sample. Large collections from one site were not allowed to dominate the mixture from throughout the zone. Seed from each collection site was inventoried by location. Seed collected from within each zone was kept separate from the other zones. Increase plots were and will be established, as seed becomes available. Each species will be released as ‘Source Identified’ germplasm from the zone in which it was collected. Evaluation and selection or plant breeding procedures has not improved ‘Source Identified’ seed.

Table #1



Discussion:

1997

The Missouri Ecotype Enhancement Program was officially started as a plant materials study with the signing of the study plan in December of 1997. This plan is an agreement between cooperators and funded by a grant from the MDC. Several meetings preceded the document signing that included MDC, NRCS, UMC, Department of Transportation, Missouri Department of Natural Resources, and other interested individuals.

The initial grant from MDC to UMC was received July 1997 and a program coordinator was hired by UMC in September 1997 to work at the Elsberry Plant Materials Center.

1998

A grant was given to UMC once again by MDC that would fund the program through August of 1999. Goals were established for 1998 collections. Some species from 1997 were recollected and new species were added.

1999

The Missouri Ecotype program continued during 1999 and the species released listed in Table #2. Beginning in September, the Lincoln County Soil and Water Conservation District took over as the administrator for the Missouri Ecotype Program replacing UMC. MDC funded the program for the 1999-2000 fiscal year.

2000

The Missouri Ecotype program continued through August until funding was depleted. The program was continued under direction of Missouri Audubon Society and MDC in cooperation with the NRCS Plant Materials Center.

2001

The Missouri Ecotype program is growing increase plots at Elsberry and also at the Charles Green Conservation area near Ashland, Missouri.

2002

MDC took over as administrator of this study and is currently still funding the program with the aid of grants. The Missouri Ecotype program is continuing to increase plots at the Elsberry PMC and Green Conservation Area. New collections are being made of both old and new species.

2003

MDC is continuing to administrate the Missouri Ecotype program. All plots are still in production and seed is being allocated. The PMC is planning to increase plots for seed production in 2004.

2004

In the spring of 2004, the plots of zones 1 and 2 sideoats gramma, *Bouteloua curtipendula*, and zone 1 river oats, *Chasmanthium latifolium*, were increased for seed production. There were no

new releases from the Missouri Ecotype Program in 2004 and there are no releases scheduled for 2005. Plans are to increase river oats-zone 1 again in 2005. Becky Erickson, Missouri Ecotype Program Coordinator, has several production plots located at the Green Conservation Area in Ashland, Missouri. The number of plots on the Green Area has increased over the past year and now sustains almost 35 different ecotypes. Plans for both the Plant Materials Center and the plots at the Green Conservation Area are to sustain the plots already established and increase the plots that display good seed production and survival. This will allow for potential releases as early as 2006.

2005

Chasmanthium latifolium, river oats from the northern zone were increased in the late summer of 2005. Plans are to increase several other better seed producing species in the spring of 2006, such as the big bluestem (northern zone), little bluestem (northern zone), sideoats gramma (northern and western zones), tall dropseed (northern and western zones), pale purple coneflower (northern and western zones), grayhead coneflower (northern zone), and Virginia wild rye (northern zone).

2006

Supplemental funding for the Missouri Ecotype Program was not extended past the agreement date of June 2007. Plans are to continue growing the species that have been previously released or have potential for high use in the commercial market. The PMC planted production plots of big bluestem (northern zone), little bluestem (northern zone), sideoats gramma (northern and western zones), tall dropseed (northern and western zones), pale purple coneflower (northern and western zones), and grayhead coneflower (northern zone) in the spring of 2006. A fall planting of Virginia wild rye (northern zone) was also planted. The plots were increased to provide more seed production and averaged ¼ to two acres in size.

2007

Production plots of sideoats grama were increased to help meet the demand for higher priority species requested by seed growers. In FY 2008, plans are to again increase these production plots. The PMC continued to put efforts into a select group of species (production species started in 2006) as they are the priority for most commercial growers interested in Missouri ecotypes. Those species are big bluestem (northern zone), little bluestem (northern zone), sideoats gramma (northern and western zones), tall dropseed (northern and western zones), pale purple coneflower (northern and western zones), and grayheaded coneflower (northern zone). Plans are to release northern Missouri sideoats grama in 2008. With growing interest in southern zone little bluestem, the Elsberry PMC will expand the production plot in 2008.

2008

The flood of 2008 dramatically affected seed production of Northern Missouri germplasm sideoats grama, Northern Missouri germplasm big bluestem, and Northern Missouri germplasm tall dropseed. All three production fields were in the floodplain (field 7) and required transplanting to salvage the plantings. The entire 0.3 acre plot of sideoats grama was moved to field #1 and approximately a 0.1 of an acre of big bluestem was moved to field #1 also.

Approximately 0.1 of an acre of tall dropseed was moved next to the northern zone of big bluestem production plot. Plans are to increase both northern and western zones of pale purple coneflower by dormant planting both plots in December or January. Northern and western zones of sideoats grama will potentially be planted with the pale purple coneflower as a companion crop to discourage heavy weed competition. Both species would be harvested for seed production. The Northern Missouri germplasm little bluestem is doing well, despite heavy inundation of water from spring rains in the south half of the production plot. Seed production remains consistent in the north half of the plot.

2009

All of the plots that were moved because of the 2008 flood survived and produced seed in 2009. The seed harvested will be used to start new production fields. The northern and western zones of pale purple coneflower were planted with a companion species, sideoats grama. Both plots were mowed during the growing season to discourage weed competition. In 2009, the northern and western zones of pale purple coneflower were released, they will be known as Northern Germplasm pale purple coneflower and Western Germplasm pale purple coneflower (*Echinacea pallida*).

2010

The Elsberry PMC is currently working approximately 15 different accessions from this study. Emphasis is focused around pollinator material and shorter growing warm season grasses for prairie restoration. Several growers are still interested in the material from this study and this continues to be the driving force behind the production of these species at the PMC.

Study 29I142G - Missouri Ecotype Releases						Table #2
Missouri Ecotype Releases from the Elsberry Plant Materials Center						
Scientific Name	Release Name	Common Name	Accession Number	Cooperating Agency(ies)	Type of Release	Year of Release
<i>Schizachyrium scoparium</i> , Michx.	Southern MO	little bluestem	9079006	MOPMC, MDC	N	2004
<i>Ratibida pinnata</i>	Northern MO	grayheaded coneflower	9079060	MOPMC, MDC	N	2004
<i>Sporobolus compositus</i> var.	Northern MO	tall dropseed	9079040	MOPMC, MDC, NAS	N	2001
<i>Coreopsis palmata</i>	Northern MO	prairie coreopsis	9079028	MOPMC, MDC, NAS	N	2001
<i>Coreopsis palmata</i>	Western MO	prairie coreopsis	9079029	MOPMC, MDC, NAS	N	2001
<i>Echinacea pallida</i>	Northern MO	pale purple coneflower	9079032	MOPMC, MDC, NAS	N	2009
<i>Echinacea pallida</i>	Western MO	pale purple coneflower	9079033	MOPMC, MDC, NAS	N	2009
<i>Liatris pycnostachya</i>	Northern MO	prairie blazing star	9079020	MOPMC, MDC, NAS	N	2001
<i>Liatris pycnostachya</i>	Western MO	prairie blazing star	9079021	MOPMC, MDC, NAS	N	2001
<i>Elymus virginicus</i> L.	Northern MO	Virginia wildrye	9079044	MOPMC,UMC,MDC,MODOT	N	1999
<i>Sorghastrum nutans</i> (L) Nash.	Northern MO	Indiangrass	9079036	MOPMC,UMC,MDC,MODOT	N	1999
<i>Sorghastrum nutans</i> (L) Nash.	Western MO	Indiangrass	9079037	MOPMC,UMC,MDC,MODOT	N	1999
<i>Andropogon gerardii</i> Vitman	Northern MO	big bluestem	9079000	MOPMC,UMC,MDC,MODOT	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Northern MO	little bluestem	9079004	MOPMC,UMC,MDC,MODOT	N	1999
Cooperating Agencies: MOPMC=Missouri Plant Materials Center; UMC=University of Missouri at Columbia; MDC=Missouri Department of Conservation; MODOT=Missouri Department of Transportation; NAS=National Audubon Society-Audubon Missouri; Grow Native.						
N= native releases; collected within the USA, occurring naturally in the USA. Generally refers to a plant which occurs naturally in a particular region, state ecosystem or habitat without direct or indirect human activity.						
Nat.=naturalized releases; collected from a population within the USA, but were originally introduced to the USA sometime in the past.						
I=introduced; means that the original collection from which the release was made was not from within the USA.						

Study ID Code: MOPMC-P-0001-WO, WL, WE

Study Title: Assembly, Evaluation and Selection of Bur Oak, *Quercus macrocarpa* Michx.

Study Leader: Cordsiemon, R.

Description:

Bur oak is a large-size tree 60-80 feet tall and 2-3 feet in diameter (max. 170 by 7 feet); crown rounded with large, heavy branches. Leaves are deciduous, oblong to ovate; 6-12 inches long; characteristically 5-9 lobed, with rounded lobes. Fruit matures in one year; acorns are 3/5-2 inches long, ellipsoidal, brown, enclosed for 1/3 to all of its length in a characteristic fringe-margined cup. Twigs are stout; yellow-brown to gray, often with characteristic corky wings. Winter buds; 1/8-1/4 inch long, hairy. Bur oak is one of the largest American Oaks. Commonly distributed throughout Missouri, Iowa and Illinois, bur oak are is important bottomland tree, frequently found in moist flats, wetlands, and undulating flood plains. Important associates of bur oak include red maple, American elm, silver maple, swamp white oak, sycamore and eastern cottonwood.

Objective:

The objective of this study is to select a local source, fast growing, and high nut producing bur oak.

Materials and Methods:

Field collections were assembled, accessioned, and held in storage until the collection period ended. The assemblage of collections began at the PMC in October 2000 and ended mid-December 2000. After the collection period was over the seed was stratified and planted in the greenhouse using the Root Pruning Method (RPM) containers. The plants will be transplanted in Field #7 on the PMC in mid to late April 2002. The design will be a randomized complete block with one plant per plot: one block for the Iowa collections, one for the Illinois collections and one block for the Missouri collections.

Discussion

2000

A total of 24 collections were made from the PMC three state service area: seven from Iowa, two from Illinois and 15 from Missouri. As these collections arrived at the PMC they were given accession numbers and placed in stratification for 120 days (cool moist storage 38 degrees Fahrenheit). At the time this report was being developed, these collections were being germinated in the greenhouse.

2001

The 24 collections of bur oaks were taken out of the germination trays and placed in containers (3 5/8" x 6") and allowed to grow to approximately one foot tall. These plants were later transplanted into one-gallon size containers and placed in the portable greenhouse. In early December 2001 the plants were transported to the root cellar for over wintering. The scheduled

planting date is April 2002. The plantings will be randomized complete block designs with one block for Iowa's collections, one block for Illinois' collections and one block for Missouri's collections. Refer to Table #1 for collection information.

2002

Three assemblies of bur oaks were planted in April 2002 representing each state's collections, Iowa, Illinois and Missouri. Iowa's collections were planted in Field #6 on April 18, 2002, Illinois' collections were planted in Field #12 on April 17, 2002, and Missouri's collections were planted in Field #7 on April 18 – 19, 2002. These collections were evaluated for height, spread, vigor, and insect and disease resistance. The evaluation data was not documented in this year's report but will be in the 2003 Annual Technical Report. Table #1 reflects collection information.

2003

The three assemblies of bur oak representing the Missouri, Illinois, and Iowa collections were evaluated in October 2003. Performance characteristics evaluated were height, spread, vigor, and insect/disease resistance. The 2003 plant performance summaries can be found in Tables #2 and #8.

2004

All three plantings, Missouri, Iowa, and Illinois, were evaluated again for height, spread, vigor, and insect/disease resistance. Evaluation data for 2004 can be found in Tables #3 and #9. The Iowa and Missouri plantings originally were planted with two trees of each collection in each replication. Not all collections had enough material to allow for two trees in each replication, but most did. In 2004, the lesser dominant tree was removed to allow the dominant tree to grow without competition. There are some replications that had trees die and the extra (non-dominant) trees were, in some cases, used to replace trees that died. The 2005 evaluations will reflect the replacement trees. Fertilizer, 13-13-13, was added to the three plantings to encourage growth and healthier plants.

2005

Evaluations were taken in September for height, spread, vigor, and insect/disease resistance on the three plantings. Fertilizer, 13-13-13, again was added to each tree. Acorns were produced on a few trees in the Missouri accession in field #7. Evaluation data for 2005 can be found in Tables # 4 and #10.

2006

In early spring, all three plots of bur oaks were sprayed with an ounce per acre rate of Oust to control grass and weed competition. There was good control through mid-summer. Late summer broadleaf weeds and grasses encroached back around the tree. Fertilizer, 13-13-13, was also added at a rate of 8 oz. per tree. An evaluation for acorn production was done on all three plots (five year old trees); Missouri plot (Rep. 3, tree MO-11) produced several acorns and in the Iowa plot (Rep. 3, tree IA-5; Rep. 6, tree IA-6; and Rep. 7, tree IA-5) all produced significant amounts of acorns. There were no height, spread, and insect/disease resistant evaluations taken this year.

2007

In 2007, the seven surviving trees from Illinois in field #12 were moved to field #11 to make room for another planting. During the summer, these seven trees stressed heavily due to drought conditions. There were no evaluations for the Illinois trees and a determination will be made in FY 2008 if there is a need to continue with an Illinois source bur oak. There were no evaluations made on the Illinois planting in FY 2007. Missouri and Iowa bur oak plots look good and are performing well. Oust again was used in the spring to control white clover and grasses around the trees. Missouri source bur oaks have been susceptible to small galls that cover the leaf surface, possibly caused by small wasps. The damage is mostly cosmetic and seems to be mainly associated with the Missouri plot, but is also evident in the Iowa planting. The Iowa and Missouri plots were evaluated again for height, spread, vigor, insect/disease resistance, and acorn production. This year the trees were measured at 24 inches high to get a diameter measurement on the trunk. This will help determine how well each tree is performing against other trees in the planting. The Iowa planting had six trees that produced acorns, while the Missouri planting did not have any trees produce in 2007. The drought conditions could possibly have been the reason for little to no acorn production. The Iowa and Missouri evaluation data can be found in Tables #5 and #11.

2008

The remaining Illinois source trees died in FY08 and plans to start a new evaluation block of trees from Illinois were discontinued. The Missouri and Iowa source bur oak trees are doing very well. Again they were evaluated and plans are to evaluate the plots once more in FY09, before selections from each plot are made. Selections will consist of using a hydraulic tree spade and physically moving the trees to a new location, where they will be planted in a new crossing block. There will be a block for both Missouri and Iowa collections. The trees will be analyzed using statistical software based on the data that has been recorded over the past 7 years. Trees from accessions possessing superior characteristics will be selected for the crossing block. Data for 2008 can be found in Tables #6 and #12.

2009

Both plantings, IA and MO, were evaluated in October of 2009. The plantings were sprayed with a non-selective herbicide around the base of the trees in early spring to allow moisture and fertilizer to access the trees roots. 13-13-13 was also added to both plantings. All trees with limbs lower than 30 inches were cut. Once the evaluations have been analyzed, selections for superior trees will be made in FY10 and a new block will be established after the trees go dormant in the fall of 2010. Data for 2009 can be found in Tables #7 and #13.

2010

Selections were made in both the Iowa and Missouri plantings and unwanted trees were removed using a chainsaw. The trees, in both plots, will be allowed to cross and acorns will be collected in order to start seedlings for field testing. In 2011, smaller trees of the selected accessions may be removed depending on their size in proportion to the rest of the plot. Release information will be compiled, as well as data from field test information in order to make a potential release by 2013-2014.

Table # 1

Study Title: Assembly, Evaluation and Selection of Bur Oak *Quercus macrocarpa* Michx.

Temporary No.	State	County	MLRA	Collector
MO-1	Missouri	Calloway	115	Thomas L. Wekenborg
MO-2	Missouri	Chariton	NA	Charles Lewis
MO-3	Missouri	Shannon	053	Randy Misser
MO-4	Missouri	Lincoln	115	Jimmy Henry
MO-5	Missouri	Lincoln	115	Jimmy Henry
MO-6	Missouri	Lincoln		Wayne Lovelace
MO-7	Missouri			
MO-8	Missouri	Pike		Keith Jackson
MO-9	Missouri	Pike		Keith Jackson
MO-10	Missouri	Pike		Keith Jackson
MO-11	Missouri	Pike		Keith Jackson
MO-12	Missouri	Howard	N/A	Robert D. Dewitt
MO-13	Missouri	Boone	N/A	Robert D. Dewitt
MO-14	Missouri	St. Charles	115	Dan Crigler
MO-15	Missouri	Moniteau	115	Douglas Wallace
IL-1	Illinois	Clark	N/A	David E. Hiatt
IL-1	Illinois	Jasper	113	Dennis D. Clency
IA-1	Iowa	Dickinson	103	Tim K. Moran
IA-2	Iowa	Dickinson	103	Tim K. Moran
IA-3	Iowa	Dickinson	103	Tim K. Moran
IA-4	Iowa	Wayne	N/A	Duane Bedford
IA-5	Iowa	Decatur	109	Kevin Reynolds
IA-6	Iowa	Bremer	104	Richard J. Cornes
IA-7	Iowa	Black	104	Rick Cordes

2003 Evaluation

Summary of Iowa Collections, Located in Field #6

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	1.1	0.4	0.8	1.1	1.3	0.6	0.8	1.0	0.8	1.0	1.2	1.2	0.93
IA-2	0.7	0.8	0.8	0.8	1.2	1.0	1.3	0.8	0.8	0.8	1.1	1.3	0.95
IA-3	0.8	0.7	0.8	1.1	0.9	0.7	0.6	0.8	0.9	1.0	0.9	0.8	0.81
IA-4	1.3	1.3	1.2	0.8	0.8	1.0	0.0	0.9	1.3	1.1	0.9	0.0	1.07
IA-5	1.1	1.3	1.3	0.7	1.3	1.2	2.0	1.9	1.0	1.4	1.7	0.8	1.30
IA-6	0.8	0.7	1.0	1.1	1.0	1.0	0.4	0.3	0.5	1.1	0.6	0.7	0.77
IA-7	1.1	1.3	0.8	1.0	1.8	1.0	0.8	0.5	0.5	1.7	0.9	1.6	1.08

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1		0.3	0.2	0.2	0.0			0.1	0.2	0.1	0.2	0.3	0.17
IA-2		0.3		0.1	0.2	0.2	0.1	0.2	0.1	0.0	0.0	0.2	0.15
IA-3			0.1	0.2	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.08
IA-4		0.2	0.1	0.1	0.1		0.0	0.1	0.1	0.2	0.2		0.14
IA-5		0.3	0.1	0.1	0.3	0.1	0.1	0.3	0.0	0.0	0.0	0.1	0.13
IA-6				0.2	0.2	0.1		0.0	0.1	0.2	0.0	0.2	0.13
IA-7	0.2	0.3	0.1		0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.11

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	4	5	4	3	4	6	4	4	6	4	4	4	4.33
IA-2	4	4	4	4	3	4	6	4	5	5	4	3	4.17
IA-3	dead	4	3	3	4	4	6	6	4	4	5	6	4.45
IA-4	3	3	3	4	4	4		4	3	4	4		3.60
IA-5	3	3	3	4	3	3	2	2	6	3	3	3	3.17
IA-6	4	4	3	3	4	4		9	7	4	6	6	4.91
IA-7	3	3	4	3	2	4	4	7	7	3	4	3	3.92

Summary of Insect and Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	3	3	2	2	2	2	5	4	3	2	3	2	2.73
IA-2	2	2	2	2	3	2	3	5	4	4	3	2	2.91
IA-3	2	3	3	3	2	3	3	3	2	4	3	3	2.91
IA-4	2	2	3	2	2		4	4	3	2	3		2.78
IA-5	3	2	3	3	3	2	2	2	4	2	2	2	2.45
IA-6	2	2	2	2	3	2	4	3	4	3	4	3	2.91
IA-7	3	3	2	2	2	3	2	3	3	3	4	3	2.73

2004 Evaluation

Summary of Iowa Collections, Located in Field #6

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	2.2	4.0	2.8	2.9	2.6	2.4	2.2	2.3	3.7	3.6	3.6	3.5	3.0
IA-2	2.9	3.1	2.4	2.7	2.9	3.1	2.4	3.6	3.5	3.4	3.7	3.8	3.1
IA-3	3.7	2.3	2.5	3.0	2.7	3.2	2.4	dead	3.6	2.5	3.3	3.4	3.0
IA-4	3.0	4.0	3.4	4.1	3.2	3.1	3.8	3.1	4.4	3.8	4.1	4.3	3.7
IA-5	4.3	3.7	4.9	3.7	3.5	3.2	3.2	3.2	3.5	dead	3.2	2.6	3.5
IA-6	2.3	3.0	2.4	2.4	3.6	3.5	dead	dead	3.4	3.1	2.2	3.5	2.9
IA-7	3.5	3.2	3.2	3.2	3.7	3.6	3.2	4.1	2.4	3.4	3.8	4.1	3.5

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	2	3.6	2.5	3.2	2.9	2.5	3	2	3.8	3.7	3	3.2	3.0
IA-2	2.7	2.7	2.4	2.1	2.2	2.8	3	3	2.6	3	2.9	2.6	2.7
IA-3	3.7	2.1	2.8	2.7	2.4	3.3	1.5	dead	3.3	3.1	2.2	3.6	2.8
IA-4	3	2.6	3.1	3.4	3	3.4	5	3.7	3	3.1	3.4	3.4	3.3
IA-5	4.4	3.2	3.3	4.1	3.8	2.5	2	3.4	2.9	dead	3.1	2.4	3.2
IA-6	2	3.6	2.2	2.4	3.4	3.4	dead	dead	3.8	3.2	2.6	2.5	2.9
IA-7	2.8	2.6	2.3	2.4	3.5	3.2	3.3	3.5	2.2	3.7	3	3.4	3.0

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	5	3	5	3	4	5	4	5	3	3	3	4	3.9
IA-2	4	3	5	5	4	4	5	3	4	4	3	4	4.0
IA-3	2	5	4	3	5	4	6	dead	3	4	5	3	4.0
IA-4	3	3	2	2	3	3	2	3	2	2	3	2	2.5
IA-5	1	3	1	2	2	4	5	3	4	dead	4	5	3.1
IA-6	5	3	5	4	3	3	dead	dead	3	3	5	4	3.8
IA-7	3	4	3	3	2	3	4	4	4	3	3	3	3.3

Summary of Insect and Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	2	3	3	3	1	4	2	3	2	2	2	2	2.4
IA-2	1	2	2	1	2	2	4	2	2	2	3	2	2.1
IA-3	2	2	3	3	2	3	2	dead	2	2	2	2	2.3
IA-4	2	3	2	2	3	2	3	2	3	2	2	1	2.3
IA-5	2	2	2	2	1	3	3	3	3	dead	5	2	2.5
IA-6	2	2	2	2	2	1	dead	dead	1	1	1	2	1.6
IA-7	2	3	2	1	2	2	2	2	2	2	4	2	2.2

Study MOPMC-P-0001-WE, WL																									
Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i> , Michx.													Table #4												
2005 Evaluation																									
Summary of Iowa Collections, Located in Field #6																									
Summary of Height (feet)																									
Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.												
IA-1	3.6	4.5	4.3	4.6	5.1	3.5	4.7	3.9	5.3	5	4.9	5.4	4.6												
IA-2	4.3	4.6	3.8	3.8	4.5	4.9	3.6	4.4	4.6	4.9	5.2	5	4.5												
IA-3	6.3	4	4	5.6	3.7	5.2	4	Dead	6	4.5	5.2	4.3	4.8												
IA-4	5.2	5.4	4.8	6	4.6	5.1	5.8	5.2	6.2	5.4	6.3	6	5.5												
IA-5	5.8	5.6	6.3	5.3	5.2	4.6	4.6	5	5.2	Dead	5	2.9	5												
IA-6	3.5	5.2	4.2	4.5	5	4.5	Dead	Dead	4.3	5.1	3.2	4.4	4.4												
IA-7	4.5	Dead	5.8	5.2	5.5	5	4.7	4.6	Dead	4.7	4.9	5.7	5.1												
Summary of Spread (feet)																									
Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.												
IA-1	2.6	3	2.3	3.6	5.1	3.4	3.8	3.6	3.7	4	4.2	3.3	3.6												
IA-2	2.6	3.2	2.6	3.6	3	3.5	3.6	3.6	3.6	3.2	3.6	3.1	3.3												
IA-3	4.2	2.6	3	4	2	3	2.6	Dead	4.9	3	2.4	4.3	3.3												
IA-4	4	3	3.5	5	3.6	4	4.7	4	3.4	4	3.4	4.4	3.9												
IA-5	4.3	3.4	4.5	4	3.6	5	3.7	4	3.2	Dead	3	3.4	3.8												
IA-6	2.2	4	2.5	3.5	2.5	4	Dead	Dead	3.6	4.2	2.4	2.6	3.2												
IA-7	2.8	Dead	2.3	3.4	3.5	3.5	4.2	3.9	Dead	3	4	4	3.5												
Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor																									
Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.												
IA-1	6	3	5	3	1	3	4	5	3	2	3	3	3.4												
IA-2	5	4	5	5	5	2	6	3	3	3	3	3	3.9												
IA-3	2	5	4	5	5	5	5	Dead	2	4	4	3	4.0												
IA-4	3	3	3	1	3	3	1	2	2	3	2	6	2.7												
IA-5	1	3	1	2	2	3	3	2	3	Dead	4	2	2.4												
IA-6	6	3	5	3	3	4	Dead	Dead	3	3	6	4	4.0												
IA-7	9	Dead	5	4		2	3	3	Dead	5	3	2	4.0												
Summary of Insect/Disease Resistance (1-9 Rating) 1=Very Good 9=Poor																									
Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.												
IA-1	3	2	2	2	2	2	2	2	3	3	2	2	2.3												
IA-2	2	2	3	3	4	2	3	2	3	3	3	2	2.7												
IA-3	3	4	2	7	3	4	1	Dead	1	3	2	2	2.9												
IA-4	1	2	1	1	1	3	1	1	1	2	1	2	1.4												
IA-5	2	3	2	2	3	1	1	3	1	Dead	2	1	1.9												
IA-6	2	3	1	2	3	2	Dead	Dead	2	2	2	4	2.3												
IA-7	9	Dead	5	3		1	1	3	Dead	2	2	2	3.1												

2007 Evaluation

Summary of Iowa Collections, Located in Field #6

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	9.8	10.2	9.0	9.7	8.4	9.0	9.8	9.5	10.1	9.5	9.1	10.6	9.56
IA-2	8.0	9.4	7.5	7.2	6.8	7.5	9.8	9.4	9.7	8.2	8.6	6.7	8.23
IA-3	11.8	8.0	8.5	8.0	7.2	8.5	8.5	dead	11.4	9.0	8.9	8.5	8.94
IA-4	9.7	9.2	10.2	9.7	9.6	9.6	11.0	10.6	11.7	12.0	11.9	10.6	10.48
IA-5	11.6	10.8	11.0	9.8	9.6	7.6	8.3	11.2	10.6	dead	9.5	6.9	9.72
IA-6	5.6	9.0	7.7	9.3	9.3	7.0	dead	dead	10.8	8.9	8.0	7.3	8.29
IA-7	9.8	7.5	9.3	9.0	9.1	9.4	10.2	9.6	5.8	9.2	11.0	11.2	9.26

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	6.7	7.3	6.9	8.5	7.0	6.8	8.0	7.6	9.7	8.2	9.7	7.2	7.80
IA-2	5.7	8.0	7.4	6.8	6.0	7.5	9.3	8.1	9.8	6.9	7.9	6.0	7.45
IA-3	10.6	6.6	6.8	8.0	5.0	6.7	7.4	dead	10.0	7.2	5.6	8.6	7.50
IA-4	6.9	6.5	8.0	9.6	9.0	7.7	11.5	9.8	8.8	9.2	10.2	10.9	9.01
IA-5	11.0	8.9	9.2	8.2	8.0	8.7	9.3	11.5	8.3	dead	10.0	6.0	9.01
IA-6	5.0	8.5	6.8	7.5	10.2	8.2	dead	dead	9.5	8.4	5.7	7.0	7.68
IA-7	7.7	4.0	8.0	7.6	8.0	8.2	10.7	10.0	4.0	6.6	10.0	9.9	7.89

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	4	5	6	3	5	6	4	5	3	3	3	3	4.17
IA-2	5	5	6	7	8	6	3	4	3	5	5	7	5.33
IA-3	3	7	6	5	7	6	5	dead	2	5	5	4	5.00
IA-4	5	5	3	3	3	5	1	1	1	1	1	1	2.50
IA-5	2	4	2	3	4	4	4	1	2	dead	3	6	3.18
IA-6	7	5	5	4	4	6	dead	dead	2	5	5	6	4.90
IA-7	4	8	6	5	5	3	2	4	7	4	1	1	4.17

2007 Evaluation

Summary of Iowa Collections, Located in Field #6

Table #5 - Continued

Summary of Insect/Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	5	3	4	3	3	5	4	3	3	3	1	3	3.33
IA-2	3	3	3	3	4	4	2	3	2	3	4	3	3.08
IA-3	4	2	3	3	3	3	1	dead	3	8	3	3	3.27
IA-4	4	3	2	2	1	2	3	1	3	3	2	2	2.33
IA-5	3	3	3	2	3	3	3	2	2	dead	3	2	2.64
IA-6	2	3	3	2	2	3	dead	dead	2	2	3	3	2.50
IA-7	3	3	5	3	3	3	3	3	2	4	2	3	3.08

Summary of Trunk Diameter Measured at 24 Inches High

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	5	6.5	5.5	8	6.25	6.75	5	5.5	7.75	6.5	7	6.25	6.33
IA-2	5.5	5.75	5.5	5	4	6.25	6.75	5.75	6.25	6.25	6	3.5	5.54
IA-3	8.25	4	5.75	5	6	5.5	4.5	dead	5.75	6.75	5.75	5.25	5.68
IA-4	9.25	8.2	7.75	8.5	7.25	8	9	8.25	7	7	9.5	9	8.23
IA-5	10	6.5	8.25	7.5	7.75	7.75	6.25	7.75	7.25	dead	6	4.5	7.23
IA-6	3.5	6.5	4.75	5.5	7.5	6	dead	dead	6.5	5.5	4.25	6	5.60
IA-7	6	3.5	5.25	5	7.5	6.75	6	5.75	2.25	6.25	6.75	5.75	5.56

Summary of Acorn Presence

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12
IA-1	no	YES	no	no	no							
IA-2	no	no	no									
IA-3	no	dead	no	no	no	no						
IA-4	no	no	no									
IA-5	YES	no	no	no	no	no	no	YES	no	dead	YES	YES
IA-6	no	no	no	YES	no	no	dead	dead	no	no	no	no
IA-7	no	no	no									

2008 Evaluation

Summary of Iowa Collections, Located in Field #6

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	11.3	12.2	10.7	10.8	10.0	9.9	10.0	11.3	12.2	10.6	10.0	11.7	10.9
IA-2	8.2	10.7	9.4	9.4	8.3	8.8	11.1	10.4	12.2	9.0	8.9	8.2	9.6
IA-3	13.2	9.1	10.3	9.4	9.7	9.4	9.3	DEAD	13.2	10.2	10.0	8.8	9.4
IA-4	12.0	12.0	12.5	13.2	11.2	11.1	12.2	12.3	12.9	13.4	12.4	12.3	12.3
IA-5	13.0	11.4	13.3	12.0	11.1	9.3	10.6	13.0	12.6	DEAD	10.4	8.8	10.5
IA-6	6.8	10.6	8.9	10.9	10.8	9.2	DEAD	DEAD	11.5	9.9	8.6	9.4	8.1
IA-7	10.2	8.3	11.1	10.0	10.0	11.6	12.0	11.7	8.4	11.6	13.3	12.2	10.9

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	7.5	8.9	6.9	7.5	8.5	7.6	7.6	8.1	9.6	8.5	7.8	6.1	7.9
IA-2	5.6	7.7	8.3	7.4	6.1	8.1	9.4	8.4	10.4	6.7	8.8	6.7	7.8
IA-3	9.9	6.4	7.4	8.7	6.0	7.6	8.4	DEAD	11.2	7.4	8.0	8.9	7.5
IA-4	7.0	7.9	11.7	8.9	10.1	8.3	11.0	8.7	9.5	12.1	9.5	9.2	9.5
IA-5	11.4	8.8	11.4	9.2	8.4	10.0	10.8	11.5	8.4	DEAD	9.6	7.5	8.9
IA-6	5.9	10.0	8.7	8.6	11.4	8.9	DEAD	DEAD	8.2	8.6	4.8	8.2	6.9
IA-7	7.1	5.3	7.0	7.5	10.7	7.9	9.6	10.3	5.2	10.3	9.8	10.4	8.4

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	8	9	7	8	8	8	8	8	8	8	9	8	8.1
IA-2	7	7	7	6	6	6	7	8	9	7	7	6	6.9
IA-3	10	7	7	7	7	7	7	DEAD	10	7	8	7	7.0
IA-4	9	9	10	10	10	9	10	10	10	10	10	10	9.8
IA-5	10	9	10	9	9	9	9	10	9	DEAD	8	7	8.3
IA-6	6	8	7	8	9	6	DEAD	DEAD	8	6	7	7	6.0
IA-7	8	5	8	7	9	9	9	8	5	8	10	10	8.0

Summary of Insect/Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	8	8	8	9	8	8	8	8	7	9	10	9	8.3
IA-2	7	8	9	9	8	8	7	8	8	9	9	7	8.1
IA-3	9	8	7	9	9	9	8	DEAD	9	8	8	9	7.8
IA-4	7	9	9	10	10	9	10	10	10	10	10	10	9.5
IA-5	7	9	9	9	9	8	10	10	10	DEAD	9	8	8.2
IA-6	7	9	8	9	10	6	DEAD	DEAD	9	8	9	10	7.1
IA-7	8	8	8	8	8	9	9	8	8	9	10	10	8.6

Summary of Diameter @ 24"

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
----------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--------	--------	--------	------

IA-1	2.00	2.50	2.25	3.00	2.50	2.50	2.00	2.25	2.50	2.75	2.50	2.50	2.44
IA-2	2.25	2.50	2.75	2.00	1.75	2.25	2.75	2.50	3.00	2.50	2.25	1.50	2.33
IA-3	3.25	1.75	2.25	1.75	2.25	2.25	2.00	DEAD	2.50	2.25	2.25	1.75	2.02
IA-4	3.25	3.50	3.25	3.75	3.00	3.25	3.75	3.25	3.00	3.50	3.50	3.25	3.35
IA-5	3.50	2.50	3.50	2.50	3.00	2.50	2.50	3.50	2.75	DEAD	2.50	2.00	2.56
IA-6	1.25	2.25	2.00	2.25	3.00	2.00	DEAD	DEAD	2.50	2.25	2.00	2.50	1.83
IA-7	2.50	1.50	2.25	2.00	2.75	2.75	2.75	2.50	1.00	2.25	3.00	2.25	2.29

Summary of Circumference

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Avg.
IA-1	6.28	7.85	7.07	9.42	7.85	7.85	6.28	7.07	7.85	8.64	7.85	7.85	7.65
IA-2	7.07	7.85	8.64	6.28	5.50	7.07	8.64	7.85	9.42	7.85	7.07	4.71	7.33
IA-3	10.21	5.50	7.07	5.50	7.07	7.07	6.28	DEAD	7.85	7.07	7.07	5.50	6.35
IA-4	10.21	10.99	10.21	11.78	9.42	10.21	11.78	10.21	9.42	10.99	10.99	10.21	10.53
IA-5	10.99	7.85	10.99	7.85	9.42	7.85	7.85	10.99	8.64	DEAD	7.85	6.28	8.05
IA-6	3.93	7.07	6.28	7.07	9.42	6.28	DEAD	DEAD	7.85	7.07	6.28	7.85	5.76
IA-7	7.85	4.71	7.07	6.28	8.64	8.64	8.64	7.85	3.14	7.07	9.42	7.07	7.20

Acorns Present

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Rep 11	Rep 12	Total
IA-1	N	N	N	N	N	N	N	N	N	N	N	N	0
IA-2	N	N	N	N	N	N	YES	N	N	N	YES	N	2
IA-3	N	N	N	YES	N	YES	N	DEAD	N	N	N	N	2
IA-4	N	N	N	N	N	N	N	N	N	N	N	N	0
IA-5	N	N	N	YES	YES	N	N	YES	YES	DEAD	YES	N	5
IA-6	N	N	N	YES	N	YES	DEAD	DEAD	N	N	N	N	2
IA-7	N	N	N	N	N	YES	YES	N	N	N	YES	N	3

2003 Evaluation

Summary of Missouri Collections, Located in Field #7

Summary of Height (Feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Average
MO-1	2.1	1.2	2.1	1.5	1.3	1.6	1.7	1.8	1.7	1.6	1.7
MO-2	1.7	2.0	1.6	2.2	1.5	0.8	0.6	0.6	0.7	1.1	1.3
MO-3	1.4	1.1	0.6	0.4	0.8	1.3	1.9	2.0	1.4	2.0	1.3
MO-4	1.6	1.3	1.7	1.9	2.1	0.7	1.5	1.3	1.5	1.2	1.4
MO-5	1.5	1.1	1.3	1.1	1.2	1.4	0.6	1.6	1.1	1.4	1.2
MO-6	1.6	0.0	0.0	0.0	0.0	2.0	1.0	2.0	1.7	1.9	1.7
MO-7	1.8	1.8	1.6	1.8	1.5		1.0	1.0	1.3	1.4	1.4
MO-8	1.1	0.7	1.2	0.5	1.1	2.1	1.6	1.9	2.1	1.6	1.4
MO-9	1.8	1.4	1.6	2.3	1.0	1.5	1.5	2.1	1.4	1.7	1.6
MO-10	2.0	1.5	2.3	2.3	1.6	0.8	1.5	1.3	1.5	1.3	1.6
MO-11	1.2	2.0	2.0	2.0	0.9	0.6	1.5	1.8	2.2	1.5	1.6
MO-12	2.0	1.9	1.7	1.1	0.8	1.3	1.5	1.5	2.0	1.5	1.5
MO-13		2.1	1.7	2.1	1.5	2.1	2.1	1.6	2.0	1.4	1.9
MO-14	1.8	1.8	2.3	1.7	1.1	1.3	1.2	1.5	1.7	1.4	1.6
MO-15			1.2	1.4	1.6						1.4

Summary of Spread (Feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Average
MO-1	0.1	0.2	0.2	0.0	0.0	0.2	0.5	0.3	0.3	0.2	0.2
MO-2	0.4	0.2	0.2	0.3	0.4	0.1	0.1	0.2	0.0	0.4	0.2
MO-3	0.1	0.0	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.1	0.1
MO-4	0.1	0.1	0.1	0.0	0.1	0.1	0.2	0.1	0.0	0.1	0.1
MO-5	0.1	0.1	0.1	0.0	0.1	0.2	0.2	0.1	0.3	0.1	0.1
MO-6	0.2								0.1	0.3	0.2
MO-7	0.2	0.1	0.1	0.2	0.3		0.0	0.3	0.1	0.2	0.2
MO-8	0.3	0.2	0.3	0.1	0.3	0.3	0.1	0.2	0.1	0.1	0.2
MO-9	0.3	0.1	0.2	0.3	0.1	0.3	0.1	0.3	0.0	0.3	0.2
MO-10	0.3	0.2	0.2	0.1	0.1		0.4	0.2	0.1	0.0	0.1
MO-11	0.1	0.2	0.2	0.1	0.0	0.1	0.1	0.1	0.1	0.3	0.1
MO-12	0.3	0.3	0.3	0.2	0.1	0.0	0.2	0.2	0.3	0.1	0.2
MO-13		0.0	0.1	0.2	0.1	0.3	0.1	0.1	0.3	0.2	0.2
MO-14	0.2	0.2	0.2	0.3	0.4	0.2			0.3	0.2	0.2
MO-15			0.1	0.1	0.1						0.1

2003 Evaluation

Table #8 - continued

Summary of Missouri Collections, Located in Field #7

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Average
MO-1	1	4	1	3	4	2	4	2	2	3	2.6
MO-2	1	1	3	1	3	7	7	8	8	5	4.4
MO-3	3	4	8	8	5	4	7	1	4	5	4.9
MO-4	3	3	2	1	1	7	3	5	3	4	3.2
MO-5	3	4	4	4	4	4	8	3	4	4	4.2
MO-6	3					2	5	1	2	1	2.2
MO-7	2	7	3	2	3		6	5	4	4	4.0
MO-8	2	7	4	8	4	1	3	1	1	3	3.4
MO-9	2	4	3	1	5	6	3	1	4	2	3.1
MO-10	1	4	1	1	3	6	3	4	2	6	3.1
MO-11	4	1	1	1	5	8	3	2	1	4	3.0
MO-12	1	1	1	1	4	4	3	3	1	4	2.3
MO-13		1	2	1	3	1	1	3	1	4	1.9
MO-14	2	1	8	2	4	4	4	3	2	4	3.4
MO-15		3		4	3						3.3

Summary of Insect and Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Average
MO-1	2	3	2	3	4	2	5	3	2	3	2.9
MO-2	2	2	4	2	3	4	4	4	5	2	3.2
MO-3	3	3	4	4	4	5	3	2	4	2	3.4
MO-4	2	3	4	2	2	4	4	3	5	2	3.1
MO-5	3	4	3	4	3	2	3	4	4	3	3.3
MO-6	4					2	3	2	3	2	2.4
MO-7	2	3	3	4	4		3	2	4	3	3.1
MO-8	2	4	4	5	3	2	4	3	2	2	3.1
MO-9	3	3	3	2	3	3	4	2	2	3	2.8
MO-10	2	2	2	2	2	4	2	5	3	4	2.8
MO-11	4	2	3	3	3	5	3	3	2	3	3.1
MO-12	2	2	2	4	3	4	2	2	4	2	2.7
MO-13		2	2	4	2	2	2	3	2	3	2.4
MO-14	3	2	3	3	2	4	3	2	3	3	2.8
MO-15		4	4	3	3						3.5

R = Tree was originally MO-6 accession, but was replaced with MO-15 accession

2004 Evaluation

Summary of Missouri Collections, Located in Field #7

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	3.0	3.2	3.2	2.8	3.7	2.4	3.9	dead	dead	2.4	3.1
MO-2	2.8	3.4	dead	3.1	3.6	dead	dead	dead	dead	2.6	3.1
MO-3	3.8	3.4	3.3	3.1	2.8	2.1	2.8	2.7	1.7	2.4	2.8
MO-4	3.6	3.1	2.9	3.3	3.0	1.3	3.1	3.1	3.2	2.8	2.9
MO-5	1.9	2.8	3.1	3.8	3.3	2.8	dead	2.8	2.9	3.3	3.0
MO-6	3.4	R	R	R	R	2.2	3.0	3.0	1.5	3.0	2.7
MO-7	3.9	dead	2.5	2.8	2.1	dead	dead	2.6	dead	3.6	2.9
MO-8	3.2	2.1	3.0	2.2	2.6	2.4	3.0	3.6	2.7	3.2	2.8
MO-9	4.3	2.8	4.2	2.6	2.4	2.5	1.3	4.2	1.6	3.4	2.9
MO-10	3.2	3.4	3.3	2.7	2.2	2.6	2.3	2.5	2.2	dead	2.7
MO-11	2.6	3.1	2.5	2.1	1.7	2.8	2.4	2.4	2.7	3.0	2.5
MO-12	2.4	2.4	3.0	2.0	2.2	2.0	2.7	2.6	2.4	3.2	2.5
MO-13	3.8	3.8	2.8	3.6	2.9	3.1	2.9	2.4	3.2	1.7	3.0
MO-14	3.5	3.3	2.6	2.8	1.7	2.1	dead	2.1	2.9	2.6	2.6
MO-15		3.9	3.1	3.0	3.1						3.3

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	2.4	3.5	3.2	2.5	3.7	2.7	3.8	dead	dead	2.2	3.0
MO-2	3.2	3	dead	3.1	2.3	dead	dead	dead	dead	3	2.9
MO-3	3	2.6	3.5	2.9	2.6	2.3	2.3	2.7	1.7	3	2.7
MO-4	2.6	2.8	2.5	2.6	2.3	1.1	2.9	2.4	3	3	2.5
MO-5	1.8	3	2.3	3.4	3.3	2.5	dead	2.5	2.6	2.6	2.7
MO-6	2.6	R	R	R	R	2	2.1	2.6	1.2	2.4	2.2
MO-7	3.3	dead	2.6	3.5	2.1	dead	dead	2.8	dead	4	3.1
MO-8	2.9	1.9	3.3	1.8	2.2	2.6	2.4	3.1	2.7	3.1	2.6
MO-9	4.2	2.7	3.2	2.3	2.4	2.1	1.3	4.3	1.5	2.9	2.7
MO-10	2.9	2.8	2.7	3	2.2	2.2	2.1	2	2.7	dead	2.5
MO-11	2.6	2.8	2.6	1.8	1.9	2.4	2	2.6	3.4	3.3	2.5
MO-12	1.5	2.8	2.7	2	2.3	2.4	2.7	3	3	4.2	2.7
MO-13	3.4	3.1	2.9	3.6	3	2.9	2.6	2.3	3.2	1.2	2.8
MO-14	2.1	4.4	2.3	3.4	2.1	2.1	dead	2.7	2.5	3.1	2.7
MO-15		3.3	3.3	2.7	3.1						3.1

2004 Evaluation
 Summary of Missouri Collections, Located in Field #7

Table #8 - continued

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	4	4	3	5	2	4	1	dead	dead	5	3.5
MO-2	5	4	dead	3	5	dead	dead	dead	dead	4	4.2
MO-3	3	4	3	4	4	5	4	4	7	4	4.2
MO-4	3	4	4	4	4	8	3	4	3	4	4.1
MO-5	6	5	4	1	3	4	dead	4	4	4	3.9
MO-6	4	R	R	R	R	5	4	4	8	4	4.8
MO-7	3	dead	5	3	6	dead	dead	5	dead	1	3.8
MO-8	3	6	3	6	5	4	4	2	4	3	4.0
MO-9	1	5	3	4	4	5	7	1	7	4	4.1
MO-10	4	4	3	3	4	5	4	6	5	dead	4.2
MO-11	4	4	5	6	6	4	5	4	3	3	4.4
MO-12	6	5	4	6	5	5	3	4	4	2	4.4
MO-13	3	3	4	2	3	3	4	6	4	7	3.9
MO-14	4	4	5	3	6	5	dead	4	4	4	4.3
MO-15		2	3	3	3						2.8

Summary of Insect and Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	3	5	3	2	4	1	2	dead	dead	3	2.9
MO-2	4	2	dead	2	4	dead	dead	dead	dead	2	2.8
MO-3	3	4	3	3	2	2	2	2	3	2	2.6
MO-4	2	2	3	3	2	1	2	2	2	1	2.0
MO-5	1	3	2	2	2	1	dead	2	2	2	1.9
MO-6	4	R	R	R	R	2	2	2	1	2	2.2
MO-7	3	dead	2	2	2	dead	dead	2	dead	1	2.0
MO-8	4	3	3	4	3	1	3	2	3	2	2.8
MO-9	2	2	3	2	1	3	1	3	2	2	2.1
MO-10	3	4	2	2	1	2	1	3	2	dead	2.2
MO-11	3	3	3	2	3	2	2	2	2	2	2.4
MO-12	2	2	4	4	2	3	1	1	3	1	2.3
MO-13	5	2	3	1	2	2	3	2	4	3	2.7
MO-14	6	3	2	3	2	2	dead	2	3	3	2.9
MO-15		2	3	3	1						2.3

R = Tree was originally MO-6 accession, but was replaced with MO-15 accession

Study MOPMC-P-0001-WE, WL											Table #9
Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i> , Michx.											
2005 Evaluation											
Summary of Missouri Collections, Located in Field #7											
Summary of Height (feet)											
Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	4.5	4.0	6.5	3.3		3.6	4.6	Dead	Dead	4.0	4.4
MO-2	3.6	4.5	Dead	4.4		Dead	Dead	Dead	Dead	5.3	4.5
MO-3	5.0	5.0	4.3	4.3		1.9	3.6	3.8	Dead	2.9	3.9
MO-4	4.9	4.4	4.3	4.6		1.0	4.3	4.4	5.3	4.5	4.2
MO-5	3.7	5.0	4.3	4.3		3.8	Dead	3.8	5.8	3.2	4.2
MO-6	6.4	R	R	R		1.0	3.9	4.0	Dead	3.5	3.8
MO-7	6.1	Dead	3.9	3.3		Dead	Dead	3.5	Dead	6.0	4.6
MO-8	6.2	4.0	3.4	3.9		Dead	Dead	4.6	4.8	4.7	4.5
MO-9	6.1	3.7	5.0	3.0		3.2	Dead	5.0	Dead	3.9	4.3
MO-10	4.0	4.2	Dead	3.2		Dead	3.6	3.0	4.0	Dead	3.7
MO-11	3.7	4.0	3.8	Dead		3.2	3.3	3.7	4.7	4.4	3.9
MO-12	3.2	6.7	3.4	3.1		Dead	4.1	4.0	3.8	4.6	4.1
MO-13	4.3	4.2	4.0	4.7		3.6	3.7	3.4	5.0	Dead	4.1
MO-14	4.0	5.0	Dead	4.0		2.8	Dead	4.8	4.0	4.0	4.1
MO-15		5.2	4.3	4.7							4.7
Summary of Spread (feet)											
Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	3.5	4.0	5.0	2.0		3.0	3.3	Dead	Dead	3.5	3.5
MO-2	3.2	3.6	Dead	3.6		Dead	Dead	Dead	Dead	4.0	3.6
MO-3	3.0	3.7	3.6	3.0		1.0	2.9	2.8	Dead	2.5	2.8
MO-4	3.3	3.1	2.7	3.0		1.0	2.8	2.3	4.0	2.9	2.8
MO-5	2.3	3.6	3.0	3.0		2.4	Dead	3.2	1.8	3.0	2.8
MO-6	3.7	R	R	R		1.0	3.0	3.0	Dead	2.9	2.7
MO-7	4.3	Dead	2.6	3.7		Dead	Dead	3.0	Dead	4.4	3.6
MO-8	4.0	3.0	2.8	2.0		Dead	Dead	2.5	3.2	2.7	2.9
MO-9	6.0	3.0	2.4	1.4		1.8	Dead	4.0	Dead	1.5	2.9
MO-10	3.0	3.7	Dead	3.0		Dead	2.7	2.5	2.0	Dead	2.8
MO-11	3.0	3.0	2.5	Dead		2.2	2.6	2.6	3.7	3.0	2.8
MO-12	1.2	3.3	2.5	2.5		Dead	3.0	3.2	3.0	4.0	2.8
MO-13	3.0	3.0	3.0	3.3		2.8	2.8	2.0	4.6	Dead	3.1
MO-14	3.0	2.2	Dead	3.6		2.3	Dead	3.2	2.5	4.0	3.0
MO-15		4.8	3.4	3.3							3.8

2007 Evaluation

Summary of Missouri Collections, Located in Field #7

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	8.3	7.3	10.7	7.9	Gone	7.8	9.9	Dead	Dead	7.5	8.5
MO-2	7.4	7.7	Dead	8.5	Gone	Dead	Dead	Dead	Dead	8.7	8.1
MO-3	10.3	8.8	8.3	8.6	Gone	3.9	8.1	6.6	Dead	7.5	7.8
MO-4	10.4	8.2	7.0	9.1	Gone	Dead	9.8	7.6	10.7	8.1	8.9
MO-5	7.9	9.8	8.9	9.6	Gone	6.7	Dead	7.9	11.1	8.2	8.8
MO-6	10.4	R	R	R	R	Dead	7.4	7.6	Dead	6.0	7.9
MO-7	11.6	Dead	8.2	8.1	Gone	Dead	Dead	6.3	Dead	10.0	8.8
MO-8	9.8	8.0	6.1	8.8	Gone	4.7	Dead	9.5	9.8	9.9	8.3
MO-9	11.0	5.9	7.3	3.0	Gone	5.1	Dead	8.8	Dead	7.5	6.9
MO-10	8.7	7.0	5.5	9.0	Gone	4.4	7.4	6.4	9.2	Dead	7.2
MO-11	6.9	9.1	8.0	4.0	Gone	7.3	8.1	7.4	10.2	9.6	7.8
MO-12	5.5	11.5	6.0	7.0	Gone	4.5	7.5	7.8	8.2	8.0	7.3
MO-13	7.8	9.1	6.6	8.5	Gone	7.6	7.6	8.9	7.7	Dead	8.0
MO-14	6.0	9.8	6.4	8.2	Gone	6.7	Dead	9.0	8.3	7.0	7.7
MO-15		9.6	9.0	8.8	Gone						9.1

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	9.3	6.2	11.5	8.8	Gone	6.0	9.5	Dead	Dead	9.0	8.6
MO-2	7.6	9.0	Dead	10.0	Gone	Dead	Dead	Dead	Dead	7.1	8.4
MO-3	9.0	8.4	10.0	7.6	Gone	3.4	8.2	7.2	Dead	7.3	7.6
MO-4	10.2	8.0	5.7	7.3	Gone	Dead	9.4	6.0	12.0	6.6	8.2
MO-5	6.5	8.0	8.0	9.0	Gone	6.0	Dead	7.0	10.3	6.8	7.7
MO-6	8.0	R	R	R	R	Dead	7.2	8.0	Dead	7.1	7.6
MO-7	11.5	Dead	9.0	7.8	Gone	Dead	Dead	8.6	Dead	9.3	9.2
MO-8	6.9	9.0	6.5	6.0	Gone	4.0	Dead	9.2	11.0	8.0	7.6
MO-9	11.5	8.0	7.0	2.8	Gone	5.0	Dead	9.0	Dead	6.0	7.0
MO-10	6.5	11.5	4.3	6.4	Gone	3.7	7.7	7.0	7.8	Dead	6.9
MO-11	6.0	10.0	6.3	3.4	Gone	6.4	7.8	7.8	10.8	7.3	7.3
MO-12	5.0	8.0	8.0	6.0	Gone	3.7	7.0	7.9	8.2	11.0	7.2
MO-13	7.2	9.0	7.0	8.8	Gone	9.2	6.7	8.0	10.1	Dead	8.3
MO-14	4.8	11.0	6.5	7.5	Gone	5.2	Dead	9.2	6.9	8.2	7.4
MO-15		11.0	9.5	8.0	Gone						9.5

2007 Evaluation
 Summary of Missouri Collections, Located in Field #7

Table 10 - Continued

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	4	5	1	5	Gone	4	4	Dead	Dead	5	4.0
MO-2	5	4	Dead	3	Gone	Dead	Dead	Dead	Dead	3	3.8
MO-3	2	3	3	3	Gone	8	4	5	Dead	5	4.1
MO-4	2	3	5	3	Gone	Dead	3	5	2	4	3.4
MO-5	6	2	4	3	Gone	4	Dead	5	1	4	3.6
MO-6	2	R	R	R	R	Dead	5	4	Dead	5	4.0
MO-7	1	Dead	4	4	Gone	Dead	Dead	6	Dead	2	3.4
MO-8	3	4	6	4	Gone	7	Dead	3	2	2	3.9
MO-9	1	6	4	9	Gone	6	Dead	2	Dead	5	4.7
MO-10	5	5	8	4	Gone	7	5	5	5	Dead	5.5
MO-11	7	3	4	8	Gone	4	3	5	2	3	4.3
MO-12	7	1	6	5	Gone	7	4	4	4	3	4.6
MO-13	6	3	7	3	Gone	4	4	4	4	Dead	4.4
MO-14	7	3	6	3	Gone	6	Dead	3	4	4	4.5
MO-15		2	3	4	Gone						3.0

Summary of Insect and Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	3	4	3	3	Gone	6	7	Dead	Dead	3	4.1
MO-2	3	2	Dead	2	Gone	Dead	Dead	Dead	Dead	4	2.8
MO-3	3	3	2	3	Gone	3	4	3	Dead	3	3.0
MO-4	3	3	3	3	Gone	Dead	3	3	2	2	2.8
MO-5	3	3	4	3	Gone	2	Dead	5	2	3	3.1
MO-6	2	R	R	R	R	Dead	2	3	Dead	3	2.5
MO-7	2	Dead	2	3	Gone	Dead	Dead	6	Dead	3	3.2
MO-8	4	3	4	3	Gone	8	Dead	3	2	3	3.8
MO-9	2	2	2	2	Gone	4	Dead	2	Dead	2	2.3
MO-10	2	4	2	3	Gone	2	3	2	3	Dead	2.6
MO-11	6	3	3	2	Gone	3	2	2	3	4	3.1
MO-12	2	3	3	5	Gone	4	2	2	7	3	3.4
MO-13	3	4	5	4	Gone	2	2	3	4	Dead	3.4
MO-14	4	1	2	3	Gone	3	Dead	3	3	4	2.9
MO-15		2	3	5	Gone						3.3

2007 Evaluation

Summary of Missouri Collections, Located in Field #7

Table #10 - Continued

Summary of Diameter of Trunk at 24 Inch Height

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	8.50	5.75	11.25	5.75	Gone	5.75	9.00	Dead	Dead	6.25	7.5
MO-2	5.25	5.25	Dead	6.75	Gone	Dead	Dead	Dead	Dead	5.75	5.8
MO-3	7.00	6.75	6.75	5.50	Gone	1.75	4.75	4.50	Dead	4.25	5.2
MO-4	8.50	5.75	5.75	6.75	Gone	Dead	8.75	6.25	6.75	5.25	6.7
MO-5	6.25	9.00	5.75	6.50	Gone	6.75	Dead	5.00	8.00	5.75	6.6
MO-6	9.25	R	R	R	R	Dead	7.75	6.50	Dead	5.00	7.1
MO-7	9.25	Dead	6.25	4.75	Gone	Dead	Dead	4.50	Dead	7.50	6.5
MO-8	7.25	7.75	4.25	7.00	Gone	2.25	Dead	6.75	9.25	8.50	6.6
MO-9	13.00	5.00	6.50	1.00	Gone	4.00	Dead	9.00	Dead	5.00	6.2
MO-10	5.50	8.00	2.50	5.00	Gone	2.75	4.25	4.50	5.00	Dead	4.7
MO-11	5.25	6.25	6.50	1.75	Gone	6.25	5.50	4.50	8.50	6.00	5.6
MO-12	3.50	8.00	5.00	3.50	Gone	2.25	6.50	5.50	5.75	8.00	5.3
MO-13	5.75	9.50	5.75	8.00	Gone	7.25	6.50	5.00	5.75	Dead	6.7
MO-14	5.50	6.00	2.50	7.50	Gone	5.25	Dead	9.75	6.50	5.25	6.0
MO-15		9.00	8.00	7.00	Gone						8.0

Summary of Visible Acorn Production

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10
MO-1	no	no	no	no	Gone	no	no	Dead	Dead	no
MO-2	no	no	Dead	no	Gone	Dead	Dead	Dead	Dead	no
MO-3	no	no	no	no	Gone	no	no	no	Dead	no
MO-4	no	no	no	no	Gone	Dead	no	no	no	no
MO-5	no	no	no	no	Gone	no	Dead	no	no	no
MO-6	no	R	R	R	R	Dead	no	no	Dead	no
MO-7	no	Dead	no	no	Gone	Dead	Dead	no	Dead	no
MO-8	no	no	no	no	Gone	no	Dead	no	no	no
MO-9	no	no	no	no	Gone	no	Dead	no	Dead	no
MO-10	no	no	no	no	Gone	no	no	no	no	Dead
MO-11	no	no	no	no	Gone	no	no	no	no	no
MO-12	no	no	no	no	Gone	no	no	no	no	no
MO-13	no	no	no	no	Gone	no	no	no	no	Dead
MO-14	no	no	no	no	Gone	no	Dead	no	no	no
MO-15		no	no	no	Gone					

R = Tree was originally MO-6 accession, but was replaced with MO-15 accession
 Replication #5 was removed and used in another study offsite of the PMC

2008 Evaluation

Summary of Missouri Collections, Located in Field #7

Summary of Height (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	9.3	8.3	11.4	Dead	Gone	8.0	7.8	Dead	Dead	10.0	9.1
MO-2	8.5	8.8	Dead	10.0	Gone	Dead	9.6	Dead	Dead	Dead	9.2
MO-3	11.1	9.8	10.4	9.2	Gone	3.5	8.3	Dead	7.8	9.3	8.7
MO-4	11.0	10.0	Dead	10.7	Gone	Dead	9.4	11.8	10.0	10.8	10.5
MO-5	7.3	10.9	10.2	10.8	Gone	8.0	9.5	11.2	8.0	Dead	9.5
MO-6	12.1	R	R	R	Gone	Dead	7.3	Dead	8.6	8.5	9.1
MO-7	12.3	Dead	9.3	Dead	Gone	Dead	11.1	Dead	8.7	Dead	10.4
MO-8	11.3	9.0	Dead	9.8	Gone	Dead	11.2	10.1	10.6	Dead	10.3
MO-9	11.8	5.8	8.3	2.0	Gone	Dead	8.5	Dead	10.6	Dead	7.8
MO-10	10.0	8.0	Dead	10.2	Gone	Dead	Dead	10.2	Dead	8.7	9.4
MO-11	6.9	9.5	9.2	Dead	Gone	9.0	10.8	11.0	8.8	9.2	9.3
MO-12	5.5	12.1	Dead	Dead	Gone	3.7	8.9	9.2	9.8	9.0	8.3
MO-13	8.8	10.0	8.2	9.3	Gone	9.1	Dead	8.9	9.2	8.5	9.0
MO-14	7.6	11.0	Dead	9.3	Gone	8.1	8.1	9.6	9.1	Dead	9.0
MO-15		11.0	10.3	10.0	Gone						10.4

Summary of Spread (feet)

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	9.1	8.4	12.5	Dead	Gone	6.0	9.0	Dead	Dead	8.8	9.0
MO-2	8.2	8.5	Dead	8.2	Gone	Dead	6.8	Dead	Dead	Dead	7.9
MO-3	11.0	7.7	11.0	8.3	Gone	1.7	7.4	Dead	6.6	7.6	7.7
MO-4	11.2	9.0	Dead	7.0	Gone	Dead	7.0	10.3	7.3	10.2	8.9
MO-5	6.6	9.8	8.0	8.6	Gone	6.2	7.5	9.7	6.4	Dead	7.9
MO-6	11.0	R	R	R	Gone	Dead	7.0	Dead	7.0	7.0	8.0
MO-7	12.0	Dead	7.3	Dead	Gone	Dead	9.2	Dead	7.0	Dead	8.9
MO-8	7.5	7.8	Dead	5.9	Gone	Dead	8.6	10.9	8.0	Dead	8.1
MO-9	11.3	7.5	8.0	1.0	Gone	Dead	6.7	Dead	9.6	Dead	7.4
MO-10	7.1	10.5	Dead	6.4	Gone	Dead	Dead	7.7	Dead	6.9	7.7
MO-11	6.0	9.7	7.2	Dead	Gone	6.0	8.0	10.0	7.5	7.5	7.7
MO-12	6.0	10.7	Dead	Dead	Gone	1.0	10.4	9.2	8.8	7.0	7.6
MO-13	7.0	10.5	8.2	8.6	Gone	7.8	Dead	9.0	6.8	6.6	8.1
MO-14	5.4	11.0	Dead	8.4	Gone	5.4	8.4	7.8	8.7	Dead	7.9
MO-15		10.7	8.8	6.8	Gone						8.8

2008 Evaluation

Summary of Missouri Collections, Located in Field #7

Table 11 - Continued

Summary of Vigor (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	7	6	9	Dead	Gone	6	5	Dead	Dead	7	6.7
MO-2	6	6	Dead	8	Gone	Dead	7	Dead	Dead	Dead	6.8
MO-3	9	7	8	7	Gone	2	6	Dead	5	7	6.4
MO-4	9	6	Dead	7	Gone	Dead	6	7	6	9	7.1
MO-5	4	8	7	7	Gone	7	5	8	5	Dead	6.4
MO-6	9	R	R	R	Gone	Dead	4	Dead	6	6	6.3
MO-7	10	Dead	6	Dead	Gone	Dead	8	Dead	5	Dead	7.3
MO-8	8	5	Dead	5	Gone	Dead	7	8	7	Dead	6.7
MO-9	10	2	5	1	Gone	Dead	5	Dead	8	Dead	5.2
MO-10	7	4	Dead	5	Gone	Dead	Dead	6	Dead	6	5.6
MO-11	3	7	7	Dead	Gone	6	7	8	6	6	6.3
MO-12	3	9	Dead	Dead	Gone	2	6	6	7	6	5.6
MO-13	6	8	4	6	Gone	7	Dead	6	6	6	6.1
MO-14	4	8	Dead	6	Gone	6	6	6	7	Dead	6.1
MO-15		8	8	7	Gone						7.7

Summary of Insect and Disease Resistance (1-9 Rating) 1=Very Good 9=Poor

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	7	7	8	Dead	Gone	6	7	Dead	Dead	7	7.0
MO-2	8	6	Dead	7	Gone	Dead	8	Dead	Dead	Dead	7.3
MO-3	7	8	8	8	Gone	5	7	Dead	6	7	7.0
MO-4	7	6	Dead	7	Gone	Dead	7	7	7	8	7.0
MO-5	7	7	8	8	Gone	7	6	8	6	Dead	7.1
MO-6	8	R	R	R	Gone	Dead	5	Dead	8	6	6.8
MO-7	8	Dead	8	Dead	Gone	Dead	7	Dead	6	Dead	7.3
MO-8	7	6	Dead	6	Gone	Dead	7	7	6	Dead	6.5
MO-9	8	4	9	9	Gone	Dead	8	Dead	7	Dead	7.5
MO-10	8	5	Dead	7	Gone	Dead	Dead	7	Dead	6	6.6
MO-11	4	8	8	Dead	Gone	5	8	8	6	7	6.8
MO-12	8	9	Dead	Dead	Gone	6	7	8	8	5	7.3
MO-13	6	9	7	8	Gone	9	Dead	8	8	6	7.6
MO-14	7	8	Dead	8	Gone	7	7	7	7	Dead	7.3
MO-15		7	8	7	Gone						7.3

2008 Evaluation

Summary of Missouri Collections, Located in Field #7

Table #11 - Continued

Summary of Diameter of Trunk at 24 Inch Height

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	2.75	2.25	4.25	Dead	Gone	2.00	2.25	Dead	Dead	3.00	2.8
MO-2	2.25	2.00	Dead	2.75	Gone	Dead	2.25	Dead	Dead	Dead	2.3
MO-3	3.50	2.50	2.50	2.25	Gone	1.00	2.00	Dead	1.75	1.75	2.2
MO-4	2.75	2.00	Dead	2.50	Gone	Dead	2.00	2.75	2.25	3.50	2.5
MO-5	1.75	3.00	2.50	2.50	Gone	2.50	1.75	3.00	1.75	Dead	2.3
MO-6	3.25	R	R	R	Gone	Dead	1.75	Dead	2.50	2.50	2.5
MO-7	3.50	Dead	2.50	Dead	Gone	Dead	2.75	Dead	2.00	Dead	2.7
MO-8	2.75	2.50	Dead	2.50	Gone	Dead	3.00	3.50	2.50	Dead	2.8
MO-9	4.50	1.25	2.25	0.25	Gone	Dead	1.75	Dead	3.50	Dead	2.3
MO-10	2.25	1.50	Dead	1.75	Gone	Dead	Dead	2.25	Dead	1.50	1.9
MO-11	1.75	2.25	2.25	Dead	Gone	2.25	2.25	3.25	1.75	2.00	2.2
MO-12	1.25	3.25	Dead	Dead	Gone	0.75	3.00	2.25	2.25	2.50	2.2
MO-13	2.25	2.75	2.00	3.00	Gone	2.75	Dead	2.50	2.00	2.50	2.5
MO-14	2.00	2.50	Dead	2.50	Gone	2.25	2.00	2.25	3.25	Dead	2.4
MO-15		4.00	3.00	2.75	Gone						3.3

Summary of Circumference

Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10	Avg.
MO-1	8.64	7.07	13.35	Dead	Gone	6.28	7.07	Dead	Dead	9.42	8.6
MO-2	7.07	6.28	Dead	8.64	Gone	Dead	7.07	Dead	Dead	Dead	7.3
MO-3	10.99	7.85	7.85	7.07	Gone	3.14	6.28	Dead	5.50	5.50	6.8
MO-4	8.64	6.28	Dead	7.85	Gone	Dead	6.28	8.64	7.07	10.99	8.0
MO-5	5.50	9.42	7.85	7.85	Gone	7.85	5.50	9.42	5.50	Dead	7.4
MO-6	10.21	R	R	R	Gone	Dead	5.50	Dead	7.85	7.85	7.9
MO-7	10.99	Dead	7.85	Dead	Gone	Dead	8.64	Dead	6.28	Dead	8.4
MO-8	8.64	7.85	Dead	7.85	Gone	Dead	9.42	10.99	7.85	Dead	8.8
MO-9	14.13	3.93	7.07	0.79	Gone	Dead	5.50	Dead	10.99	Dead	7.1
MO-10	7.07	4.71	Dead	5.50	Gone	Dead	Dead	7.07	Dead	4.71	5.8
MO-11	5.50	7.07	7.07	Dead	Gone	7.07	7.07	10.21	5.50	6.28	7.0
MO-12	3.93	10.21	Dead	Dead	Gone	2.36	9.42	7.07	7.07	7.85	6.8
MO-13	7.07	8.64	6.28	9.42	Gone	8.64	Dead	7.85	6.28	7.85	7.8
MO-14	6.28	7.85	Dead	7.85	Gone	7.07	6.28	7.07	10.21	Dead	7.5
MO-15		12.56	9.42	8.64	Gone						10.2

2008 Evaluation

Summary of Missouri Collections, Located in Field #7

Table #11 - Continued

Summary of Visible Acorn Production


Acc. No.	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Rep 9	Rep 10
MO-1	N	N	YES	Dead	Gone	N	N	Dead	Dead	N
MO-2	N	YES	Dead	N	Gone	Dead	N	Dead	Dead	Dead
MO-3	N	N	N	N	Gone	N	N	Dead	N	N
MO-4	N	N	Dead	N	Gone	Dead	N	N	N	N
MO-5	N	N	N	N	Gone	N	N	N	N	Dead
MO-6	N	R	R	R	Gone	Dead	N	Dead	N	N
MO-7	N	Dead	N	Dead	Gone	Dead	N	Dead	N	Dead
MO-8	N	N	Dead	N	Gone	Dead	N	N	N	Dead
MO-9	N	N	N	N	Gone	Dead	N	Dead	N	Dead
MO-10	N	N	Dead	N	Gone	Dead	Dead	N	Dead	N
MO-11	N	YES	N	Dead	Gone	N	N	YES	N	N
MO-12	N	YES	Dead	Dead	Gone	N	N	N	N	N
MO-13	N	N	N	N	Gone	N	Dead	N	N	N
MO-14	N	N	Dead	N	Gone	N	N	N	N	Dead
MO-15		N	N	N	Gone					

R = Tree was originally MO-6 accession, but was replaced with MO-15 accession
 Replication #5 was removed and used in another study offsite of the PMC

Study MOPMC-P-001 Assembly, Evaluation and Selection of Bur Oak, *Quercus macrocarpa*, Michx.

Plot Layout: 5-11-02

Field 7 - Missouri Collections



**FOURTEEN ACCESSIONS PER REPLICATION. FIVE REPLICATIONS PER ROW.
TWO ROWS TOTALING 10 REPLICATIONS AND 70 PLANTS.**

Study MOPMC-P-001 Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i> , Michx.						
Plot Layout: 5-30-02				Table # 13		
Two plants planted per location			BP=Border Plant (only one plant planted)			
FIELD 6 - Iowa Collections						
		2 ROWS				NORTH ↑
REPS	3 (BP)		3 (BP)		REPS	
12	2		6		1	
	5		2			
	6		7			
	4		1			
	7		3			
11	1		5		2	
	3		4			
	4		5			
	7		3			
	5		7			
10	1		2		3	
	3		6			
	5		1			
	6		4			
	7		3			
9	4		7		4	
	3		1			
	7		4			
	1		6			
	2		2			
8	6		5		5	
	1		4			
	6		2			
	5		5			
	3		3			
7	7		1		6	
	4		6			
	2		7			
	1		2			
	6*		5			
	4		7			
	2		1			
	7		3			
	3		6*			
	5		4			
	2 (BP)		1 (BP)			
*ONLY ONE OF PLANT 6 PLANTED IN THESE LOCATIONS. RAN OUT OF PLANT 6. (BP=Border Plant)						

Study ID Code: MOPMC-T-0105, PA

Study Title – Compatibility Study Using Native Warm Season and Cool Season Grasses with Native Legumes and Forbs

Study Leader: Bruckerhoff, S. B.

Introduction:

Herbaceous plantings using native species are often a single grass species or a mixture of grasses with few legumes or forbs. These types of plantings are typical for forage, conservation cover or even wildlife plantings. Many native forbs and legumes are compatible with native grass species in a native prairie. In a planting using native species it is important to know which ones are most likely to compete with the grasses during the establishment period. Forb and legume seed are more expensive than the grass seed and most plantings lack diversity.

Problem:

There is little to no documented information regarding the compatibility of native warm and cool season grasses with native legumes and forbs in a pasture or range seeding. As a result of the lack of this needed information, the PMC Advisory Committee has directed the PMC to initiate this study.

Objective:

The objective of this study is to determine which native forbs and legumes will establish the easiest and persist the longest with specific native grasses.

Procedure:

Secure seed of the following native cool and warm season grasses, forbs, and legumes.

- **Cool Season Grasses:** Virginia wildrye, Western wheatgrass, Junegrass, and Porcupinegrass.
- **Warm Season Grasses:** Eastern gamagrass, Little bluestem, Big bluestem, Indiangrass, and Switchgrass.
- **Forbs:** Oxeye daisy, Prairie coreopsis, and Grayheaded coneflower.
- **Legumes:** Bush clover, Showy tick trefoil, Purple prairie clover, White prairie clover, Illinois bundleflower, Goat's rue, Wild senna, and Lead plant.

Plots of a native warm season grass mixture, native cool season grass mixture and warm and cool season grass mixture will be established in four replications. Native legumes and forb mixtures will be planted with the grass mixtures. Plots will be planted in the spring and also as winter dormant plantings. All species will also be planted at the same time in the spring and winter except one warm and cool season grass mixture.

Plots will be mowed for weed control during the establishment year. The forage will be removed two to three times a year from half the plot the following years to assimilate rotational grazing.

Discussion:

2001

A site was prepared on the PMC using glyphosate to kill existing vegetation that consisted of mostly annual weedy species. The area was then plowed, disked and planted to an annual covercrop of 80% oats and 20% wheat. Plot composition of species can be seen in Table #1. Seeding rates are 40 pure live seed per square foot with 60% being the grass component and 40% being the forb and legume component.

2002

The winter dormant plots were planted January 8 and 9, 2002 using a plot planter. Seeding depth was one fourth inch for all species except the eastern gamagrass which was planted three fourths to one inch deep. The spring plots were planted May 20 and 21. All seed was planted at a depth of one fourth to one half inch with the exception of eastern gamagrass which again was planted at a depth of three fourths to one inch. All species that required treatment were stratified and/or scarified and inoculated. The plot map of the winter dormant planting is Table #2 and the spring planting is Table #3.

Mowing throughout the summer was the weed control method used. The plots were mowed when vegetation reached six to eight inches. Mowing height was three to four inches.

All plots were evaluated toward the end of the growing season for species composition. Most of the grasses were represented in the plots but in very low densities. Only sideoats gramma and Virginia wildrye appeared in plots in densities in the moderate range. The only legumes/forbs that were identified even at low densities were winter dormant planting Illinois bundleflower, grayheaded coneflower, and prairie coreopsis. Spring planting was Illinois bundleflower and wild senna.

2003/2004

The plots were evaluated for specie density during 2003 and 2004 (see Tables 4 and 5). A winter burn was conducted on all plots in early 2004.

Most species that were planted were identified in the plots although some in very low densities. The specie in the legume/forb mixture showing up in the highest concentration is grayheaded coneflower. Others most consistently found were showy tick trefoil, oxeye daisy, Illinois bundleflower(spring seeding only), and purple prairie clover (spring seeding only).

Most of the grass components of the plots established well but were not very thick stands. The sideoats gramma was high density and the plots with western wheatgrass, Junegrass, and porcupine grass were very poor or none at all.

2005/2006

The plots were again evaluated in 2005 and 2006 (see Tables 4 and 5). The only maintenance to the plots during this period was a burn in March of 2005.

Most of the species planted could be found in the 2006 evaluation although most of the legumes and forbs were in very low densities. Evaluations were visual estimates on a one to nine scale with plants to square feet estimates on the scale. The 2006 evaluation was done by the visual estimate method and also actual counts of three, random one square foot samples per plot. The

table below is actual counts. In comparing the evaluation methods it showed the visual estimates somewhat under estimated the grass species, especially when the densities were high; and over estimated some of the very low densities. Visual estimates were above zero when a few scattered plants could be seen in a plot, but when only three actual counts of one square foot each were made, several had counts of zero.

Species density (stems/sq ft) five years after planting (planted 2002, density at end of 2006)

Winter Dormant Seeding (1/8/2002)				Spring Planting (5/26/2002)			
Grasses	St/sqft	Legumes	St/sqft	Grasses	St/sqft	Legumes	St/sqft
Eastern gamagrass (Plot #8)	13.3	Grayheaded coneflower	2.575	Switchgrass (Plot #4)	24.2	Grayheaded coneflower	1.338
Virginia wildrye (Plot #6)	8.6	Oxeye false sunflower	0.250	Siedoats grama	18.2	Bush clover	0.325
Switchgrass (Plot #1)	7.3	Wild senna	0.113	Virginia wildrye (Plot #1)	15.1	Desmodium Showy tick trefoil	0.275
Eastern gamagrass (Plot #3)	5.8	Desmodium	0.100	Switchgrass (Plot #11)	13.4	Oxeye false sunflower	0.125
Indiangrass	5.5	Illinois bundleflower	0.013	Indiangrass	11.7	Purple prairie clover	0.113
Virginia wildrye (Plot #5)	5.0	Bush clover	0.013	Eastern gamagrass (Plot #3)	11.2	Illinois bundleflower	0.050
Little bluestem	3.9	Purple prairie clover	0	Eastern gamagrass (Plot #8)	8.3	White prairie clover	0
Switchgrass (Plot #4)	2.7	White prairie clover	0	Virginia wildrye (Plot #5)	2.1	Goat's rue	0
Sideoats grama	2.3	Goat's rue	0	Big bluestem	1.4	Lead plant	0
Big bluestem	0.3	Lead plant	0	Little bluestem	0	Prairie coreopsis	0
Western wheatgrass	0	Prairie coreopsis	0	Western wheatgrass	0	Wild senna	0 not planted
Porcupine grass	0			Porcupine grass	0		
Junegrass	0			Junegrass	0		
Total stems/sq ft In 8 plots	54.7		3.064		105.6		2.226

Total stem counts for grass species were higher in the spring planting than the winter dormant planting with 105.6 compared to 54.7. Total stem counts of the forb legume mix were higher in the winter dormant planting with 3.064 compared to 2.226 even though the seed was stratified.

2007

The plots were not burned in 2007 and no stem counts were made. A mid summer (June or July) mowing of plots is planned for 2008.

2008/2009

No evaluations were taken during 2008 or 2009. The plots were mowed once each year in May for weed control and to delay flowering. Mowing was at a height of 8 to 12 inches. A final evaluation for plant composition will be done in 2010 or 2011.

2010

No evaluations were taken from the plots. This study is scheduled for a final evaluation and final report in 2011 or 2012.

8/16/01

Spring planting Randomized complete block 4 Replications
 Winter dormant planting Randomized complete block 4 Replications \1

Plot 1 8' BB , SG Legume Forb Mixture	Plot 2 8' LB , SO Legume Forb Mixture	Plot 3 EG Legume Forb Mixture	Plot 4 T , SG Legume Forb Mixture \1	Plot 5 VW , IG Legume Forb Mixture	Plot 6 VW , WW Legume Forb Mixture	Plot 7 JG , PG Legume Forb Mixture	Plot 8 EG Kura clover	Plot 9 Check
--	--	---	---	--	--	--	--------------------------------	-----------------

WS grass components

big bluestem (BB)
 little bluestem (LB)
 switchgrass (SG)
 sideoats gramma (SO)
 eastern gamagrass (EG)
 Indiangrass (IG)

CS grass components

Virginia wildrye (VW)
 western wheatgrass (WW)
 junegrass (JG)
 porcupine grass (PG)
 timothy (T)

Legume components

bush clover
 purple prairie clover
 white prairie clover
 desmodium
 goat's rue
 wild senna
 Illinois bundleflower
 lead plant

Forb components

oxeye daisy
 grayheaded coneflower
 prairie coreopsis

Fall planted oats covercrop on winter dormancy plantings
 plot size 10' X 20'

\1 This plot will not have a winter dormant planting but rather a late summer planting.

Study MOPMC-PA-0105		Compatability Study										Table #4				
		Winter Planted Plots					Stems/Square Ft Per Plot					Planted 1/8/02				
Plot #1		Plot #2		Plot #3		Plot #4		Plot #5		Plot #5		Table #4				
2003 2004 2005 2006		2003 2004 2005 2006		2003 2004 2005 2006		2003 2004 2005 2006		2003 2004 2005 2006		2003 2004 2005 2006		2003 2004 2005 2006				
WS grass components																
big bluestem (BB)	0.75	1.25	3.00	0.30												
little bluestem (LB)				0.50	0.38	0.40	3.90									
switchgrass (SG)	1.75	1.75	6.80	7.30				0.50	0.50	0.90	2.70					
sidecoats gamma (SO)				1.00	0.38	2.20	2.30									
eastern gamagrass (EG)						0.10										
Indiangrass (IG)								3.00	2.25	9.50	5.80					
												1.75	1.00	9.40	5.50	
CS grass components																
Virginia wildrye (VW)																
western wheatgrass (WW)																
junegrass (JG)																
porcupine grass (PG)																
timothy (T)												0.25	1.75			
Legume components																
bush clover	0.13	0.13	0.10	0.00	0.13	0.25	0.10	0.13	0.25	0.30		0.13	0.10			
purple prairie clover	0.13							0.13								
white prairie clover																
desmodium	0.38	0.38	0.10	0.00	0.25	0.50	0.10	0.25	0.50	0.10	0.20	0.13	0.13	0.38	0.10	
goat's rue																
wild senna	0.38	0.25	0.10	0.10	0.38	0.50	0.10	0.38	0.50	0.10		0.25	0.50	0.38	0.10	
Illinois bundleflower							0.10					0.13				
lead plant																
kura clover																
Forb components																
oxeye daisy	0.50	0.50	0.20	0.10	0.50	0.50	0.30	0.10	0.50	0.20	0.20	0.50	0.50	0.20	0.10	
grayhead coneflower	0.50	2.00	2.00	0.30	0.50	2.00	4.00	0.60	0.50	2.50	3.80	1.90	0.50	2.00	0.50	4.50
prairie coreopsis	0.13					0.50				0.13				0.25		0.25
2003, 2004, 2005 estimated on 1-9 scale																
2006 actual plant count - 3 counts/plot were averaged																
Study MOPMC-PA-0105		Compatability Study										Table #4				

Study ID Code: MOPMC-T-0106, BU

Study Title: Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips

Study Leader: Cordsiemon, R.

Description:

A need has developed out of a three-state technical review committee and approved by the State Conservationists Advisory Committee to evaluate different native cool season grasses and sedges for filter strips.

Grasses and sedges to be considered are Virginia wildrye, *Elymus canadensis*; Canada wildrye, *Elymus canadensis*; Junegrass, *Koeleria cristata*; bluejoint, *Calamagrostis canadensis*; sweet woodreed, *Cinna arundinacea*; river oats, *Uniola latifolia*; longhair sedge, *Carex cosmosa*; Frank sedge, *Carex frankii*; shoreline sedge, *Carex hyalinolepis*; wheat sedge, *Carex atherodes*; raven's foot sedge, *Carex crus-corvi* Shuttlew.; short sedge, *Carex shortina*; hop sedge, *Carex lupulina* Muhl.; crested sedge, *Carex cristatella* Britton; bristle bract sedge, *Carex tribuloides*; and greater straw sedge, *Carex normalis*.

Objective:

There is little to no documented information regarding native cool season grasses and sedges being used in filter strip situations. In an attempt to respond to this lack of information, the PMC has been directed to initiate this study. Depending upon the performance of selected native cool season grasses and sedges in filter strip situations, previous recommendations may change to include those native cool season grasses and sedges performing excellently in this situation.

Discussion:

2001

The study plan for this study was initiated and approved by the State Conservationists' Advisory Committee in August of 2001. Selected field offices in the PMC service area will be contacted in the spring of 2002 requesting their participation in this collection, however everyone is welcomed to participate. One to three collections per state per species are being requested, both seeds and plants. The plants will be grown in the PMC greenhouse and later transplanted in randomized complete blocks. Each block will be one foot wide and five feet long with approximately 30 plants per plot. The spacing of the plants in the blocks will be six inches x six inches.

2002-2003

Collections of native sedges and cool season grasses began on July 2, 2002. The following chart reflects a listing of the collections made as of the time this report was developed. The collection period was extended one more year to make collections of those species that have not been made or those species needing more collections. Fourteen additional collections were made in the state of Missouri and eleven were made in Iowa during 2003. Samples of seed from each different species were planted in the greenhouse to determine the germination percentage. The results will be documented in

the 2004 Annual Technical Report. Field #10 on the PMC has been selected as the site for this study because of the access to water. Two collections of river oats were planted (vegetatively) on September 9, 2002. Both collections were performing with fair to good vigor.

2004

The planting site for this project was changed from Field #10 to Field #7. There is still available water and space. There were two separate wetland cells constructed by using a landscaper in order to simulate a wetland environment. The wetland cells measure 20 feet x 200 feet and are made up of several different individual blocks. The blocks themselves measure 5 feet x 20 feet (refer to Table #2 for map). Collections that did well in the greenhouse were stepped up in plug containers. They were planted in Field #7 on May 3 and were evaluated for percent stand, percent cover, lodging, and survival in late June (refer to Table #3). The west cell contains 17 blocks that include 100 plants per block of a particular species. The east cell contains 27 different collections consolidated into 11 different blocks. These collections were added together because they did not contain 100 plants. All blocks were planted on one foot by one foot spacing. Each collection will be evaluated three times in 2005, (late winter, summer, and fall) for percent stand and cover, lodging, and survival. The cells will be kept fairly damp throughout the growing season and will be treated with a pre-emergent grass herbicide in the sedge plots to help control annual grasses.

2005

Two evaluations were completed (refer to Table # 5) in 2005. Six species were selected based on evaluations for increase plantings. The species selected for increase are larger straw sedge (*Carex normalis*), Crested Sedge (*Carex cristella*), Fox Sedge (*Carex vulpinoidea*), Crowfoot Sedge (*Carex crus-corvi*), Franks Sedge (*Carex frankii*), and Green Bulrush (*Scirpus atrovirens*). The increase plantings are scheduled for January 2006.

2006

On February 15, 2006 the six selected species were planted into production blocks. The blocks were 0.25 acre in size. The plots were planted to cereal rye the season prior to planting and mowed in the fall, plowed in January 2005, disked and rolled prior to planting. Plots were planted in 8" rows with the plot planter on the surface to 1/4" deep. All six species (larger straw sedge (*Carex normalis*), Crested Sedge (*Carex cristella*), Fox Sedge (*Carex vulpinoidea*), Crowfoot Sedge (*Carex crus-corvi*), Franks Sedge (*Carex frankii*), and Green Bulrush (*Scirpus atrovirens*)) were planted at an estimated rate of 40 pure live seed per square foot. See figure 1 for production plantings in field 7A. Plots were sprayed with both Poast (grass herbicide) and 2,4-D (broadleaf herbicide). When the plots were sprayed no sedges were present, weed competition was extreme with white clover dominating the plots. The plots were evaluated several times throughout the growing season, but the selected sedges were not observed. Plans are to replant the same plots with the same species at a later date (April) in 2007. The selected species will be put in the germinator to check germination percent.

2007

On April 25, 2007, the six selected species were planted again in the same areas of field 7 in an attempt to establish production plots of each. Again all six species (larger straw sedge (*Carex normalis*), crested sedge (*Carex cristella*), fox sedge (*Carex vulpinoidea*), crowfoot sedge (*Carex crus-corvi*), franks sedge (*Carex frankii*), and green bulrush (*Scirpus atrovirens*)) were planted at an estimated rate of 40 pure live seed per square foot. Neither Poast, nor 2,4-D were used; instead a new chemical called Stinger was used to control weed competition. Several seedlings were identified, but the plots still struggled to establish. Weed competition again became a problem by late summer and drought conditions did not help in the survival of these wetland species. After speaking with Chris Hoag, Aberdeen PMC in Idaho, it was determined that water would play a huge factor in establishing the different species of sedges.

Seed from the PMC evaluation plots was depleted after planting plots in April 2007. The decision was made to make more collections of the same species from the US Fish and Wildlife Refuge in Annada, Missouri. Candy Chambers, assistant manager of the Clarence Cannon Refuge, assisted in collecting the seed. Collections were made from June to August 2007; some by hand and two were harvested by using the plot combine. Plans for FY 2008 are to plant production plots at the Clarence Cannon Refuge in Annada where the amount and level of water is easily controlled.

Species of Sedges Collected from Clarence Cannon Refuge in 2007		
Scientific Name	Common Name	Amount Collected in Lbs.
<i>Carex vulpinoidea</i>	Fox Sedge	82.9
<i>Carex hyalinolepis</i>	Thinscale Sedge	6.1
<i>Carex frankii</i>	Franks Sedge	0.3
<i>Carex crus-corvi</i>	Crowfoot sedge	1.2
<i>Carex lupulina</i>	Hops Sedge	0.3
<i>Carex cristella</i>	Crested Sedge	0.2
<i>Scirpus atrovirens</i>	Green Bulrush	0.1

2008

In 2008, plans were to establish 5 plots at the Clarence Cannon Refuge in Annada, MO in late June. The flood, that occurred most of the summer, engulfed most of the bottom ground. The 20 x 40 ft. plots would have to wait until November in order to be planted. The five species that will be planted are crested sedge, ravenfoot sedge, Franks sedge, hops sedge, and green bulrush. They will be evaluated for how well they establish and ability to survive in the controlled water levels.

The six plots in field 7 at the PMC were planted in April of 2007 and were inundated with water for 4 weeks during the flood. When the flood waters had receded, it was obvious that Franks, crowfoot, fox, and crested sedge had survived and showed signs of flourishing without the extra competition.

2009

The plots of Franks, crowfoot, fox, and crested sedge were maintained in field 7. There is an emphasis on seed production from these four species so that field testing may begin. Thinscale sedge is showing promise of competing well vegetatively and might be a possible plant material to use in place of Reeds canarygrass. There was very little seed harvested in 2009 and only crested sedge and green bulrush were harvested from field 7. The production plots in field 7 are filling in well and seed production in the future years should be good.

2010

The plots in field 7 of Franks, crowfoot, fox, and crested were maintained. Seed from the production plot of Franks sedge (*Carex frankii*) were harvested and a total of 29 bulk pounds were cleaned. The production field of (*Carex crus-corvi*) crowfoot sedge produced 0.6 pounds of bulk seed.

Table #1
Study MOPMC-T-0106, BU - Collection and Evaluation of Native Cool Season
Grasses and Sedges for Filter Strips

Initial Collections

Scientific Name	Common Name	Collector	City, State	Temp. Acc. No
<i>Carex crus-corvi</i>	Ravens foot sedge	Dennis Shirk	Vienna, MO	MO-1
<i>Carex grayii</i>	Gray sedge	Dennis Shirk	Vienna, MO	MO-2
<i>Carex atherodes</i>	Slough Sedge	Dennis Shirk	Vienna, MO	MO-3
<i>Carex vulpinoidea</i> Michx.	Fox sedge	Dennis Shirk	Vienna, MO	MO-4
<i>Carex vulpinoidea</i> Michx.	Fox sedge	Kaiser & Henry	Elsberry, MO	MO-5
<i>Carex hyalinolepis</i> Steud.	Thinscale scale	Kaiser & Henry	Elsberry, MO	MO-6
<i>Carex crus-corvi</i> Shuttlew	Crowfoot sedge	Kaiser & Henry	Elsberry, MO	MO-7
<i>Carex hyalinolepis</i> Steud.	Thinscale sedge	Paul Freese	Albany, MO	MO-8
<i>Carex vulpinoidea</i> Michx	Fox sedge	Kaiser & Henry	Elsberry, MO	MO-9
<i>Scirpus atrovirens</i>	Green bulrush	Kaiser & Henry	Elsberry, MO	MO-10
<i>Scirpus atrovirens</i>	Green bulrush	Kaiser & Henry	Elsberry, MO	MO-11
<i>Carex frankii</i> Kunth.	Franks sedge	Paul Freese	Albany, MO	MO-12
<i>Carex lupulina</i> Muhl.	Hop sedge	Raleigh Redman	Warrensburg, MO	MO-13
<i>Carex grayii</i>	Gray's sedge	Raleigh Redman	Warrensburg, MO	MO-14
<i>Carex hyalinolepis</i> Steud.	Thinscale sedged.	Raleigh Redman	Warrensburg, MO	MO-15
<i>Carex frankii</i> Kunth	Frank's sedge	Lingwall & Ellis	Ralls Co., MO	MO-17
<i>Carex crus-corvi</i>	Crowfoot sedge	Lingwall & Ellis	Ralls Co., MO	MO-18
<i>Carex hyalinolepis</i> Stued.	Thinscale sedge	Lingwall & Ellis	Ralls Co., MO	MO-19
<i>Carex frankii</i> Kunth	Frank's sedge	Raleigh Redman	Warrensburg, MO	MO-20
<i>Chasmanthium latifolium</i>	River oats	J. Kaiser	Troy, MO	MO-21

Table # 1-Study MOPMC-T-0106, BU - cont.

Scientific Name	Common Name	Collector	City, State	Temp. Acc. No
<i>Chasmanthium latifolium</i>	River oats	Travis Dinsdale	Springfield, MO	MO-22
<i>Chasmanthium latifolium</i>	River oats	Rodney Doolen	Puxico, MO	MO-23
<i>Chasmanthium latifolium</i>	River oats	J. Kaiser	Troy, MO	MO-24
<i>Chasmanthium Latifolium</i>	River oats	William Brouk	Benton, MO	MO-25
<i>Carex crus-corvi Shuttlew</i>	Ravensfoot sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-26
<i>Carex shartina</i>	Short sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-27
<i>Carex</i>	Shoreline sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-28
<i>Carex hyalinoepris</i>	Thinscale sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-29
<i>Carex vulpinoidea Michx.</i>	Fox sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-30
<i>Carex crus-corvi Shuttlew</i>	Ravensfoot sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-31
<i>Carex vulpinoidea Michx</i>	Fox sedge	J. Kaiser J. Henry	BK Leach Wildlife Area	MO-32
<i>Scirpus atrovirens</i>	Green bulrush	Aaron Jeffries	Howard Co, MO	MO-33
<i>Carex frankii</i>	Frank's sedge	Aaron Jeffries	Howard Co, MO	MO-34
<i>Carex lupulina</i>	Hop sedge	Aaron Jeffries	Howard Co, MO	MO-35
<i>Carex shortina</i>	Short sedge	Aaron Jeffries	Howard Co, MO	MO-36
<i>Scirpus acutus</i>	Hard-stemmed bulrush	Aaron Jeffries	Howard Co, MO	MO-37
<i>Scirpus atrovirens</i>	Green bulrush	Paul Frese	Gentry Co, MO	MO-38
<i>Chasmanthium latifolium</i>	River oats	Travis Dinsdale	Webster Co, MO	MO-39
<i>Carex hyalinoepris Steud.</i>	Thinscale sedge	Dave Hiatt	Martinsville, IL	IL-1


Table 1-Study MOPMC-T-0106, BU - cont.

Scientific Name	Common Name	Collector	City, State	Temp. Acc. No
<i>Carex lupulina</i> Muhl.	Hop sedge	Christine Talige	Fairfield, IA	IA-1
<i>Carex cristatella</i> Britton	Crested sedge	Tim Meyer	Williamsburg, IA	IA-2
<i>Carex cristatella</i> Britton	Crested sedge	Tim Meyer	Williamsburg, IA	IA-3
<i>Carex vulpineidea</i>	Fox sedge	Tim Meyer	Williamsburg, IA	IA-4
<i>Scirpus atrovirens</i>	Green bulrush	Tim Meyer	Williamsburg, IA	IA-5
<i>Juncus interior</i> Weigand	Inland rush	Tim Meyer	Williamsburg, IA	IA-6
<i>Calamagrostis Canadensis</i>	Bluejoint	Tim Meyer	Williamsburg, IA	IA-7
<i>Scirpus atrovirens</i>	Green bulrush	Tim Meyer	Williamsburg, IA	IA-8
<i>Carex normalis</i>	Larger straw sedge	Tom Hurford	Atlantic, IA	IA-9
<i>Carex tribuloides</i>	Bristle bract sedge	Tom Hurford	Atlantic, IA	IA-10
<i>Carex normalis</i>	Larger straw sedge	Tom Hurford	Atlantic, IA	IA-11
<i>Scirpus atrovirens</i>	Green bulrush	Tom Hurford	Atlantic, IA	IA-12

Sedge, Rush, and Cool Season Grass Plot - Field #7

Table #2

North



Planted between 5/3/04 & 5/10/04
 Each individual plot is 5 foot by 20 foot.
 Each plant is planted 1 foot apart in a 5 x 20 foot block.
 Plots on the west side, were planted with a complete 100 plant block. Plots on the east side are made up of partial collections.

Study ID Code: MOPMC-T-0106, BU			Table #3
Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips			
MISSOURI COLLECTIONS			
Collection	Common Name	Scientific Name	# of Plants
MO-1	Crowfoot Sedge	Carex crus-corvi	100 Plants
MO-3	Slough Sedge	Carex obnupta	7 plants
MO-4	Fox Sedge	Carex vulpinoidea	100 Plants
MO-5	Fox Sedge	Carex vulpinoidea	100 Plants
MO-6	Thinscale Sedge	Carex hyalinolepis	8 Plants
MO-7	Crowfoot Sedge	Carex crus-corvi	47 Plants
MO-9	Franks Sedge	Carex frankii	45 Plants
MO-10	Green Bulrush	Scirpus atrovirens	100 Plants
MO-11	Green Bulrush	Scirpus atrovirens	100 Plants
MO-12	Franks Sedge	Carex frankii	100 Plants
MO-13	Hop Sedge	Carex lupulina	25 Plants
MO-15	Thinscale Sedge	Carex hyalinolepis	3 Plants
MO-16	Franks Sedge	Carex frankii	75 Plants
MO-17	Franks Sedge	Carex frankii	76 Plants
MO-18	Crowfoot Sedge	Carex crus-corvi	11 Plants
MO-19	Thinscale Sedge	Carex hyalinolepis	3 Plants
MO-20	Franks Sedge	Carex frankii	54 Plants
MO-21	River Oats	Chasmathium latifolium	76 Plants
MO-22	River Oats	Chasmathium latifolium	100 Plants
MO-23	River Oats	Chasmathium latifolium	100 Plants
MO-24	River Oats	Chasmathium latifolium	100 Plants
MO-25	River Oats	Chasmathium latifolium	100 Plants
MO-26	Crowfoot Sedge	Carex crus-corvi	6 Plants
MO-27	Bottlebrush Sedge	Carex comosa	100 Plants
MO-28	Thinscale Sedge	Carex hyalinolepis	9 Plants
MO-29	Thinscale Sedge	Carex hyalinolepis	13 Plants
MO-30	Fox Sedge	Carex vulpinoidea	100 Plants
MO-31	Crowfoot Sedge	Carex crus-corvi	11 Plants
MO-32	Fox Sedge	Carex vulpinoidea	100 Plants
MO-35	Hop Sedge	Carex lupulina	19 Plants
MO-36	Squarrose Sedge	Carex squarrosa	6 Plants
MO-37	Hard-stemmed Sedge	(hard-stemmed bulrush) Schoenoplectus acutus	18 Plants
MO-39	River Oats	Chasmathium latifolium	100 Plants

Table 3 - continued			
	<u>IOWA COLLECTIONS</u>		
<u>Collection</u>	<u>Common Name</u>	<u>Scientific name</u>	<u># of Plants</u>
IA-1	Hop Sedge	Carex lupulina	23 Plants
IA-2	Crested Sedge	Carex cristatella	52 Plants
IA-3	Crested Sedge	Carex cristatella	100 Plants
IA-4	Fox Sedge	Carex vulpinoidea	100 Plants
IA-5	Green Bulrush	Scirpus atrovirens	100 Plants
IA-6	Inland Rush	Juncus interior	17 Plants
IA-7	Bluejoint	Calamagrostis canadensis	23 Plants
IA-9	Larger Straw Sedge	Carex normalis	76 Plants
IA-8	Green Bulrush	Scirpus atrovirens	38 Plants
IA-11	Larger Straw Sedge	Carex normalis	76 Plants
IA-12	Green Bulrush	Scirpus atrovirens	60 Plants
Shoreline sedge is the same as thinscale sedge			
Crowfoot sedge is the same as ravenfoot sedge			

Study ID Code: MOPMC-T-0106, BU

Table #4

Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips

Sedge, Cool Season Grass, and Bulrush Evaluation

DATE: 6/22/04

Collection #	Name	Number of Plants	Percent Stand	Percent Cover	Lodging (1-9 Rating)	Notes
MO-1	Crowfoot Sedge	100	100	20	1	
MO-4	Fox Sedge	100	100	20	1	
MO-5	Fox Sedge	100	100	20	1	
MO-10	Green Bulrush	100	100	15	1	
MO-11	Green Bulrush	100	100	15	1	
MO-12	Franks Sedge	100	100	25	1	
MO-22	River Oats	100	80	5	1	
MO-23	River Oats	100	100	5	1	
MO-24	River Oats	100	95	5	1	
MO-25	River Oats	100	95	5	1	
MO-27	Bottlebrush Sedge	100	100	20	1	
MO-30	Fox Sedge	100	100	15	1	
MO-32	Fox Sedge	100	100	15	1	
MO-39	River Oats	100	80	5	1	
IA-3	Crested Sedge	100	100	30	1	
IA-4	Fox Sedge	100	100	15	1	
IA-5	Green Bulrush	100	100	20	1	
MO-13	Hop Sedge	25	100	25	1	
MO-16	Franks Sedge	75	100	25	1	
MO-9	Franks Sedge	45	100	25	1	
MO-20	Franks Sedge	54	100	20	1	
MO-29	Thinscale Sedge	13	100	15	1	
MO-6	Thinscale Sedge	8	100	20	1	
MO-15	Thinscale Sedge	3	66	15	1	
MO-19	Thinscale Sedge	3	33	15	1	
MO-18	Crowfoot Sedge	11	100	15	2	
MO-7	Crowfoot Sedge	47	100	20	2	
MO-17	Franks Sedge	76	100	10	1	
MO-37	Hard-stemmed Sedge	18	100	10	1	
MO-35	Hop Sedge	20	100	20	1	
MO-26	Crowfoot Sedge	6	100	10	2	
MO-31	Crowfoot Sedge	11	100	25	1	
MO-28	Thinscale Sedge	9	100	10	1	
MO-36	Squarrose Sedge	6	85	15	1	
MO-3	Slough Sedge	7	100	15	1	
IA-12	Green Bulrush	60	100	10	1	

Table #4 - continued

Collection #	Name	Number of Plants	Percent Stand	Percent Cover	Lodging (1-9 Rating)	Notes
IA-8	Green Bulrush	38	100	10	1	
IA-9	Larger Straw Sedge	76	100	20	1	
IA-11	Larger Straw Sedge	76	100	10	1	
IA-1	Hop Sedge	23	100	15	1	
IA-2	Crested Sedge	52	100	25	1	
IA-7	Bluejoint	21	92	15	1	
IA-6	Inland Rush	16	96	15	1	
MO-21	River Oats	76	95	5	1	

1-9 Rating 1 = No Lodging 9 = Severe Lodging

Study MOPMC-T-0106, BU - Collection and Evaluation of Native Cool Season Grasses and Sedges

Table #5

2005 Evaluation Averages For Each Species

Name	Collection #	% Stand	% Cover	Lodging	Vigor
Bluejoint	IA-7	62.5	47.5	4	6
Bottlebrush Sedge	MO-27	100	82.5	4	6
Crested Sedge	IA-2	100	54.5	5	1
Crested Sedge	IA-3	100	85	5.5	1
Crowfoot Sedge	MO-1	100	90	3	1
Crowfoot Sedge	MO-7	100	95	4.5	2
Crowfoot Sedge	MO-18	100	90	4	2
Crowfoot Sedge	MO-26	100	90	5	3
Crowfoot Sedge	MO-31	100	95	7	2
Fox Sedge	IA-4	100	72.5	4.5	3
Fox Sedge	MO-4	99	95	5	2
Fox Sedge	MO-5	100	87.5	5	2
Fox Sedge	MO-30	100	87.5	5.5	3
Fox Sedge	MO-32	100	80	4.5	4
Franks Sedge	MO-9	100	90	3	3
Franks Sedge	MO-12	100	85	2.5	1
Franks Sedge	MO-16	100	92.5	3	3
Franks Sedge	MO-17	98.5	75	3	5
Franks Sedge	MO-20	100	85	3	3
Green Bulrush	IA-5	100	57.5	4.5	4
Green Bulrush	IA-8	100	87.5	5	4
Green Bulrush	IA-12	100	72.5	4.5	4
Green Bulrush	MO-10	99.5	52.5	3	5
Green Bulrush	MO-11	100	50	3	6
Hop Sedge	IA-1	100	92.5	7	2
Hop Sedge	MO-13	100	85	2	3
Hop Sedge	MO-35	100	90	3	2
Inland Rush	IA-6	95		3.5	4
Larger Straw Sedge	IA-9	100	95	5.5	2
Larger Straw Sedge	IA-11	100	92.5	5	1
River Oats	MO-21				9
River Oats	MO-22	70	15	1	9
River Oats	MO-23	97	25	1	9
River Oats	MO-24	95	20	1	9
River Oats	MO-25	90	20	1	9
River Oats	MO-39	85	15	1	9
Slough Sedge	MO-3	100	62.5	3.5	5

Study MOPMC-T-0106, BU - Collection and Evaluation of Native Cool Season Grasses and Sedges

Table #5 - continued

2005 Evaluation Averages For Each Species

Name	Collection #	% Stand	% Cover	Lodging	Vigor
Squarrose Sedge	MO-36	100	50	5	5
Thinscale Sedge	MO-6	100	80	4.5	1
Thinscale Sedge	MO-15	100	75	2	2
Thinscale Sedge	MO-19	100	65	2	5
Thinscale Sedge	MO-28	100	70	2	4
Thinscale Sedge	MO-29	100	77.5	1.5	2

Lodging

1 = No Lodging 9= Severe Lodging

Vigor

1= Highly Vigorous 9= Low Vigor

Study: MOPMC-P-0613-PA, WL

Study Title: Evaluation and Release of a Shade Tolerant Big Bluestem, *Andropogon gerardii*, L., for Silvopasture

Study Leaders: Van Sambeek, J., D. Wallace, G. Garrett, S. Bruckerhoff

Introduction:

Big bluestem is one of the most widespread important forage grasses of the North American tallgrass prairie region. It is usually associated with one or more of the other three dominant species; Indiangrass, switchgrass, and little bluestem. Big bluestem occurs on sub-irrigated lowlands, nearly level to gently undulating glacial till plains, overflow sites, level swales and depressions, residual and glacial uplands, and stream terraces and bottomlands along rivers and tributaries. The abundant, leafy forage is palatable to all classes of livestock.

The Elsberry PMC initiated a big bluestem collection from the Ozark region (Missouri, Arkansas, and Oklahoma) in 1988 to develop an improved big bluestem that would be better adapted to this region. The collection effort resulted in an assembly of 370 big bluestem accessions and three releases, OZ70 germplasm, a selected release for forage; Refuge germplasm, a medium height selection for buffers, filters and wildlife; and OH370 germplasm, a source identified release for diversity.

Problem:

There is limited information available on species selected for forage to be primarily used for savannas and silvopasture systems throughout Missouri.

Objective:

An existing PMC big bluestem collection displays an amazing range of morphological and phonological characteristics – color, plant height, blade width, stem erectness, rust resistance, spring emergence, anthesis, etc. This collection has never been evaluated for shade attributes that might be valuable in silvopasture systems or with savanna restoration efforts. This study will select and evaluate for forage production in relation to shade tolerance.

Cooperators:

The USDA Natural Resources Conservation Service, Plant Materials Center and the University of Missouri, Columbia, Horticulture and Agroforestry Research Center, HARC.

Procedure:

Randomized complete block design with three replications, five shade levels: full sun, 30% shade, 55% shade, 78% shade with sunflecks, and 80% shade.

Twenty-two accessions of big bluestem were selected based on original collection site descriptions and phenotypic characteristics such as leaf width, and chlorophyll content. These collections came from previous work (Study 29I097G) at the Elsberry and Arkansas PMC's and were assembled in 2005 at the University of Missouri Agroforestry Center (HARC). Additional species being evaluated are eastern gamagrass, Canada wildrye, riveroats, cluster fescue, and tall fescue. Plants were started in the greenhouse and planted in replicated plots the spring 2006.

Forty-five pots for each accession were randomized within each treatment. Above ground dry weight, leaf weight, and forage quality (acid detergent fiber, neutral detergent fiber, crude protein) will be collected for each accession.

Discussion:

2005/2006

Key Findings:

All big bluestem accessions exhibited growth reductions in harvested seasonal biomass (summer plus fall harvest in 2005) under reduced light, although four accessions had produced greater biomass at 70% of full sunlight than under full sunlight (Table 1). Response of most accessions closely approximated a linear relationship between harvested biomass and percentage reduction in light from full sunlight. When fitted to the equation $Y = MX + B$ where Y is g/pot harvested biomass and X is percent reduction from full sunlight, M approximates decrease in biomass per percent reduction in full sunlight and B approximates yield under full sun. We hypothesize the best accessions for most agroforestry practices will exhibit a compromise between relatively low values for M and relatively high values for B such as exhibited by PMC-6925, a high producing shade tolerant accession, compared to PMC-6974, an accession that did poorly under increasing shade (fig. 1). Yield data was also fitted to a second order polynomial and first derivative solved to estimate the percent of full sunlight produced maximum yields (estimated peak biomass production was at 76 % and >100% of full sunlight for PMC-6925 and PMC-6974, respectively).

Analysis of summer 2005 big bluestem biomass for forage quality showed highly significant interactions for Crude Protein (CP), Acid Detergent Fiber (ADF), and Relative Feed Value (RFV) under differing shade levels with and overall increase in CP and ADF and an overall decrease in RFV as shade levels increased (Table 2). Neutral Detergent Fiber (NDF) and Total Digestible Nutrients (TDN) also showed significant and very significant responses, respectively, to increasing shade levels.

Table 1.—Seasonal biomass production in 2005 for 21 accessions of big bluestem under five light regimes, coefficients for linear regression for each accession, and calculated percentage of full sunlight for maximum biomass yield.

State of origin	PMC number	2005 combined summer and fall biomass (g/pot) under five light regimes					Reg. coeff. for mx + b		Max. yield % fs
		Full sun	70%	45%	20%	Sunfleck	b	x	
AR	6967(02)	71	70	51	31	38	77	-0.51	>100
MO	6832(04)	85	81	67	39	32	92	-0.66	89
MO	6812(05)	75	85	61	37	44	86	-0.52	80
MO	6885(06)	96	86	79	32	42	104	-0.76	88
AR	6896(07)	94	82	73	43	50	98	-0.61	95
AR	6972(08)	88	81	78	42	52	94	-0.53	85
MO	6807(10)	91	76	64	47	46	92	-0.57	>100
AR	6974(12)	94	73	53	23	27	97	-0.89	>100
AR	6902(13)	90	80	58	32	46	94	-0.68	>100
MO	6802(14)	110	89	89	52	60	112	-0.66	>100
AR	6905(15)	88	75	67	40	45	91	-0.58	>100
OK	7049(16)	103	92	74	54	53	107	-0.65	>100
MO	6741(17)	87	93	77	52	56	96	-0.47	79
AR	6925(19)	91	96	89	65	58	100	-0.41	76
MO	6838(21)	96	96	78	45	57	105	-0.64	86
MO	6704(22)	73	67	57	37	35	77	-0.48	98
AR	6935(24)	85	64	60	26	37	86	-0.65	>100
OK	7039(25)	78	72	57	52	53	79	-0.34	>100
MO	OZ-70(26)	88	62	54	18	26	89	-0.83	>100
MO	Rountree	85	88	79	55	56	92	-0.41	78
OK	7007(29)	67	54	32	17	20	68	-0.62	>100

Figure 1.—Individual pot biomass in 2005 for shade tolerant PMC-6925 (squares) and intolerant PMC-6974 (diamonds) and fitted linear and quadratic responses as a function of full sunlight.




Table 2.– Summer 2005 forage quality mean and standard deviation (SD) averaged across 21 big bluestem (BBS) under five light regimes for percent crude protein (CP=true protein plus non-protein nitrogen), acid detergent fiber (ADF=highly indigestible fiber), neutral detergent fiber (NDF=cell wall or structural components), total digestible nutrients (TDN=111.8 + 0.95 CP - 0.36 ADF- 0.7 NDF), and relative feed value (RFV=(%DDM x %DMI)/1.29 or relative to full bloom alfalfa set at 100).

Light regime	CP (%)		ADF (%)		NDF (%)		TND (%)		RFV	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Full sun	6.2	1.0	36.0	3.2	63.5	2.6	60.2	3.3	89.4	6.8
70 % of full sun	6.2	1.5	39.2	3.6	65.7	3.3	57.6	4.0	83.0	8.0
45 % of full sun	6.9	1.9	40.0	3.9	65.9	3.4	57.7	4.4	81.8	8.1
20% of full sun	8.7	1.9	40.8	3.0	66.4	2.4	58.9	3.9	80.2	5.9
22% sunfleck	8.3	1.9	40.2	2.8	65.6	2.5	59.3	3.8	81.9	5.7
Sign: Shade	***		***		*		**		***	
5% LSD	0.5		1.2		1.5		1.3		2.5	
Interaction sign.: Shade x Acces.	***		ns		ns		**		ns	

2007

Key Findings:

Analysis of 2006 forage quality data compared to 2005 results (Table 3) showed that general trends for 2006 were similar to 2005. TDN and RFV values decreased as shade levels increased; while crude protein, ADF and NDF, values increased for all combined accessions.

Table 3. Seasonal forage quality in 2005 and 2006 for combined accessions of big bluestem under four light regimes.

Light regime	CP (%)		ADF (%)		NDF (%)		TDN (%)		RFV	
	2005 Mean	2006 Mean								
Full sun	6.2	5.0	36.0	35.6	63.5	62.9	60.2	59.7	89.4	90.4
70 % of full sun	6.2	4.4	39.2	36.2	65.7	64.0	57.6	58.2	83.0	88.4
45 % of full sun	6.9	4.3	40.0	37.7	65.9	65.3	57.7	56.6	81.8	85.0
20% of full sun	8.7	4.6	40.8	39.1	66.4	66.5	58.9	55.6	80.2	82.0
5% LSD	0.5	0.4	1.2	1.1	1.5	1.4	1.3	1.3	2.5	2.5

Substantial variation exists among the 21 big bluestem accessions for total forage production (sum of boot-stage plus fall harvest) in both 2005 and 2006 (Table 4). Both boot-stage and fall biomass yields for big bluestem plants were substantially lower in 2006 than in 2005 when the plants were first established. Examination of the root systems after the 2006 growing season indicated plants were heavily root-bound with many roots circling the inside of the 2-gallon pots. The number of days to mid-boot or late-boot (just before inflorescence emerges from the sheath of a flag leaf) showed greater variability among the big bluestem accessions during the 2006 growing season than the 2005 growing season. In addition, four accessions (PMC-6832, 6896, 6295, and 7007) in 2006 had plants that failed to flower especially at the lowest light levels.

Table 4. Seasonal biomass production in 2005 and 2006 for 21 accessions of big bluestem under five light regimes with sunfleck treatment yielding 22% of full sunlight.

State of origin	PMC number	2005 combined summer and fall biomass (g/pot) under five light regimes					2006 combined summer and fall biomass (g/pot) under five light regimes				
		Full sun	70%	45%	20%	Sun fleck	Full sun	70%	45%	20%	Sun fleck
AR	6967(02)	71	70	51	31	38	28	23	24	29	28
MO	6832(04)	85	81	67	39	32	38	32	26	27	31
MO	6812(05)	75	85	61	37	44	31	33	28	32	34
MO	6885(06)	96	86	79	32	42	41	45	41	27	34
AR	6896(07)	94	82	73	43	50	21	19	21	17	22
AR	6972(08)	88	81	78	42	52	19	22	31	29	24
MO	6807(10)	91	76	64	47	46	31	25	35	36	38
AR	6974(12)	94	73	53	23	27	67	56	51	42	43
AR	6902(13)	90	80	58	32	46	24	24	34	31	27
MO	6802(14)	110	89	89	52	60	58	43	49	52	50
AR	6905(15)	88	75	67	40	45	20	29	26	29	28
OK	7049(16)	103	92	74	54	53	23	33	39	46	48
MO	6741(17)	87	93	77	52	56	24	30	30	37	30
AR	6925(19)	91	96	89	65	58	31	37	45	32	43
MO	6838(21)	96	96	78	45	57	30	30	39	38	43
MO	6704(22)	73	67	57	37	35	31	35	40	39	34
AR	6935(24)	85	64	60	26	37	26	28	40	26	30
OK	7039(25)	78	72	57	52	53	29	19	24	27	22
MO	OZ70(26)	88	62	54	18	26	27	37	41	28	25
MO	Rountree	85	88	79	55	56	37	47	46	42	41
OK	7007(29)	67	54	32	17	20	30	31	33	31	32
Means	-----	87	79	67	40	44	29	32	33	31	31

2010

Doug Wallace, study leader, took a different position within NRCS, therefore no data has been reported for this study in the last two years. Data has been collected and the Elsberry PMC personnel is working with Jerry Van Sambeek, Forest Service, to compile the data and it will be reported in the 2011 ATR.

Study: MOPMC-T-0716-BF

Study Title: In-Field Weathering Effects on Biomass Yield and Biofuel Quality of Warm Season Grasses.

Study Leaders: Cordsiemon, R., Douglas, J., and Bruckerhoff, S.

Introduction:

Dedicated energy crops grown for direct combustion or gasification to generate electricity; ethanol production for transportation fuel; or thermochemical conversion into other by products, require different biofuel quality (McLaughlin et al., 1996). For direct combustion or gasification, biofuel quality needs to have low concentration of alkali metals, especially potassium, low levels of total ash-forming materials, and higher calcium content to mitigate slagging and fouling of conventional boilers (Baxter et al., 1998, Miles et al., 1996). Conversely, for ethanol fermentation and thermochemical conversion by gasification, the biofuel quality must have low moisture, nitrogen, and ash content and a high concentration of lignocellulose in the biomass (McKendry, 2002).

Time and frequency of harvest play a major role in biofuel quality. Nitrogen and ash content of 'Alamo' switchgrass (*Panicum virgatum* L.), 'Highlander' eastern gamagrass (*Tripsacum dactyloides* [L.] L.) and caucasian bluestem (*Bothriochloa baldhii* [Retz.] S.T. Blake) were reduced in a single fall harvest regime compared to a two harvest regime, which consisted of an early summer and early fall harvest in Mississippi (Grabowski et al., 2004). Deferring native grass species for two years in Canada produced the greatest biomass yield with highest cellulosic content for ethanol production as compared to three and four year deferral period, which seem to favor livestock forage for beef cattle (Jefferson et al., 1999). Exposing standing biomass to natural field weathering has shown to be advantageous for achieving biofuel quality. Delaying switchgrass harvest from the fall to spring in Pennsylvania reduced moisture and mineral content to a level suitable for all biofuel conversion systems (Alder et al., 2006). However, these authors reported lower yields due to loss of leaves and panicles during the winter months, and difficulty during the harvesting operations because of the brittleness of the biomass and lodging from snowfall. In contrast, Boe and Lee (2005) found little to no difference in biomass yield in the northern Great Plains from fall to spring, but clipping height was adjusted from 10 cm in the fall to near ground level in the spring, resulting in higher yields due to heavier stems near the base of the plant.

Problem:

There is limited information available on biofuel quality and biomass yield of warm season grasses in relation to effects of weathering in the field throughout the winter.

Objective:

Cultivars/selections of warm season grasses will be compared in replicated plots to evaluate the effects of fall, winter and early spring harvest on dry matter production and biofuel quality.

Cooperators:

Steve Bruckerhoff, USDA-NRCS, Elsberry, Missouri
 Bill Kuentler, USDA-NRCS, Fort Worth, Texas
 Jerry Lemunyon, USDA-NRCS, Fort worth, Texas
 Jim Kiniry, USDA-ARS, Temple, Texas
 Brian Baldwin, Mississippi State University, Starkville, Mississippi
 Ray Cragar, USDA-NRCS, Knox City, Texas
 Joel Douglas, USDA-NRCS, Fort Worth, Texas

Procedure:

Randomized split plot design with four replications.

Species/cultivars or selections (see list below) from appropriate sources will be planted by seed or propagules (*miscanthus*) into plots containing seven rows with 36” spacing and 20 feet long. Interior five rows will be clipped for biomass and grab samples will be taken for fuel quality estimates. Outside rows will be border rows. Plots will be planted April/May 2007. Irrigation will be applied as needed during the establishment year only. Timing and rate of fertilizer amendments will be determined at a later date.

Species/cultivars to be tested	Seeding rate(seeds/row ft--#/ac
‘Cave-in-rock’ switchgrass	50 PLS/row ft - 2.80#PLS/ac
‘Kanlow’ switchgrass	50 PLS/row ft - 2.80#PLS/ac
‘Rumsey’ Indiangrass	50 PLS/row ft - 4.16#PLS/ac
9083274 big bluestem (MOPMC)	50 PLS/row ft - 5.03#PLS/ac
‘Alamo’ switchgrass	50 PLS/row ft - 2.8#PLS/ac
<i>Miscanthus</i> (sterile)	3 ft spacing within and between rows

Supporting evaluations

Stems or plants per row foot at end of first growing season

Harvest Treatments

A 14’ 6” swath will be clipped from the center of each plot beginning at seed maturity (2008-2010) and every six weeks until spring (2009-2011).

Harvest Dates (approximate)
September 15
November 1
December 15
February 1
March 15

Biomass production and biofuel quality of N, lignin, ADF, NDF, ash (total) caloric value, Ca, Mg, S, P, K and gross energy will be determined at each harvest date.

This study is being replicated at four locations, Elsberry, Missouri; Knox City, Texas; Temple, Texas and Starkville, Mississippi.


Discussion:

2007

The study site was fallowed in 2006 using roundup and tillage. The site was then tilled and rolled in 2007 prior to establishment of the plots April 19, 2007. Preemergant chemical weed control was used after planting using atrazine on the switchgrass, plateau on the big bluestem and Indiangrass, and prowl on the *miscanthus*. The study was planted using a randomized split plot design with four replications. (See table #1.) The switchgrass plots encountered herbicide damage and were replanted June 4, 2007. During establishment, the plots were irrigated, weeded, and roto-tilled as needed. The plots were evaluated for first year establishment, stem counts, and a visual rating of percent stand. (See table #2.) Any segments of a row missing plants will be filled in with live plants the spring of 2008 to ensure all rows being complete.

Plot Layout MOPMC-T-0716-BF
North


Table #1



Plot #

1 'Cave-In-Rock'	Switchgrass	4	Miscanthus
2 9083274	Big bluestem	5 'Kanlow'	Switchgrass
3 'Alamo'	Switchgrass	6 'Rumsey'	Indiangrass

Individual Plot



Discussion

2008

The plots established very well in 2007. During 2008 the plots were first burned, rototilled between the rows, and 100# actual N (nitrogen) applied as urea and incorporated. Greenhouse grown plugs were planted into any open space of about 4 inches or more within the row, very few plants were necessary. The plots also received chemical weed control.

The plots were harvested one row at a time (1/1000 acre) and evaluated for lodging, and biomass production. (See Table #3) Rows were harvested randomly with row one being harvested at anthesis (at least 50% of seed mature). Row 2 was harvested six weeks after row 1 and that sequence maintained through row 5.

2009

The plots were harvested and evaluated again in 2009 as they were in 2008. All plots received 50# actual N (nitrogen) in May. Biomass production is located in table #4 and element contents are in tables #5-10.

2010

The plots were harvested and evaluated in 2010 as they were in 2008 and 2009. All plots received 50#/acre actual nitrogen (N) in April. Data for all years will be compiled for a final report in 2011.

MOPMC-T-0716					Table #2
Biofuel Study	Establishment Evaluation			11/19/2007	
Cave-in-Rock switchgrass Stm ct	Rep #1	Rep#2	Rep#3	Rep #4	Ave
(Stems per row foot) row #2	67	34	63	72	59
row #3	48	55	61	65	57.25
row #4	47	72	59	63	60.25
row #5	66	93	71	51	70.25
row #6	49	46	76	53	56
Cave-in-Rock switchgrass % Std	Rep #1	Rep#2	Rep#3	Rep #4	Ave
row #2	100	85	92	95	93
row #3	96	100	99	85	95
row #4	95	100	100	95	97.5
row #5	98	100	97	95	97.5
row #6	99	100	95	80	93.5
Alamo switchgrass Stm ct	Rep #1	Rep#2	Rep#3	Rep #4	Ave
(Stems per row foot) row #2	88	99	106	120	103.25
row #3	65	58	105	104	83
row #4	167	118	121	88	123.5
row #5	103	64	106	61	83.5
row #6	118	85	98	114	103.75
Alamo switchgrass % Std	Rep #1	Rep#2	Rep#3	Rep #4	Ave
row #2	100	100	100	100	100
row #3	100	100	100	100	100
row #4	100	100	100	100	100
row #5	100	100	100	95	98.75
row #6	100	100	100	100	100
Kanlow switchgrass Stm ct	Rep #1	Rep #2	Rep #3	Rep #4	
(Stems per row foot) row #2	91	80	77	88	84
row #3	81	39	42	42	51
row #4	95	19	48	78	60
row #5	63	48	111	85	76.75
row #6	67	77	64	61	67.25
Kanlow switchgrass % Std	Rep #1	Rep #2	Rep #3	Rep #4	
row #2	100	95	99	100	98.5
row #3	100	100	97	95	98
row #4	100	100	99	100	99.75
row #5	100	100	100	95	98.75
row #6	98	100	98	95	97.75
Rumsey Indiangrass Stm ct	Rep #1	Rep #2	Rep #3	Rep #4	
(Stems per row foot) row #2	83	129	98	95	101.25
row #3	85	81	84	106	89
row #4	107	72	116	76	92.75
row #5	66	79	85	94	81
row #6	76	82	124	73	88.75

Runsey Indiangrass % Std		Rep #1	Rep #2	Rep #3	Rep #4	
row #2		100	100	100	100	100
row #3		96	100	100	100	99
row #4		100	100	100	95	98.75
row #5		100	100	96	100	99
row #6		100	100	100	100	100
Big Bluestem Stm ct		Rep #1	Rep #2	Rep #3	Rep #4	
(Stems per row foot) row #2		21	22	23	19	21.25
row #3		14	12	18	18	15.5
row #4		18	7	31	24	20
row #5		24	20	27	33	26
row #6		19	13	25	22	19.75
Big Bluestem % Std		Rep #1	Rep #2	Rep #3	Rep #4	
row #2		96	80	95	75	86.5
row #3		92	70	98	65	81.25
row #4		90	60	93	90	83.25
row #5		97	100	94	85	94
row #6		98	30	98	90	79
Miscanthus Stm ct		Rep #1	Rep #2	Rep #3	Rep #4	
(Stems per Plant) row #2		6	40	18	30	23.5
row #3		5	28	12	20	16.25
row #4		18	26	19	19	20.5
row #5		43	25	34	23	31.25
row #6		30	26	25	45	31.5
Miscanthus % Std		Rep #1	Rep #2	Rep #3	Rep #4	
row #2		100	100	30	100	82.5
row #3		100	100	20	100	80
row #4		100	100	19	80	74.75
row #5		80	100	23	40	60.75
row #6		100	100	45	100	86.25

2008/09 Averages for Lodging and Biomass Quantity Table #3

	Lodging at Harvest						Yield (Pounds/Acre)					
	1=No Lodging 5=Extreme Lodging											
	Seed Maturity Cutting	2 nd Cutting	3 rd Cutting	4 th Cutting	5 th Cutting	Ave.	Seed Maturity Cutting	2 nd Cutting	3 rd Cutting	4 th Cutting	5 th Cutting	Ave.
Cave-In-Rock Switchgrass	1.3	1.3	1.4	1.5	1.5	1.4	9,944	10,177	11,628	6,749	6,880	9076
Big Bluestem 9083274 (Epic)	2.0	2.0	2.8	3.6	3.8	2.8	11,118	12,264	10,802	7,788	8,698	10134
Alamo Switch	3.7	2.8	1.9	2.0	1.5	2.4	15,894	15,818	12,944	12,879	12,514	14010
'Freedom' Miscanthus	1.3	1.0	1.1	1.0	1.0	1.1	21,630	24,388	23,619	29,011	24,515	24633
Kanlow Switch	2.3	2.3	3.4	2.1	1.6	2.3	15,835	15,124	15,869	12,182	12,624	14327
Rumsey Indiangrass	4.3	4.3	4.3	4.3	4.4	4.3	13,150	12,811	13,528	10,097	11,192	12156

The data in the chart above shows lodging ratings and amount of biomass harvested starting at seed maturity and continuing every six weeks for four more cuttings. These cuttings started approximately mid Sept. and ran through the beginning of March.

2009/10 Averages for Lodging and Biomass Quantity

Table #3

	Lodging at Harvest						Yield (Pounds/Acre)					
	1=No Lodging 5=Extreme Lodging											
	Seed Maturity Cutting	2 nd Cutting	3 rd Cutting	4 th Cutting	5 th Cutting	Ave.	Seed Maturity Cutting	2 nd Cutting	3 rd Cutting	4 th Cutting	5 th Cutting	Ave.
Cave-In-Rock Switchgrass	3.5	2.6	3.4	3.5	3.1	3.22	9,219	9,329	9,717	9,238	8,543	9,209
Big Bluestem 9083274 (Epic)	1.5	1.8	3.8	4.1	4	3.04	9,951	10,994	10,553	6,997	8,245	9,348
Alamo Switch	1.6	1.9	2.5	2.4	2.6	2.20	12,605	12,810	11,967	11,873	11,560	12,163
'Freedom' Miscanthus	1.1	1	1	1	1.1	1.04	22,756	27,880	18,434	25,780	23,271	23,624
Kanlow Switch	2.5	2.9	4	3.8	3.1	3.26	11,694	11,950	14,123	10,277	12,492	12,107
Rumsey Indiangrass	3.1	3.6	3.5	4.3	3.5	3.60	11,048	12,269	11,157	10,084	9,925	12,107

The data in the chart above shows lodging ratings and amount of biomass harvested starting at seed maturity and continuing every six weeks for four more cuttings. These cuttings started approximately mid Sept. and ran through the beginning of March.

LSD All - Pairwise Comparison Tests for Harvest Date

Table #5

Test of: adf

Common Name	Harvest Date	Mean	Homogeneous Groups
Alamo	30-Mar	55.639	A
	5-Jan	54.524	A
	17-Feb	53.855	AB
	26-Nov	52.301	AB
	14-Oct	50.681	B
Big bluestem	24-Mar	57.072	A
	9-Feb	56.761	A
	30-Dec	55.467	A
	18-Nov	54.292	A
	6-Oct	49.376	B
Cave-in-Rock switchgrass	24-Feb	59.652	A
	12-Jan	58.556	A
	2-Dec	55.607	B
	21-Oct	48.528	C
	10-Sep	44.29	D
Indiangrass	9-Feb	56.689	A
	30-Dec	56.229	A
	24-Mar	55.669	A
	18-Nov	54.120	A
	6-Oct	49.779	B
Kanlow - switchgrass	26-Jan	55.301	A
	5-Nov	53.16	B
	17-Dec	52.87	B
	1-Oct	48.39	C
Miscanthus	30-Mar	64.703	A
	5-Jan	63.888	A
	17-Feb	63.678	AB
	26-Nov	60.49	B
	14-Oct	53.456	C

Study: MOPMC-P-0717-PA, WL**Study Title:** Evaluation and Release of a Shade Tolerant Little Bluestem for Silvopasture**Study Leaders:** Van Sambeek, J., D. Wallace, G. Garrett, S. Bruckerhoff**Introduction:**

Little bluestem is one of the most widely distributed native grasses in North America. It will grow on a wide variety of soils but is very well adapted to well-drained, medium to dry, infertile soils. The plant has excellent drought and fair shade tolerance, and fair to poor flood tolerance. Little bluestem is considered less valuable for grazing due to its tussock growth form and perceived lower forage quality. In Missouri, it is often found growing as an understory plant in open woodlands, savannas, and transitional forested areas.

The Elsberry PMC initiated a little bluestem collection from the Ozark region (Missouri, Arkansas, and Oklahoma) in 1996 to develop an improved little bluestem that would be better adapted to this region. The collection effort resulted in an assembly of 130 little bluestem accessions.

Problem:

There is limited information available on species selected for forage to be primarily used for savannas and silvopasture systems throughout Missouri.

Objective:

An existing PMC little bluestem collection displays an amazing range of morphological and phenological characteristics – color, plant height, blade width, stem erectness, rust resistance, spring emergence, anthesis, etc. This collection has never been evaluated for shade attributes that might be valuable in silvopasture systems or with savanna restoration efforts. This study will select and evaluate for forage production in relation to shade tolerance.

Cooperators:

The USDA Natural Resources Conservation Service, Plant Materials Center and the University of Missouri, Columbia, Horticulture and Agroforestry Research Center, HARC.

Procedure:

A randomized complete block design with three replications, five shade levels: full sun, 30% shade, 55% shade, 55% shade with sunflecks, and 80% shade will be used for the study.

Twenty-eight accessions of little bluestem are selected based on original collection site descriptions that indicated collections were associated with woodland conditions. These collections came from previous work (Study 29I141G) at the Elsberry PMC and will be assembled in 2008 at the University of Missouri Agroforestry Center (HARC). Additional varieties being evaluated are the little bluestem Northern region assembly and the bluestem Southern region assembly. Plants will be started in the greenhouse and planted in replicated plots the spring 2008.

Forty-five pots for each accession will be randomized within each treatment. Above ground dry weight, leaf weight, and forage quality (acid detergent fiber, neutral detergent fiber, crude protein) will be collected for each accession.

Discussion:

2007

Collection reports from PMC study 29I141G were reviewed and 28 accessions were selected based on woodland site descriptions (Table #1). Two selections under evaluation at the PMC and two established cultivars will also be included.

Study MOPMC-P-0717-PA, WL				Table #1
PMC Accessions Selected For Shade Tolerance Evaluation				
		Location Description		
Accn No.	State Code	Soil Type	County	Site
9078847	IA-1	Gara	Clarke	Woods Edge (WE)
9078848	IA-2	Gara	Clarke	WE
9078850	IA-4	Monona	Woodbury	WE
9078852	IA-6	Hamburg	W. Pottawattamie	Woods (W)
9078858	IA-12	Haydon	Dallas	Open Woods (OW)
9078961	IA-27	Cresco	Scott	OW
9078873	IL-1	Rodman	Kane	W
9078875	IL-3	Miami Casco	Kane	OW
9078876	IL-4	Rozetta	Pike	WE
9078880	IL-8	Ashbury	Will	Ditch
9078882	IL-10	Hamburg	Henderson	W
9078887	IL-15	Bloomfield	Mason	OW
9078888	IL-16	Ava	Marion	W/G Lode
9078891	IL-19	SOGN	Calhoun	Glade
9078894	MO-1	Winnegan	Chariton	W
9078895	MO-3		Cape	
9078898	MO-6	Keswick	Putnum	OW
9078899	MO-7	Midco	Ripley	OW
9078902	MO-10	Captina	Carter	WE
9078905	MO-13	Clarksville	Butler	W (Row)
9078915	MO-23		Perry	
9078917	MO-25	Gorin	Monroe	OW
9078921	MO-29	Goss	Montgomery	
9078956	MO-64	-	Shannon	Glade
9078959	MO-67	-	Wayne	OW
9078960	MO-68	Keswick	Pike	WE
9083271			Composite Selection (South Region)	
9083317			Composite Selection (North Region)	
421553	'Aldous'	Kansas Flinthills		
421552	'Cimarron'	Eastern Kansas and Oklahoma		

2010

Doug Wallace, study leader, took a different position within NRCS, therefore no data has been reported for this study in the last two years. Data has been collected and the Elsberry PMC personnel is working with Jerry Van Sambeek, Forest Service, to compile the data and it will be reported in the 2011 ATR.

Study: MOPMC-T-0718-WE, WL

Study Title: Evaluation of the Flood Tolerance of Planted Oak Seedlings Derived from Different Seed Origins

Study Leaders: Cogshall, M., University of Missouri
Cordsiemon, R., Elsberry Plant Materials Center

Introduction:

Many plant species exhibit wide amplitude in terms of their capacity to occupy a range of positions along a hydrologic gradient. The presence or absence of a particular tree species in a bottomland hardwood forest is primarily dependent upon soil moisture gradients, stream deposition patterns, and flooding season and duration (Hodges, 1997). In the Midwestern U.S., there is considerable interest in restoring native ecosystems on former bottomland sites that are now being devoted to marginal agriculture (Kruse and Groninger, 2003). Oak seedlings are a major component of these restoration efforts

Problem:

Little is known about whole tree responses to a variety of flooding treatments for different oak seedlings used in WRP and other wetland programs.

Objective:

The specific objectives of the proposed research are: 1) to determine the effect of controlled flooding treatments, including inundation, on the survival and regrowth of planted oak seedlings in the Flood Laboratory located at the NRCS Plant Materials Center (PMC) in Elsberry, Missouri; 2) to contrast these flooding results with additional flood evaluations using planted seedlings in the Flood Tolerance Laboratory (FTL) at the Horticulture and Agroforestry Research Center (HARC) in New Franklin, Missouri; and 3) initiate a potted seedling study as a means to further expand our knowledge base of how plants respond to controlled flooding events.

Procedure:

A total of 1,746 test trees were established as 1-0 bare root seedlings in the PMC Flood Laboratory (Field #4) in May 2007. Spacing was 4' x 10'. The entire test planting covers 1.60 acres. A total of seven different oak species were planted including: bur oak (12 seedlots), chinkapin oak (2 seedlots), pin oak (6 seedlots), northern red oak (9 seedlots), shumard oak (8 seedlots), swamp white oak (27 seedlots) and white oak (25 seedlots). Each seedlot contains 18 seedlings which were established in a series of 18 single tree plots. All seedlots planted in the PMC planting were also common to the FTL planting at HARC. The experimental design for this planting is a randomized complete block design with 18 replications. Replication boundaries were defined based on slope position within the Elsberry Flood Laboratory to the extent possible.

Discussion:

2007

Collection of first year seedling survival, height and diameter data in the PMC planting will be made in the winter of 2007/2008. Flooding treatments will coincide with the natural flooding for calendar year 2007 of the Mississippi River, based upon flood stage data from the nearest river gauging station to the Elsberry planting site. Survival data will be collected immediately following the end of the first flooding period and re-growth potential. Flooding will be assessed as a function of shoot growth in the year following. This data will be contrasted with results obtained from the flood treatments that are planned in the HARC FTL, which will be initiated following the completion of the first growth flush in the spring 2008.

2008

In 2008, the PMC staff flooded the evaluation plot from 2 -24 inches. The water was left on the trees for 16 days, beginning the first part of April to mimic the flood stage of the Mississippi River in 2007 at the Winfield Lock and Dam. Mark Coggeshall and students collected data in late summer.

2009

The wetland cell was saturated for most of the growing season and impossible to maintain. Mr. Coggeshall and his team did not make it to the PMC to do any further evaluations. In 2010, a decision will be made to determine whether to continue with the study or not.

2010

Mr. Coggeshall and his team from the University of Missouri cut the remaining material within the wetland cell and collected data. The PMC is working with Jerry Van Sambeek, Forest Service at the U. of Mo. HARC farm, to organize data collected from the evaluations that took place on the selected oak species. A final report will be organized for this study and documented in the 2011 ATR.

Study: MOPMC-P-0820-UR

Study Title: Evaluation and Release of Native Plants for Urban Landscaping.

Study Leader: Bruckerhoff, S. B.

Introduction:

Plants for urban landscaping have been an ever increasing market for many years. Recently the use of herbaceous native species is also increasing. This type of vegetation can be used for its aesthetics and also in areas to help control erosion, runoff and improve water quality.

Problem:

There are few current varieties of little bluestem, big bluestem or broomsedge bluestem on the market that have been specifically selected for landscape purposes. There is demand in the nursery industry for this type of plant material and current collections already in place on the PMC can provide desirable germplasm.

Objective:

The objective is to assemble, evaluate, develop and release adapted selections of little bluestem, big bluestem and broomsedge bluestem for landscape use in Missouri, Iowa, and Illinois.

Procedure:

Native plants with desirable landscape and/or ornamental characteristics will be assembled and evaluated. Plants will be selected from existing assemblies at the PMC and also collected from native prairie remnants within the service area. Plants selected for evaluation will be collected vegetatively or by seed depending on the species and trait being selected for.

Existing assemblies of big and little bluestem will be evaluated for color, upright growth form, and stability during the dormant season. During the growing season colors may be deep and rich, or abnormally pale. Autumn colors will have reddish tints to leaves and elaborate seedheads.

Selected plants will be vegetatively collected, increased in the greenhouse, and transplanted into increase plots. Plants maintaining desired characteristics when mature will be divided (cloned) and made available to commercial growers as improved native landscape plants.

Seed will be collected of broomsedge bluestem (and other native species that naturally exhibit attractive characteristics for urban landscaping) and placed in the germinator. Selections will be made on quick germination. Selected plants will be placed in an evaluation nursery where undesirable plants will be culled. Remaining plants will remain for seed production to establish an increase field. This selection will be for color and quick establishment. Seed from the increase field will be made available to commercial growers.

Discussion:

2008/2009

Plants of big and little bluestem in evaluation plantings at the PMC were visually evaluated for color and form. After going dormant, the plants were dug in the fall of 2008, divided into multiple plants and grown out in the greenhouse. These plants (approximately 700 little bluestem and 300 big bluestem) were put on three foot spacings in a block for each species. After a year establishment period, the better plants will be selected for increase.

Seed of broomsedge bluestem was collected from 17 locations throughout Missouri and placed in the seed germinator. Seed germinating within a seven day period was transplanted into plug flats and transplanted into a crossing block including plants from all locations.

2010

The little bluestem and big bluestem selections were placed in an evaluation nursery, where they were allowed to grow. There are concerns on how to handle this material, since it does not follow the program protocol for a typical release. There is very little data collected in order to make a release. A determination will be made as to whether or not this study will continue in 2011.

The selected broomsedge plants were planted in the fall of 2009, but did not survive the winter. Inferior root development could have been a possibility for poor survival. It has been determined that broomsedge will be discontinued.

Study: MOPMC-P-0821-BF

Study Title: Evaluation and Release of an Iowa Source, stiff stemmed Indiangrass, *Sorghastrum nutans*, for Biofuel.

Study Leader: Cordsiemon, R., Elsberry Plant Materials Center

Introduction:

The Northern Germplasm, Central Germplasm, and Southern Germplasm production fields of Iowa Ecotype Indiangrass will be evaluated for biofuel potential. This study will develop a new selection of Indiangrass for Iowa that has high biomass production, lodging resistance, and is quick to establish.

Objective:

Switchgrass (*Panicum virgatum* L.) and giant cane (*Miscanthus giganteus*) are commonly used in the evaluation of herbaceous species for biofuels. Another warm season species that is showing potential is Indiangrass (*Sorghastrum nutans*). The objective is to develop an Indiangrass that has the potential for use in the biofuel industry.

Discussion:

2008

In March, 2008, plants were selected out of each zone from the Iowa Ecotype Study. The ecotype zones break the state of Iowa up into 3 separate zones. There is a northern, central, and southern zone. The PMC has production plots of each zone of Indiangrass and from these plots, 3 to 4 plants were selected. They were selected based on their visual robust, tall standing stature and ability not to lodge. These plants were separated into plugs and propagated in the greenhouse. There were 3 plants from the northern zone, 4 plants from the central zone, and 3 plants from the central zone. 40 plugs were started from each plant that was selected. An evaluation planting was established at Forrest Keeling Nursery with 10 plants per replication and 4 replications. The plants were allowed to cross pollinate and an equal amount of seed was taken off each plant. The seed will be used to start seedlings in the germinator and selections will be made based on how quick the seed germinates.

2009

An evaluation block was established using the seedlings that germinated within 7 days of entering the growth chamber. The seedlings were transplanted into conetainers and allowed to establish in the greenhouse and were then planted in the southeast corner of field #7F. There are 224 plants in the evaluation block. The plants will be evaluated individually for basal spread, height at seedhead, height of foliage, 5 random stem counts for diameter, and late winter evaluation for lodging and overall plant rating. Data will be

collected at the beginning of spring to determine ratings for plant lodging and the overall plant rating. The plants will be analyzed and selections will be made from the data collected. The 2010 annual technical report will show collection data and selections made for the new crossing block.

2010

In the spring of 2010, the final evaluations were made for lodging and overall plant rating. The data that was collected in 2009 and 2010 was used in an arbitrary formula to weigh out the different plants in the 224 plant plot. Approximately the top 10 percent will be selected for superior growth characteristics. The emphasis of this study was to select an Iowa source Indiangrass with biofuel potential. Due to Agriculture Research Service and the Forest Service having primary responsibility for biomass/biofuel selections, the PMC will evaluate the potential for this selection as a possible benefit to wildlife and/or living snow fence or wind barrier. Selected plants have been isolated in a crossing block and seed should be available next year for a foundation block and field testing. Data collected is found in table #1.

Iowa Biofuel Indiangrass (Selected from FKN plot)
 Planted in Spring 2009 - Field 7F



101	102	103	104	105	106	107	108	109	110	111	112	113	114
201	202	203	204	205	206	207	208	209	210	211	212	213	214
301	302	303	304	305	306	307	308	309	310	311	312	313	314
401	402	403	404	405	406	407	408	409	410	411	412	413	414
501	502	503	504	505	506	507	508	509	510	511	512	513	514
601	602	603	604	605	606	607	608	609	610	611	612	613	614
701	702	703	704	705	706	707	708	709	710	711	712	713	714
801	802	803	804	805	806	807	808	809	810	811	812	813	814
901	902	903	904	905	906	907	908	909	910	911	912	913	914
1001	1002	1003	1004	1105	1006	1007	1008	1009	1010	1011	1012	1013	1014
1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114
1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214
1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314
1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414
1501	1502	1503	1504	1505	1506	1507	1508	1509	1510	1511	1512	1513	1514
1601	1602	1603	1604	1605	1606	1607	1608	1609	1610	1611	1612	1613	1614

Iowa Indiangrass Biofuel Study
11-2-2009 Evaluation
3-8-2010 Evaluation for Lodging and Overall Rating ONLY

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average							Lodging	Overall Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average				
					~to the nearest thousandth of an inch~									
1009	9	8	85	38	0.249	0.211	0.205	0.229	0.224	0.2236	1	1	306	
809	7	9	73	36	0.178	0.192	0.135	0.157	0.168	0.166	1	1	188	
405	6	7	80	25	0.225	0.218	0.154	0.223	0.21	0.206	1	1	108	
610	7	7	74	27	0.128	0.202	0.1	0.16	0.128	0.1436	1	1	95	
1101	7	7	72	22	0.15	0.161	0.191	0.162	0.181	0.169	1	1	91	
702	8	8	66	29	0.194	0.222	0.196	0.188	0.232	0.2064	1	3	64	
202	9	9	67	28	0.199	0.181	0.116	0.151	0.181	0.1656	1	3	63	
1007	7	8	72	26	0.225	0.207	0.231	0.232	0.25	0.229	1	3	56	
302	8	9	63	23	0.166	0.222	0.215	0.176	0.192	0.1942	1	3	54	
1309	8	7	74	26	0.188	0.189	0.132	0.202	0.184	0.179	3	1	43	
1509	8	7	77	26	0.167	0.227	0.099	0.172	0.209	0.1748	3	1	42	
109	7	5	75	38	0.175	0.231	0.156	0.177	0.205	0.1888	3	1	42	
909	7	8	72	27	0.164	0.15	0.181	0.168	0.167	0.166	3	1	42	
806	7	7	68	28	0.114	0.178	0.17	0.214	0.202	0.1756	3	1	40	
1107	7	7	69	26	0.189	0.16	0.202	0.193	0.2	0.1888	1	3	40	
614	7	6	71	26	0.228	0.199	0.18	0.208	0.199	0.2028	1	3	37	
1411	7	7	69	24	0.181	0.131	0.223	0.173	0.198	0.1812	1	3	36	
310	7	7	80	25	0.204	0.131	0.159	0.158	0.2	0.1704	3	1	35	
410	6	7	74	26	0.198	0.178	0.203	0.159	0.186	0.1848	3	1	34	
1105	7	7	64	22	0.199	0.163	0.182	0.201	0.131	0.1752	3	1	31	
1514	6	7	60	25	0.198	0.161	0.181	0.219	0.116	0.175	1	3	31	
1111	6	6	79	28	0.232	0.113	0.142	0.204	0.196	0.1774	1	3	30	

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average								Lodging 1 - 3 - 5 Rating	Overall Rating 1 - 3 - 5 Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average					
	~to the nearest inch~		~to the nearest inch~		~to the nearest thousandth of an inch~										
209	7	6	73	26	0.118	0.189	0.208	0.143	0.141	0.1598	1	3	29		
902	7	7	63	21	0.176	0.138	0.162	0.218	0.132	0.1652	1	3	28		
1102	6	7	67	22	0.182	0.167	0.196	0.173	0.173	0.1782	1	3	27		
808	6	7	66	21	0.161	0.195	0.171	0.203	0.2	0.186	3	1	27		
1014	7	7	72	22	0.208	0.116	0.124	0.135	0.153	0.1472	1	3	26		
1609	7	7	64	22	0.102	0.164	0.149	0.153	0.139	0.1414	3	1	25		
912	5	6	75	31	0.132	0.082	0.203	0.199	0.131	0.1494	1	3	23		
409	6	6	72	23	0.181	0.175	0.157	0.154	0.161	0.1656	1	3	23		
1304	8	7	65	17	0.174	0.092	0.166	0.12	0.148	0.14	1	3	22		
1314	5	5	64	25	0.197	0.149	0.097	0.153	0.202	0.1596	1	3	17		
501	6	6	59	18	0.163	0.166	0.164	0.157	0.104	0.1508	1	3	16		
203	7	5	62	19	0.126	0.146	0.166	0.138	0.156	0.1464	1	3	16		
505	8	8	68	25	0.202	0.232	0.165	0.168	0.118	0.177	3	3	16		
1607	7	7	72	27	0.254	0.145	0.213	0.166	0.227	0.201	3	3	15		
714	7	7	71	24	0.225	0.156	0.212	0.232	0.225	0.21	3	3	14		
1613	7	7	62	19	0.187	0.148	0.132	0.112	0.157	0.1472	1	5	14		
914	7	8	59	24	0.207	0.177	0.163	0.208	0.147	0.1804	3	3	13		
901	6	7	72	28	0.19	0.2	0.23	0.182	0.206	0.2016	3	3	13		
1311	7	6	71	32	0.206	0.187	0.135	0.146	0.195	0.1738	3	3	13		
807	7	6	69	25	0.198	0.227	0.214	0.218	0.243	0.22	3	3	13		
512	6	5	67	25	0.204	0.153	0.216	0.15	0.126	0.1698	1	5	13		
1114	6	6	59	21	0.189	0.144	0.168	0.145	0.19	0.1672	1	5	13		
1401	7	7	69	25	0.126	0.185	0.204	0.172	0.23	0.1834	3	3	12		
1512	7	7	77	24	0.227	0.172	0.16	0.197	0.196	0.1904	3	3	12		
709	7	8	68	24	0.195	0.193	0.119	0.156	0.169	0.1664	3	3	12		
814	6	6	60	18	0.236	0.132	0.215	0.153	0.216	0.1904	1	5	12		
1209	7	7	73	22	0.201	0.172	0.208	0.205	0.211	0.1994	3	3	12		
805	7	6	75	26	0.209	0.189	0.169	0.181	0.194	0.1884	3	3	11		
908	7	7	66	21	0.182	0.153	0.23	0.215	0.219	0.1998	3	3	11		

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average							Lodging	Overall Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average				
					~to the nearest thousandth of an inch~									
1508	7	6	70	24	0.194	0.227	0.161	0.221	0.213	0.2032	3	3	11	
1513	7	7	61	22	0.218	0.178	0.19	0.167	0.196	0.1898	3	3	11	
1409	8	7	65	21	0.136	0.116	0.164	0.19	0.248	0.1708	3	3	11	
1412	6	6	64	24	0.255	0.188	0.247	0.245	0.222	0.2314	3	3	11	
1611	7	7	61	24	0.112	0.191	0.2	0.169	0.162	0.1668	3	3	11	
802	9	8	61	22	0.157	0.19	0.21	0.225	0.249	0.2062	3	5	11	
813	6	6	64	20	0.171	0.149	0.076	0.18	0.18	0.1512	1	5	11	
601	5	5	55	18	0.16	0.177	0.12	0.106	0.146	0.1418	1	3	11	
1511	7	7	74	24	0.142	0.156	0.169	0.184	0.161	0.1624	3	3	11	
604	7	7	54	25	0.124	0.178	0.139	0.165	0.168	0.1548	3	3	11	
102	6	6	67	28	0.18	0.23	0.171	0.202	0.156	0.1878	3	3	11	
211	7	8	66	24	0.158	0.16	0.207	0.096	0.081	0.1404	3	3	10	
1105	6	7	72	24	0.166	0.177	0.133	0.226	0.221	0.1846	3	3	10	
707	6	6	59	25	0.191	0.214	0.196	0.208	0.219	0.2056	3	3	10	
210	7	6	67	25	0.18	0.168	0.145	0.198	0.183	0.1748	3	3	10	
1407	6	6	78	26	0.168	0.205	0.173	0.224	0.19	0.192	3	3	10	
1605	6	7	68	19	0.248	0.208	0.144	0.272	0.224	0.2192	3	3	10	
306	6	8	79	22	0.188	0.195	0.149	0.158	0.134	0.1648	3	3	10	
1207	6	7	64	20	0.187	0.206	0.222	0.168	0.247	0.206	3	3	10	
108	7	7	70	21	0.235	0.149	0.092	0.153	0.201	0.166	3	3	9	
110	6	7	65	23	0.176	0.195	0.186	0.152	0.167	0.1752	3	3	9	
1507	6	7	72	20	0.241	0.191	0.207	0.126	0.239	0.2008	3	3	9	
609	6	6	70	25	0.173	0.177	0.178	0.226	0.182	0.1872	3	3	9	
1504	8	8	63	19	0.129	0.14	0.159	0.14	0.106	0.1348	3	3	9	
811	7	6	62	24	0.106	0.19	0.179	0.172	0.148	0.159	3	3	9	
1214	7	6	66	22	0.175	0.183	0.123	0.194	0.184	0.1718	3	3	9	
1313	7	6	66	22	0.225	0.129	0.204	0.097	0.2	0.171	3	3	9	
1213	6	6	67	23	0.184	0.186	0.198	0.167	0.209	0.1888	3	3	9	
907	5	5	66	19	0.197	0.122	0.213	0.205	0.168	0.181	1	5	9	

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average							Lodging 1 - 3 - 5 Rating	Overall Rating 1 - 3 - 5 Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average				
	~to the nearest inch~		~to the nearest thousandth of an inch~											
911	6	6	71	24	0.233	0.212	0.177	0.117	0.129	0.1736	3	3	8	
106	5	6	60	18	0.098	0.172	0.192	0.174	0.133	0.1538	1	5	8	
1106	6	6	75	23	0.185	0.154	0.19	0.2	0.171	0.18	3	3	8	
1003	7	7	60	18	0.2	0.173	0.111	0.142	0.203	0.1658	3	3	8	
1510	6	7	70	24	0.154	0.123	0.123	0.149	0.175	0.1448	3	3	8	
1109	6	6	63	22	0.207	0.163	0.189	0.169	0.174	0.1804	3	3	8	
407	6	6	65	20	0.151	0.232	0.243	0.167	0.17	0.1926	3	3	8	
703	7	7	50	21	0.144	0.171	0.128	0.126	0.102	0.1342	3	3	8	
504	6	6	72	21	0.188	0.21	0.184	0.167	0.146	0.179	3	3	8	
1310	7	7	72	21	0.211	0.226	0.21	0.202	0.231	0.216	3	5	7	
301	5	6	66	24	0.237	0.173	0.119	0.196	0.193	0.1836	3	3	7	
510	6	6	70	22	0.212	0.204	0.167	0.119	0.111	0.1626	3	3	7	
213	6	7	60	24	24	0.178	0.16	0.185	0.189	0.127	3	3	7	
112	6	6	66	24	0.157	0.153	0.191	0.15	0.084	0.147	3	3	7	
1201	7	6	66	19	0.176	0.174	0.202	0.19	0.047	0.1578	3	3	7	
303	6	7	60	23	0.091	0.193	0.085	0.138	0.139	0.1292	3	3	7	
708	6	6	63	21	0.152	0.182	0.2	0.133	0.148	0.163	3	3	7	
1307	6	7	68	21	0.234	0.241	0.225	0.252	0.141	0.2186	3	5	6	
207	5	6	61	24	0.156	0.167	0.186	0.157	0.096	0.1524	3	3	6	
1408	6	5	67	18	0.182	0.204	0.151	0.242	0.227	0.2012	3	3	6	
1108	6	6	70	24	0.18	0.242	0.182	0.242	0.201	0.2094	3	5	6	
1414	5	4	68	23	0.143	0.154	0.129	0.118	0.098	0.1284	1	5	6	
905	5	7	68	18	0.167	0.225	0.141	0.161	0.149	0.1686	3	3	6	
1608	6	7	71	20	0.206	0.237	0.175	0.164	0.258	0.208	3	5	6	
1306	7	6	70	20	0.19	0.234	0.183	0.219	0.202	0.2056	3	5	6	
1208	5	6	73	22	0.128	0.15	0.184	0.131	0.175	0.1536	3	3	6	
1203	6	7	49	17	0.157	0.133	0.118	0.106	0.19	0.1408	3	3	6	
413	6	6	59	18	0.185	0.135	0.129	0.166	0.157	0.1544	3	3	6	
1211	6	6	63	26	0.165	0.164	0.171	0.164	0.224	0.1776	3	5	6	

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average								Lodging 1 - 3 - 5 Rating	Overall Rating 1 - 3 - 5 Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average					
	~to the nearest inch~		~to the nearest inch~		~to the nearest thousandth of an inch~										
1606	5	5	67	23	0.133	0.167	0.191	0.189	0.15	0.166	3	3	5		
1010	6	7	66	23	0.21	0.129	0.123	0.184	0.174	0.164	3	3	5		
705	5	5	68	23	0.194	0.154	0.157	0.141	0.178	0.1648	3	3	5		
304	6	6	63	19	0.157	0.16	0.145	0.127	0.096	0.137	3	3	5		
1301	6	7	56	23	0.169	0.158	0.176	0.141	0.163	0.1614	3	3	5		
305	6	5	73	20	0.108	0.18	0.169	0.187	0.127	0.1542	3	3	5		
1103	7	7	46	16	0.147	0.086	0.091	0.106	0.157	0.1174	3	3	5		
1505	6	7	68	17	0.207	0.208	0.241	0.217	0.179	0.2104	3	3	5		
1008	5	5	73	24	0.18	0.166	0.125	0.099	0.162	0.1464	3	3	5		
1604	7	7	58	18	0.22	0.133	0.126	0.196	0.128	0.1606	3	3	5		
801	6	6	63	18	0.245	0.175	0.205	0.197	0.265	0.2174	3	3	5		
1413	6	6	61	21	0.201	0.181	0.137	0.219	0.182	0.184	3	3	5		
208	5	6	60	25	0.201	0.194	0.193	0.163	0.163	0.1828	3	3	5		
1308	6	6	64	23	0.128	0.15	0.201	0.172	0.176	0.1654	3	3	5		
1506	6	6	72	23	0.198	0.147	0.226	0.137	0.105	0.1626	3	3	5		
204	6	5	63	20	0.116	0.136	0.166	0.115	0.134	0.1334	3	3	4		
1013	7	6	56	17	0.16	0.193	0.182	0.205	0.194	0.1868	3	3	4		
1302	6	7	63	19	0.168	0.161	0.193	0.172	0.135	0.1658	3	3	4		
1205	7	6	64	18	0.172	0.199	0.154	0.153	0.183	0.1722	3	3	4		
313	6	6	59	19	0.187	0.183	0.227	0.139	0.213	0.1898	3	3	4		
1303	6	7	63	18	0.18	0.163	0.176	0.151	0.184	0.1708	3	3	4		
906	6	6	66	19	0.239	0.207	0.215	0.104	0.176	0.1882	3	3	4		
904	7	7	62	19	0.094	0.117	0.137	0.199	0.144	0.1382	3	3	4		
1602	6	8	58	19	0.082	0.167	0.149	0.155	0.127	0.136	3	3	4		
1012	6	6	65	23	0.144	0.148	0.162	0.138	0.147	0.1478	3	3	4		
1305	6	6	67	18	0.205	0.19	0.229	0.103	0.208	0.187	3	3	4		
1410	6	6	72	22	0.097	0.169	0.174	0.108	0.195	0.1486	3	3	4		
810	5	6	65	20	0.182	0.193	0.189	0.249	0.159	0.1944	3	3	4		
1212	7	6	77	15	0.234	0.128	0.171	0.174	0.202	0.1818	3	3	4		

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average							Lodging	Overall Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average				
	~to the nearest inch~		~to the nearest inch~		~to the nearest thousandth of an inch~									
1004	6	7	61	15	0.157	0.182	0.204	0.221	0.122	0.1772	3	5	4	
1404	7	6	68	18	0.129	0.081	0.209	0.192	0.119	0.146	3	5	4	
502	7	6	66	15	0.181	0.18	0.171	0.177	0.165	0.1748	5	3	4	
910	6	5	68	19	0.17	0.193	0.243	0.228	0.129	0.1926	3	5	4	
711	6	6	58	18	0.169	0.185	0.175	0.175	0.116	0.164	3	5	4	
412	5	5	74	23	0.199	0.191	0.168	0.177	0.184	0.1838	3	5	4	
803	6	7	59	20	0.143	0.124	0.136	0.12	0.105	0.1256	3	5	4	
1406	5	6	55	17	0.211	0.145	0.206	0.247	0.222	0.2062	3	5	4	
1011	6	5	69	19	0.204	0.175	0.171	0.171	0.19	0.1822	3	5	3	
1610	6	6	59	20	0.154	0.164	0.11	0.116	0.173	0.1434	3	5	3	
602	6	5	54	22	0.165	0.118	0.176	0.16	0.155	0.1548	3	5	3	
1210	5	5	63	23	0.164	0.217	0.2	0.142	0.162	0.177	3	5	3	
701	6	6	54	19	0.156	0.145	0.138	0.135	0.166	0.148	3	5	3	
205	5	5	79	22	0.167	0.19	0.169	0.181	0.198	0.181	5	3	3	
913	6	6	63	21	0.152	0.128	0.12	0.106	0.146	0.1304	3	5	3	
1202	5	7	56	12	0.209	0.176	0.234	0.228	0.242	0.2178	3	5	3	
613	6	5	66	19	0.176	0.156	0.154	0.195	0.116	0.1594	3	5	3	
1001	5	5	65	17	0.21	0.136	0.232	0.242	0.236	0.2112	3	5	3	
1402	6	7	63	14	0.14	0.14	0.148	0.177	0.145	0.15	3	5	3	
403	6	6	57	18	0.148	0.133	0.144	0.124	0.127	0.1352	3	5	3	
1612	6	6	72	24	0.135	0.172	0.185	0.173	0.159	0.1648	5	5	3	
1405	6	6	66	19	0.164	0.207	0.226	0.219	0.211	0.2054	5	5	3	
804	6	6	57	18	0.126	0.179	0.137	0.116	0.092	0.13	3	5	3	
611	5	5	60	18	0.166	0.242	0.149	0.162	0.202	0.1842	3	5	3	
1502	5	6	49	17	0.134	0.178	0.169	0.159	0.141	0.1562	3	5	3	
607	5	6	58	16	0.207	0.177	0.182	0.137	0.121	0.1648	3	5	3	
503	5	5	73	22	0.165	0.183	0.089	0.128	0.152	0.1434	3	5	3	
608	5	5	60	16	0.204	0.158	0.169	0.251	0.179	0.1922	3	5	3	
506	6	5	63	16	0.189	0.107	0.183	0.158	0.158	0.159	3	5	3	

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average							Lodging	Overall Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average				
	~to the nearest inch~		~to the nearest inch~		~to the nearest thousandth of an inch~									
513	5	5	64	19	0.173	0.185	0.173	0.147	0.125	0.1606	3	5	3	
606	5	5	66	18	0.159	0.134	0.148	0.201	0.202	0.1688	3	5	3	
1503	7	7	56	20	0.158	0.108	0.089	0.134	0.143	0.1264	5	5	2	
309	5	5	63	18	0.179	0.195	0.152	0.145	0.106	0.1554	3	5	2	
111	6	6	56	18	0.187	0.181	0.198	0.196	0.135	0.1794	5	5	2	
509	5	5	57	17	0.181	0.131	0.207	0.165	0.126	0.162	3	5	2	
404	5	6	56	16	0.169	0.146	0.107	0.167	0.118	0.1414	3	5	2	
1603	7	6	64	20	0.151	0.1	0.134	0.07	0.212	0.1334	5	5	2	
201	5	4	63	21	0.165	0.125	0.187	0.121	0.154	0.1504	3	5	2	
107	5	6	52	21	0.183	0.138	0.214	0.15	0.148	0.1666	5	5	2	
704	6	4	52	17	0.15	0.119	0.128	0.176	0.174	0.1494	3	5	2	
1002	5	6	59	17	0.242	0.15	0.225	0.19	0.165	0.1944	5	5	2	
308	5	4	60	21	0.182	0.124	0.161	0.13	0.098	0.139	3	5	2	
903	5	5	59	21	0.171	0.154	0.18	0.253	0.15	0.1816	5	5	2	
1501	6	6	54	16	0.156	0.17	0.212	0.181	0.103	0.1644	5	5	2	
212	5	5	56	22	0.162	0.24	0.168	0.131	0.124	0.165	5	5	2	
612	4	7	67	14	0.136	0.103	0.15	0.093	0.168	0.13	3	5	2	
1006	5	6	62	17	0.173	0.181	0.11	0.15	0.165	0.1558	5	5	2	
1113	5	5	59	20	0.145	0.166	0.148	0.12	0.192	0.1542	5	5	2	
406	5	5	56	18	0.165	0.189	0.18	0.163	0.123	0.164	5	5	1	
713	6	6	30	16	0.158	0.08	0.129	0.142	0.109	0.1236	5	5	1	
103	5	6	40	11	0.1	0.125	0.138	0.123	0.15	0.1272	3	5	1	
1312	5	6	61	13	0.202	0.193	0.205	0.21	0.067	0.1754	5	5	1	
1403	5	5	66	16	0.116	0.11	0.139	0.233	0.212	0.162	5	5	1	
311	4	5	57	20	0.192	0.138	0.183	0.153	0.097	0.1526	5	5	1	
603	4	4	63	26	0.206	0.144	0.13	0.145	0.106	0.1462	5	5	1	
812	5	5	39	14	0.174	0.173	0.191	0.126	0.17	0.1668	5	5	1	
1601	5	5	43	17	0.12	0.145	0.112	0.131	0.137	0.129	5	5	1	
712	5	5	33	14	0.141	0.134	0.144	0.104	0.203	0.1452	5	5	1	

1=Best and 5=Poor

Plant ID #	Basal Spread		Height		Stem Diameter Average							Lodging	Overall Rating	Selection Formula
	N-S	E-W	Seedhead	Foliage	Stem 1	Stem 2	Stem 3	Stem 4	Stem 5	Average				
	~to the nearest inch~		~to the nearest inch~		~to the nearest thousandth of an inch~							1 - 3 - 5 Rating		
206	5	4	68	14	0.23	0.183	0.182	0.149	0.155	0.1798	5	5	1	
710	5	5	47	12	0.175	0.158	0.194	0.149	0.127	0.1606	5	5	1	
508	4	5	56	15	0.14	0.131	0.15	0.215	0.162	0.1596	5	5	1	
1206	5	5	42	15	0.109	0.127	0.14	0.104	0.103	0.1166	5	5	1	
1204	4	4	47	14	0.152	0.179	0.162	0.138	0.192	0.1646	5	5	1	
1112	5	5	46	14	0.097	0.127	0.094	0.098	0.087	0.1006	5	5	1	
1104	5	4	46	15	0.118	0.119	0.105	0.11	0.112	0.1128	5	5	1	
307	4	4	44	18	0.151	0.121	0.117	0.133	0.123	0.0988	5	5	1	
605	4	5	52	12	0.14	0.103	0.097	0.094	0.089	0.1046	5	5	1	
101	3	4	34	9	0.169	0.125	0.116	0.078	0.077	0.113	3	5	0	
104	4	4	41	12	0.143	0.101	0.084	0.094	0.06	0.0964	5	5	0	
1110	4	3	47	10	0.12	0.14	0.115	0.158	0.146	0.1358	5	5	0	
408	3	3	40	10	0.121	0.097	0.141	0.082	0.117	0.1116	5	5	0	
105	3	3	45	8	0.149	0.095	0.078	0.107	0.081	0.102	5	5	0	
401	dead									0				
402	lodged									0	5	5		
706	dead									0				
507	very small									0	5	5		
411	dead									0				
511	dead									0				
312	dead									0				
113	dead									0				
114	dead									0				
214	dead									0				
314	dead									0				
414	dead									0				
514	dead									0				
1614	dead									0				

Study: MOPMC-P-0822-PA,WL

Study Title: Inter Center Strain Trial – Yield and Persistence of 11 Big Bluestem Sources in Kansas, Missouri, Arkansas, and Mississippi

Study Leader: Cordsiemon, R., Elsberry Plant Materials Center

Introduction:

Big bluestem cultivars developed by NRCS and ARS, and NRCS prevarietal releases and selections will be evaluated for yield and persistence in replicated plots in Booneville, AR; Coffeerville, MS; Elsberry, MO and Manhattan, KS. There have been several prevarietal releases of big bluestem made in recent years by Booneville, AR and Elsberry, MO PMCs. Comparative evaluations of these prevarietal releases and a selection from the Manhattan, KS PMC are needed to further document their performance and adaptation in other geographical regions. Information gained from these plantings may be used to provide data to support elevating lower class releases (e.g. source id and selected class) to a higher release category (e.g. tested class or cultivar).

In addition to these releases, standard big bluestem cultivars commonly used in NRCS conservation plantings and programs will be included in the trial along with other cultivars developed by the USDA-ARS, Lincoln, NE.

Procedure:

Big bluestem entries will be established in 6-ft x 9-ft plots at PMCs in Booneville, AR; Elsberry, MO; Coffeerville, MS; and Manhattan, KS in 2008. All PMCs, except Coffeerville, will send a packet of seed of their respective entry to participating PMCs (see table above). Participating PMC will start seed in the greenhouse and grow out approximately 100 seedlings of each entry. Plants will be transplanted in the field in late spring or early summer using the same plot layout on page 6.

Elsberry will provide live propagules of Refuge germplasm for each location (72 plants x 4 locations = 288 plus 15 -25 extras per location for replacements).

Evaluations will start when 50% of the first entry reaches the late boot stage (1st seed head emerging) of growth. Height measurements will be recorded from the center 4 plants from each of the four replications. Yield will be determined by sampling four plants from the center of each row with a cutting height of 8 inches. The average plot weight (average weight of the 4 plants) will be determined for each big bluestem entry by replication and a grab sample collected for dry matter. After harvest, all remaining plants will be cut and removed from the plot. A second harvest will be made within 2-4 weeks after the first killing frost. The average plant height of entry in each of the four replications from the center 4 plants will be measured and recorded.

List of entries included in the ICST.

Name/Accession#	Source	PMC Responsible For Securing and Distributing Seed to Participating PMCs
Hampton Germplasm (9056854)	Booneville	Randy King
OZ-70 Germplasm (9078831)	Elsberry	Steve Bruckerhoff
Refuge Germplasm (9078832)	Elsberry	Steve Bruckerhoff
9083274	Elsberry	Steve Bruckerhoff
Local ecotype (9079000)	Elsberry	Steve Bruckerhoff
Rountree (474216)	Elsberry	Steve Bruckerhoff
Kaw (421276)	Manhattan	Rich Wynia
483446	Manhattan	Rich Wynia
Pawnee	Stock Seed	Rich Wynia
Bonanza	Stock Seed	Rich Wynia
Goldmine	Sharp Bros.	Rich Wynia

Discussion:

2008

In the spring of 2008, the 4 Plant Materials Centers provided plant material of the big bluestem that they were working with or were responsible for. Seedling plugs were provided in most cases. In 2008, the evaluation plot was given a year to establish and evaluations are to start in 2009.

2009

The big bluestem ICST plot was fertilized with 50 lb. per acre (8 lb. bulk for the plot) and 100 lb. of pellet lime on April 21, 2009. On July 31 and November 3, the plots were cut and 4 plants were weighed and calculated to determine the amount of forage per acre. Grab samples were taken to determine dry matter yield. The July 31 represents the first cutting and November 3 represents the amount of re-growth. Also on those dates, the plots were evaluated for plant height, vigor, and disease and insect resistance. See table #1 for a summary of the evaluations for 2009 at the Elsberry PMC.

2010

On July 13 and November 15, the plots were cut and 4 plants were weighed and calculated to determine the amount of forage per acre. Grab samples were taken to determine dry matter yield. The July 13 represents the first cutting and November 15 represents the amount of re-growth. Also on those dates, the plots were evaluated for plant height, vigor, and disease and insect resistance. See table #2 for a summary of the evaluations for 2010 at the Elsberry PMC.

Summary of Data for Big Bluestem Inter-Center Strain Trial for 2009 Table #1

Release Name	Accession	Dry Material			Vigor		Height	
		7/31/09 Tons/Acre	11/3/09 Tons/Acre	Total Wt Tons/Acre	7/31/2009	11/3/2009	7/31/2009	11/3/2009
Northern	9079000	4.9	0.3	5.17	2.5	3.5	7.7	1.2
OZ-70	9078831	4.8	0.3	5.12	2.0	1.5	6.9	1.7
Bonanza	641701	4.7	0.2	4.96	3.0	5.0	6.9	0.9
Hampton	9056854	4.4	0.2	4.61	1.0	4.5	7.1	1.1
Epic	9083274	4.2	0.3	4.55	2.0	2.5	6.7	1.6
Goldmine	641702	4.2	0.2	4.37	1.5	4.5	6.8	1.0
Rountree	474216	4.1	0.2	4.29	4.0	3.5	7.7	1.1
Kaw	421276	4.1	0.2	4.29	3.0	3.5	6.4	1.1
Kansas	483446	3.6	0.2	3.80	2.5	4.0	6.0	1.2
Refuge	9078832	2.9	0.1	3.01	1.5	3.5	5.9	1.1
Pawnee	9055679	2.5	0.2	2.72	4.0	5.0	6.6	0.9

Vigor Ratings 1=Excellent 3=Good 5=Fair 7=Poor

Summary of Data for Big Bluestem Inter-Center Strain Trial for 2010										Table #2	
Release Name	Accession	Dry Material			Total Wt Tons/Acre	Vigor		Height (ft)			
		7/13/10 Tons/Acre	11/15/10 Tons/Acre	11/15/10 Tons/Acre		7/13/2010	11/15/2010	7/13/2010	11/15/2010		
Northern Missouri Ecotype	9079000	3.91	1.42	5.33		2.00	3.00	4.43	2.06		
OZ-70											
Germplasm	9078831	2.43	1.66	4.09		2.00	2.00	3.85	2.10		
Bonanza	641701	2.86	0.61	3.47		5.00	4.00	3.50	1.75		
Hampton											
Germplasm	9056854	2.81	1.21	4.02		2.00	1.50	3.78	6.90		
Epic											
Germplasm (not released)	9083274	2.04	1.31	3.35		2.00	1.00	3.55	2.52		
Goldmine	641702	3.17	1.44	4.61		1.50	3.00	3.83	2.06		
Rountree	474216	3.12	0.94	4.06		4.00	2.50	4.18	2.00		
Kaw	421276	1.59	0.72	2.32		3.50	5.00	3.28	1.38		
Kansas											
Selection	483446	2.23	1.29	3.52		4.50	3.00	3.33	2.23		
Refuge											
Germplasm	9078832	3.26	0.82	4.08		1.00	1.00	4.18	1.98		
Pawnee	9055679	1.63	0.41	2.03		4.50	4.00	3.45	1.69		

Vigor Ratings 1=Excellent 3=Good 5=Fair 7=Poor

Herbaceous and Woody Seed and Plant Production at the Elsberry PMC 2010

The plant and seed inventory at the Elsberry PMC is used for field plantings, special plantings, demonstration plantings, research studies and commercial release.

Name	Seed Harvested in 2009	Seed Harvested in 2010
Herbaceous Species	Bulk (Pounds)	Bulk (Pounds)
<u>'Rountree' big bluestem</u> <i>Andropogon gerardii</i>	310.5 Foundation	149.4 Foundation
<u>Refuge Big Bluestem</u> (9078832) <i>Andropogon gerardii</i>	0.5 Re-establishing Plot	1.0 Re-establishing Plot
<u>'Rumsey' Indiangrass</u> <i>Sorghastrum nutans</i>	43.0 Foundation	0 Did Not Harvest
<u>'Cave-In-Rock' switchgrass</u> <i>Panicum virgatum</i>	1470.9 Foundation	0 Did Not Harvest – Poor Seed
<u>'OH-370' big bluestem</u> <i>Andropogon gerardii</i>	0 Did Not Harvest	0 Did Not Harvest
<u>'OZ-70' big bluestem</u> <i>Andropogon gerardii</i>	265.7 SG1	416.5 SG1
<u>'Alexander' tick trefoil</u> <i>Desmodium canadense</i>	0 SG1	1.0 SG1
<u>Flood tolerant switchgrass</u> <i>Panicum virgatum</i>	1.2 Re-establishing Plot	0 Destroyed Seed - Poor
<u>Low growing switchgrass</u> <i>Panicum virgatum</i>	3.4 Re-establishing Plot	0 Destroyed Seed - Poor
<u>'Bobwhite' soybean</u> <i>Glycine spp.</i>	0 Destroyed by Flood	0 Destroyed by Flood
<u>Cuivre River Virginia Wildrye</u> <i>Elymus virginicus</i>	0 Destroyed Seed - Poor	0 Did Not Harvest – Poor Seed
<u>Iowa Ecotype Plantings</u> (6 Species; 7 Plots)	6.0 Total	30.4 Total
<u>Missouri Ecotype Plantings</u> (6 Species; 12 Plots)	446.7 Total	539.4 Total

Herbaceous and Woody Seed and Plant Production – continued

Name:	Seed Harvested & Plant Inventory as of December 2009		Seed Harvested & Plant Inventory as of December 2010	
	Plants	Seed Bulk (Pounds)	Plants	Seed Bulk (Pounds)
Woody Species				
<u>'Union' tulip tree</u> <i>Liriodendron tulipifera</i> (9055584)	0	0	0	0
<u>American hazelnut</u> (9083247) (Composite) <i>Corylus americana</i>	0	11	0	10
<u>American plum</u> (9083241) (Composite) <i>Prunus Americana</i>	0	8.8	25	47.3
<u>'Redstone' Cornelian cherry</u> (9055585) <i>Cornus mas</i>	0	0	0	1.8
<u>Chokecherry</u> (9083259) <i>Prunus virginicus</i>	0	0	0	0
<u>Missouri Covey False Indigo Bush</u> <i>Amorpha fruticosa</i> (9083248)	0	6.1	50	0
<u>Iowa Covey False Indigo Bush</u> <i>Amorpha fruticosa</i> (9083249)	0	8	50	0
<u>Illinois Covey False Indigo Bush</u> <i>Amorpha fruticosa</i> (9083250)	0	8.7	50	0

FY2010 PLANT RELEASE
UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE
ELSBERRY, MISSOURI

**NOTICE OF RELEASE OF OZARK LITTLE BLUESTEM
SELECTED CLASS OF NATURAL GERMPLASM**

The Natural Resources Conservation Service (NRCS), U.S. Department of Agriculture, announces the release of a selected class of little bluestem (*Schizachyrium scoparium*, Michx.) for southern counties in Missouri and Illinois.

As a selected class release, this plant will be referred to as Ozark Germplasm little bluestem to document its original area of collections. Ozark Germplasm little bluestem is released as a selected class release of certified seed (genetically manipulated). It has been assigned the NRCS accession number 9083271.

This alternative release procedure is justified because there are no existing commercial sources of little bluestem collected from numerous native sites throughout this specific region. Propagation material of specific ecotypes is needed for conservation programs, wildlife cover and forage for livestock, as well as roadside plantings and prairie restoration and enhancement. The potential for immediate use is high.

Collection Site Information: Collections were taken from native prairie remnants within the counties in Missouri south of the Missouri River and southern Illinois in the Ozark region. Vegetative materials were collected from 51 native sites throughout the two states. A minimum of three collections per Major Land Resource Area for each state was requested. Three Major Land Resource Areas (MLRA's) were represented. Ozark Highland, (116A) had collections from 6 counties of Perry, Cape Girardeau, Ripley, Taney, Miller, and 2 from Maries. Cherokee Prairies, (112) had collections from Barton and 2 collections from Pettis. Southern Illinois Claypan Area (113) had 1 collection from Wayne County.

Ecotype Description: Ozark Germplasm fits within the description of little bluestem. Little bluestem is a medium tall, warm season, bunch type grass with coarse stems at basal leaves (Hughes, Heath and Metcalf, 1951). As a perennial it begins growth in late spring and continues through the hot summer period until the first killing frost. It is easily mistaken for common broomsedge, *Andropogon virginicus*. Plants are slender to robust, compressed, 50 to 150 cm. tall, erect, the upper half freely branching; sheaths and blades commonly glabrous or nearly so, frequently sparsely pilose at their junction, rarely pubescent to villous throughout, the blades 3 to 6 mm. wide, flat; raceme 3 to 6 mm. long, mostly curved, the filiform peduncles mostly wholly or partly included in the sheaths, commonly spreading, the rachis slender, flexuous, pilose, sometimes copiously so; sessile spikelet mostly 6 to 8 mm. long, scabrous, the awn 8 to 15 mm. long; pedicellate spikelet usually reduced, short-awned, spreading, the pedicel pilose (Steyermark, 1968). It develops full stands where moisture is sufficient, but gets clumpy on drier sites. It has value as a persistent low maintenance cover plant and as summer forage.

Collections of little bluestem from east to west across Missouri improves the adaptation of the release to the entire zone.

Method of Selection: The eleven collections that comprise Ozark Germplasm little bluestem (Accession 9083271) were selected from a collection of 46 accessions of little bluestem originating in Missouri and Illinois (Table 1). These 46 accessions included 12 plants each for a total 612 plants in the evaluation. Eighty-seven of the 612 were selected for forage quality analysis and 17 plants were selected that comprise accession 9079130 (Table 2). Plants were selected for forage quality and apparent quantity, and regrowth potential following defoliation, late maturity, vigor, and quick seedling emergence. Initial evaluation selection data was taken in 1999-2000 (cycle 1). The eleven selections were placed in a poly cross block and seed harvested was used to start cycle 2. Selections from cycle 1 were based on quick emerging progeny that were placed into a new evaluation nursery. Selections from this evaluation nursery (cycle 2) comprise the breeders (G0) block of accession 9083271.

Ecological Considerations and Evaluation: Ozark Germplasm little bluestem is a collection of naturally occurring germplasm. Ozark Germplasm little bluestem did not meet the assessment of a plant that could become invasive based on guidelines adopted by the NRCS Plant Materials Program. Ozark Germplasm little bluestem was “okay to release” when evaluated through the “Worksheet for Conducting an Environmental Evaluation of Plant Materials Releases”.

Anticipated Conservation Use: Ozark Germplasm can be used in a range seeding, a pasture seeding, prairie restoration, prairie landscaping, wildlife cover and roadside restoration.

Field Testing: In 2008 a field planting was done in Osage County in MLRA 116A, Ozark Highland area, to test the performance in comparison between 9083271 and Aldous little bluestem. The first year performance on seedling vigor was good for 9083271 with some seedlings obtaining 1 foot in height, averaging 6 inches in height, compared to Aldous with a fair vigor and seedlings only averaging 4 inches in height. In 2009, the field planting was evaluated for stand density and height. Accession 9083271 densities were 80% and a height of 5 feet compared to Aldous density of 40% and a height of 4 feet.



In 2009, a field planting was conducted in MLRA 115, Central Mississippi Wooded Slopes, in Lincoln County. The planting was a 1 acre field border practice. This planting comparison was between 9083271 little bluestem on 0.5 acre to 0.5 acre of Aldous little bluestem. The field border was planted into existing crop residue and annual vegetation, which had glyphosate applied at 2qts/acre on April 22, 2009. The two little bluestems were planted, using a no-till drill, on 0.5 acres each on June 2, 2009, followed by an application of imazapic at 8 oz/acre after planting. By mid July, 9083271 had obtained 6 inches in height, with rows emerging within good

weed control in the field border. Aldous was 3 inches in height with rows, however emergence was spotty; again weed control was good in the field border. In November 2009 after the growing season, stand density counts results for 9083271 average 62% stand compared to Aldous with 39% stand.

Potential Area of Adaptation: Ozark Germplasm was collected from a wide variety of soils, so there is a potential for a broad range of adaptation. It grows well on deep to shallow, sandy to fine-textured and rocky soils. It tolerates low rainfall and is often found on droughty sites. It is not tolerant to heavy shade.

Little bluestem occurs in prairies, glades, rocky open woods, abandoned fields, sandy open ground, waste places, and along railroads (Hitchcock, 1951). Ozark Germplasm originated throughout the southern counties of the Missouri Ozarks within portions of MLRA's 112, 114, 116a and 116b. It is found in USDA plant hardiness zones 5 and 6.

Availability of Plant Materials: G1 material is being produced in limited supply by the Elsberry Plant Materials Center. For information contact USDA, NRCS, Plant Materials Center, 2803 N. Hwy 79, Elsberry, Missouri 63343 (573 898-2012).

References:

Flora of Missouri; p. 244 (Seventh Printing, 1996), Steyermark, J. A; Iowa State University Press, Ames, IA 1968.

Manual of Grasses of the United States; pp. 753-755, Hitchcock, A.S., United States Department of Agriculture, Washington, DC, 1951.


Forages; p. 514, Hughes, H.D.; Heath, M.E.; Metcalfe, D.S., The Iowa State College Press, Ames, Iowa, 1951.

Prepared by:

Ron Cordsiemon, USDA NRCS Plant Materials Center, 2803 North Hwy 79, Elsberry, Missouri, 63343.

Signatures for release of:

Ozark Germplasm little bluestem (*Schizachyrium scoparium*, Michx.)




J.R. Flores
State Conservationist
United States Department of Agriculture
Natural Resources Conservation Service
Columbia, Missouri

4-28-10
Date



William J. Gradle
State Conservationist
United States Department of Agriculture
Natural Resources Conservation Service
Champaign, Illinois

5-20-2010
Date



Richard Sims
State Conservationist
United States Department of Agriculture
Natural Resources Conservation Service
Des Moines, Iowa

05/07/2010
Date

Michael Hubbs
Director, Ecological Sciences Division
United States Department of Agriculture
Natural Resources Conservation Service
Washington, D.C.

Date

Table 1. Initial collection of little bluestem accessions in Missouri and Illinois. Accessions in *italic* comprise Ozark Germplasm USDA-NRCS Elsberry, MO.

ACCESSION	REFERENCE		MLRA	COUNTY	STATE
	NUMBER	COLLECTOR			
<i>9078895</i>	<i>MO-3</i>	<i>Joe Tousignant</i>	<i>N116B</i>	<i>Cape Girardeau</i>	<i>Missouri</i>
<i>9078899</i>	<i>MO-7</i>	<i>Tommy Robins/ Jim Hofer</i>	<i>116</i>	<i>Ripley</i>	<i>Missouri</i>
9078900	MO-8	Grant P. Butler	N116B	Jefferson	Missouri
9078901	MO-9			Iron	Missouri
9078903	MO-11	Arch J. Mueller	M115	Ste. Genevieve	Missouri
9078904	MO-12			St. Francois	Missouri
9078905	MO-13	J. Mark Mitchell		Butler	Missouri
9078906	MO-14	Randy C. Miller	N116A	Shannon	Missouri
9078907	MO-15	Tom Johnson	N116B	Bollinger	Missouri
9078908	MO-16	Tom Johnson	N116A	Bollinger	Missouri
9078909	MO-17	Randy C. Miller	N116B	Reynolds	Missouri
9078911	MO-19	Tom Johnson	N116A	Wayne	Missouri
9078912	MO-20	Mark E. Nussbaum	N116B	Cape Girardeau	Missouri
<i>9078915</i>	<i>MO-23</i>	<i>Claude F. Peifer</i>	<i>116B</i>	<i>Perry</i>	<i>Missouri</i>
9078916	MO-24	Grant P. Butler/ Bryan L. Westfall	N116A	Washington	Missouri
9078925	MO-33	Gary J. Barker	M109	Gentry	Missouri
9078926	MO-34			Vernon	Missouri
9078928	MO-36	Todd E. Mason	M109	Worth	Missouri
<i>9078942</i>	<i>MO-50</i>	<i>Ian S. Kurtz</i>	<i>116A</i>	<i>Taney</i>	<i>Missouri</i>
9078943	MO-52	Dennis Shirk/ Ed Gillmore	115	Gasconade	Missouri
9078944	MO-53	Dennis Shirk/ Ed Gillmore	116	Osage	Missouri
9078945	MO-54	Raleigh Redman	112	Henry	Missouri
9078946	MO-55	Dennis Shirk/ Ed Gillmore	116	Maries	Missouri
9078947	MO-56	Jerry Cloyed	M112	Barton	Missouri
9078948	MO-57	Ian S. Kurtz	116A	Taney	Missouri
9078949	MO-58	Ben A. Reed	M112	Barton	Missouri
<i>9078950</i>	<i>MO-59</i>	<i>Jerry Cloyed</i>	<i>M112</i>	<i>Barton</i>	<i>Missouri</i>
<i>9078952</i>	<i>MO-60</i>	<i>M. Denise Brown</i>	<i>N116A</i>	<i>Miller</i>	<i>Missouri</i>
9078953	MO-61	M. Denise Brown	N116B	Miller	Missouri
9078954	MO-62	Howard L. Coambes	N116B	Cedar	Missouri
9078955	MO-63	Howard L. Coambes	N116B	Cedar	Missouri
9078958	MO-66	Rod Doolen		Wayne	Missouri
9078959	MO-67	Rod Doolen		Wayne	Missouri
9078963	MO-69	Maurice Davis/ Steve Clubine		Pettis	Missouri
	MO-71	Maurice Davis/ Steve Clubine		St. Clair	Missouri
	MO-72	Maurice Davis/ Steve Clubine		Benton	Missouri

Table 1 (con't). Initial collection of little bluestem accessions in Missouri and Illinois. Accessions in italic comprise Ozark Germplasm USDA-NRCS Elsberry, MO.

ACCESSION	REFERENCE NUMBER	COLLECTOR	MLRA	COUNTY	STATE
<i>9078964</i>	<i>MO-73</i>	<i>Maurice Davis/</i>		<i>Pettis</i>	<i>Missouri</i>
<i>9078965</i>	<i>MO-74</i>	<i>Maurice Davis/</i> Steve Clubine		<i>Pettis</i>	<i>Missouri</i>
9078966	MO-77	Maurice Davis/ Steve Clubine		Maries	Missouri
9078967	MO-78	Dennis Shirk		Maries	Missouri
<i>9078968</i>	<i>MO-79</i>	<i>Steve Clubine</i>		<i>Maries</i>	<i>Missouri</i>
<i>9078969</i>	<i>MO-80</i>	<i>Maurice Davis</i>		<i>Maries</i>	<i>Missouri</i>
9078970	MO-81			Lawrence	Missouri
9078889	IL-17	Michael Stanfill/	115	Monroe	Illinois
9078892	IL-20	Michael Stanfill/ Marty Kemper	113	Washington	Illinois
<i>9078893</i>	<i>IL-21</i>	<i>Remington T. Irwin</i>	<i>114</i>	<i>Wayne</i>	<i>Illinois</i>

Table 2. Plant performance ratings and forage quality estimates of accessions comprising Ozark Germplasm little bluestem 1999-2000, USDA-NRCS Elsberry, MO.

Plant Number ¹	1999						2000		
	Aug Forage Rating ²	Aug Vigor Rating ³	First Seedhead ⁴	Quality Rating			Oct Regrowth Rating ⁸	Jun Forage Rating ⁹	Jun Vigor Rating ¹⁰
				% N ⁵	% CP ⁶	% ADF ⁷			
MO-23	3	5	6-Sep	2.1	13	33	4	1	2
IL-21	3	3	12-Sep	2	13	34	3	4	3
IL-21	4	5	8-Sep	2	13	33	2	3	3
MO-23	2	4	27-Aug	2	13	34	1	1	3
MO-3	4	3	26-Aug	2	13	35	3	3	2
MO-79	3	3	14-Sep	1.9	12	37	2	1	3
MO-3	7	4	13-Sep	1.9	12	34	3	3	2
MO-7	1	2	10-Sep	1.9	12	36	3	2	2
MO-60	3	3	8-Aug	1.9	12	38	3	2	2
IL-21	4	7	7-Sep	1.8	11	36	3	4	3
MO-7	1	2	25-Aug	1.8	11	36	4	1	1
MO-73	2	3	25-Jul	1.8	11	36	2	1	2
MO-59	5	4	27-Jul	1.7	10	35	2	2	2
MO-50	2	4	10-Sep	1.6	10	37	1	1	2
MO-80	1	3	28-Jul	1.6	10	35	2	1	2
MO-50	3	5	15-Sep	1.5	9	37	2	1	3
MO-74	1	4	12-Aug	1.5	9	36	2	1	3

1)Plant Number describes the collection the plant originated from; 2)Forage Rating and 3)Vigor Rating are a subjective rating given to each individual plant based on their appearance in Aug 1999; 4)First Seedhead is the date the recorded when the initial seedheads were noticed; Quality Rating – 5)% Nitrogen, 6)% Crude Protein, and 7)% ADF (Acid Detergent Fiber) were measured from grab samples taken from each individual plant; 8)Regrowth Rating is a subjective rating taken in October 2000 based on the amount of regrowth present after an early forage harvest; 9)Forage Rating and 10)Vigor Rating are a subjective rating given to each individual plant based on their appearance in June 2000. Ratings are based on a scale of 1-9. 1 = Excellent and 9 = Poor

Releases From Elsberry Plant Materials Center						
Yearly Release						
Scientific Name	Release Name	Common Name	Accession Number	Secondary Agency(ies)	Type of Release	Year of Release
<i>Panicum virgatum</i> L.	Cave-In-Rock	switchgrass	469228	MOA	N	1974
<i>Glycine</i> sp. L	** Bobwhite	soybean	421822	MOPMC,ARS, MOA,	I	1975
<i>Acer ginnala</i> Maxim.	Flame	Amur maple	483442		I	1978
<i>Andropogon gerardii</i> Vitman.	Rountree	big bluestem	474216	MOA	N	1983
<i>Sorghastrum nutans</i> (L.) Nash.	Rumsey	Indiangrass	315747	MOA	N	1983
<i>Ulmus parvifolia</i> Jacq.	Elsmo	lace bark elm	9004438		I	1990
<i>Cornus mas</i> L.	Redstone	cornelian cherry dogwood	516476		I	1991
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Central Iowa	sideoats grama	9062279	UNI,IARV,IAT,ICIA	N	1995
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Northern Iowa	sideoats grama	9062278	UNI,IARV,IAT,ICIA	N	1995
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Southern Iowa	sideoats grama	9062280	UNI,IARV,IAT,ICIA	N	1995
<i>Elymus canadensis</i> L.	Northern Iowa	Canada wildrye	9062275	UNI,IARV,IAT,ICIA	N	1995
<i>Elymus canadensis</i> L.	Central Iowa	Canada wildrye	9062276	UNI,IARV,IAT,ICIA	N	1995
<i>Elymus canadensis</i> L.	Southern Iowa	Canada wildrye	9062277	UNI,IARV,IAT,ICIA	N	1995
<i>Heliopsis helianthoides</i> (L.) Sweet	Central Iowa	oxeye false sunflower	9068606	UNI,IARV,IAT,ICIA	N	1995
<i>Panicum virgatum</i> L.	* Shawnee	switchgrass	591824		N	1995
<i>Heliopsis helianthoides</i> (L.) Sweet	Northern Iowa	oxeye false sunflower	9068605	UNI,IARV,IAT,ICIA	N	1996
<i>Lespedeza capitata</i> Michx.	Central Iowa	roundhead lespedeza	9062282	UNI, IARV, IAT, ICIA	N	1996
<i>Sorghastrum nutans</i> (L.) Nash	Central Iowa	Indiangrass	9062317	UNI,IARV,IAT,ICIA	N	1996
<i>Sorghastrum nutans</i> (L.) Nash	Northern Iowa	Indiangrass	9062316	UNI,IARV,IAT,ICIA	N	1996
<i>Sporobolus compositus</i> (Poir.) Merr.	Central Iowa	tall dropseed	9062314	UNI,IARV,IAT,ICIA	N	1996
<i>Andropogon gerardii</i> Vitman.	OH-370	big bluestem	9062323	ARPMC	N	1997
<i>Desmodium canadense</i> L.	Alexander	showy tick trefoil	9057110		N	1997
<i>Heliopsis helianthoides</i> (L.) Sweet	Southern Iowa	oxeye false sunflower	9068607	UNI,IARV,IAT,ICIA	N	1997
<i>Lespedeza capitata</i> Michx.	Southern Iowa	roundhead lespedeza	9062283	UNI, IARV, IAT, ICIA	N	1997
<i>Liriodendron tulipifera</i> L.	Union	tulip poplar	9055584		N	1997
<i>Schizachyrium scoparium</i> , Michx.	Central Iowa	little bluestem	9062320	UNI,IARV,IAT,ICIA	N	1997
<i>Andropogon gerardii</i> Vitman.	Central Iowa	big bluestem	9068615	UNI,IARV,IAT,ICIA	N	1998
<i>Dalea purpurea</i>	Central Iowa	purple prairie clover	9068609	UNI,IARV,IAT,ICIA	N	1998
<i>Eryngium yuccifolium</i> Michx.	Northern Iowa	rattlesnake master	9068602	UNI,IARV,IAT,ICIA	N	1998
<i>Solidago rigida</i> L.	Northern Iowa	rigid goldenrod	9068617	UNI,IARV,IAT,ICIA	N	1998
<i>Sorghastrum nutans</i> (L.) Nash.	Southern Iowa	Indiangrass	9062318	UNI,IARV,IAT,ICIA	N	1998
<i>Andropogon gerardii</i> Vitman.	Southern Iowa	big bluestem	9068616	UNI, IARV, IAT, ICIA	N	1999
<i>Andropogon gerardii</i> Vitman.	Northern MO	big bluestem	9079000	UMC,MDC,MODOT	N	1999
<i>Elymus virginicus</i> L.	Northern MO	Virginia wildrye	9079044	UMC,MDC,MODOT	N	1999
<i>Eryngium yuccifolium</i> Michx.	Southern Iowa	rattlesnake master	9068604	UNI, IARV, IAT, ICIA	N	1999
<i>Eryngium yuccifolium</i> Michx.	Central Iowa	rattlesnake master	9068603	UNI, IARV, IAT, ICIA	N	1999
<i>Liatris pycnostachya</i> , Michx	Northern Iowa	prairie blazing star	9068626	UNI, IARV, IAT, ICIA	N	1999
<i>Liatris pycnostachya</i> , Michx	Central Iowa	prairie blazing star	9068627	UNI, IARV, IAT, ICIA	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Northern Iowa	little bluestem	9062319	UNI, IARV, IAT, ICIA	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Southern Iowa	little bluestem	9962321	UNI, IARV, IAT, ICIA	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Northern MO	little bluestem	9079004	UMC,MDC,MODOT	N	1999
<i>Sorghastrum nutans</i> (L.) Nash.	Northern MO	Indiangrass	9079036	UMC,MDC,MODOT	N	1999
<i>Sorghastrum nutans</i> (L.) Nash.	Western MO	Indiangrass	9079037	UMC,MDC,MODOT	N	1999
<i>Andropogon gerardii</i> Vitman.	Northern Iowa	big bluestem	9068614	UNI,IARV,IAT,ICIA	N	2000
<i>Lespedeza capitata</i> Michx.	Northern Iowa	roundhead lespedeza	9062284	UNI, IARV, IAT, ICIA	N	2000
<i>Liatris pycnostachya</i> , Michx	Southern Iowa	prairie blazing star	9068628	UNI, IARV, IAT, ICIA	N	2000
<i>Sporobolus compositus</i> (Poir.) Merr.	Northern Iowa	tall dropseed	9062313	UNI, IARV, IAT, ICIA	N	2000
<i>Coreopsis Palmata</i> Nutt.	Northern MO	prairie coreopsis	9079028	MDC, NAS	N	2001
<i>Coreopsis Palmata</i> Nutt.	Western MO	prairie coreopsis	9079029	MDC, NAS	N	2001
<i>Liatris pycnostachya</i> , Michx	Northern MO	prairie blazing star	9079020	MDC, NAS	N	2001
<i>Liatris pycnostachya</i> , Michx.	Western MO	prairie blazing star	9079021	MDC, NAS	N	2001
<i>Sporobolus compositus</i> var. comp.	Northern MO	tall dropseed	9079040	MDC, NAS	N	2001
<i>Aster novae-angliae</i> L.	Central Iowa	New England Aster	9068682	UNI, IARV, IAT, ICIA	N	2002
<i>Aster novae-angliae</i> L.	Northern Iowa	New England Aster	9068681	UNI, IARV, IAT, ICIA	N	2002
<i>Aster novae-angliae</i> L.	Southern Iowa	New England Aster	9068683	UNI, IARV, IAT, ICIA	N	2002
<i>Echinacea pallida</i> Nutt.	Northern Iowa	pale purple coneflower	9068611	UNI, IARV, IAT, ICIA	N	2002
<i>Echinacea pallida</i> Nutt.	Southern Iowa	pale purple coneflower	9068613	UNI, IARV, IAT, ICIA	N	2002
<i>Elymus virginicus</i> L.	Cuivre River	Virginia wildrye	9803169	MDC	N	2002
<i>Solidago rigida</i> L.	Southern Iowa	rigid goldenrod	9068619	UNI, IARV, IAT, ICIA	N	2002

Releases From Elsberry Plant Materials Center						
Alphabetically						
Scientific Name	Release Name	Common Name	Accession Number	Secondary Agency(ies)	Type of Release	Year of Release
<i>Acer ginnala</i> Maxim.	Flame	Amur maple	483442		I	1978
<i>Amorpha fruticosa</i> L.	Missouri Covey False Indigo Bush	false indigo bush	9083248		N	2005
<i>Amorpha fruticosa</i> L.	Iowa Covey False Indigo Bush	false indigo bush	9083249		N	2005
<i>Amorpha fruticosa</i> L.	Illinois Covey False Indigo Bush	false indigo bush	9083250		N	2005
<i>Andropogon gerardii</i> L.	OZ-70	big bluestem	9078831		N	2004
<i>Andropogon gerardii</i> Vitman.	Northern Iowa	big bluestem	9068614	UNI,IARV,IAT,ICIA	N	2000
<i>Andropogon gerardii</i> Vitman.	Southern Iowa	big bluestem	9068616	UNI, IARV, IAT, ICIA	N	1999
<i>Andropogon gerardii</i> Vitman.	Northern MO	big bluestem	9079000	UMC,MDC,MODOT	N	1999
<i>Andropogon gerardii</i> Vitman.	Central Iowa	big bluestem	9068615	UNI,IARV,IAT,ICIA	N	1998
<i>Andropogon gerardii</i> Vitman.	OH-370	big bluestem	9062323	ARPMC	N	1997
<i>Andropogon gerardii</i> Vitman.	Rountree	big bluestem	474216	MOA	N	1983
<i>Andropogon gerardii</i> Vitman.	Refuge	big bluestem	9078832		N	2006
<i>Aster novae-angliae</i> L.	Central Iowa	New England Aster	9068682	UNI, IARV, IAT, ICIA	N	2002
<i>Aster novae-angliae</i> L.	Northern Iowa	New England Aster	9068681	UNI, IARV, IAT, ICIA	N	2002
<i>Aster novae-angliae</i> L.	Southern Iowa	New England Aster	9068683	UNI, IARV, IAT, ICIA	N	2002
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Central Iowa	sideoats grama	9062279	UNI,IARV,IAT,ICIA	N	1995
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Northern Iowa	sideoats grama	9062278	UNI,IARV,IAT,ICIA	N	1995
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Southern Iowa	sideoats grama	9062280	UNI,IARV,IAT,ICIA	N	1995
<i>Bouteloua curtipendula</i> (Michx.) Torr.	Northern MO	sideoats grama	9079072		N	2008
<i>Coreopsis Palmata</i> Nutt.	Northern MO	prairie coreopsis	9079028	MDC, NAS	N	2001
<i>Coreopsis Palmata</i> Nutt.	Western MO	prairie coreopsis	9079029	MDC, NAS	N	2001
<i>Cornus mas</i> L.	Redstone	cornelian cherry dogwood	516476		I	1991
<i>Corylus, americana</i> Walt.	Sun Harvest	American hazelnut	9083247		N	2007
<i>Dalea purpurea</i>	Northern Iowa	purple prairie clover	9068608	UNI, IARV, IAT, ICIA	N	2003
<i>Dalea purpurea</i>	Central Iowa	purple prairie clover	9068609	UNI,IARV,IAT,ICIA	N	1998
<i>Desmodium spp.</i>	Northern MO	showy tick trefoil	9079012	MDC	N	2004
<i>Desmodium canadense</i> L.	Alexander	showy tick trefoil	9057110		N	1997
<i>Echinacea pallida</i> Nutt.	Northern Iowa	pale purple coneflower	9068611	UNI, IARV, IAT, ICIA	N	2002
<i>Echinacea pallida</i> Nutt.	Southern Iowa	pale purple coneflower	9068613	UNI, IARV, IAT, ICIA	N	2002
<i>Echinacea pallida</i> Nutt.	Central Iowa	pale purple coneflower	9068612	UNI, IARV, IAT, ICIA	N	2006
<i>Echinacea pallida</i> (Nutt.)	Northern MO	pale purple coneflower	9079032		N	2009
<i>Echinacea pallida</i> (Nutt.)	Western MO	pale purple coneflower	9079033		N	2009
<i>Elymus canadensis</i> L.	Northern Iowa	Canada wildrye	9062275	UNI,IARV,IAT,ICIA	N	1995
<i>Elymus canadensis</i> L.	Central Iowa	Canada wildrye	9062276	UNI,IARV,IAT,ICIA	N	1995
<i>Elymus canadensis</i> L.	Southern Iowa	Canada wildrye	9062277	UNI,IARV,IAT,ICIA	N	1995
<i>Elymus virginicus</i> L.	Cuivre River	Virginia wildrye	9803169	MDC	N	2002
<i>Elymus virginicus</i> L.	Northern MO	Virginia wildrye	9079044	UMC,MDC,MODOT	N	1999
<i>Eryngium yuccifolium</i> Michx.	Southern Iowa	rattlesnake master	9068604	UNI, IARV, IAT, ICIA	N	1999
<i>Eryngium yuccifolium</i> Michx.	Central Iowa	rattlesnake master	9068603	UNI, IARV, IAT, ICIA	N	1999
<i>Eryngium yuccifolium</i> Michx.	Northern Iowa	rattlesnake master	9068602	UNI,IARV,IAT,ICIA	N	1998
<i>Glycine</i> sp. L **	Bobwhite	soybean	421822	MOPMC,ARS, MOA,	I	1975
<i>Heliopsis helianthoides</i> (L.) Sweet	Southern Iowa	oxeye false sunflower	9068607	UNI,IARV,IAT,ICIA	N	1997
<i>Heliopsis helianthoides</i> (L.) Sweet	Northern Iowa	oxeye false sunflower	9068605	UNI,IARV,IAT,ICIA	N	1996
<i>Heliopsis helianthoides</i> (L.) Sweet	Central Iowa	oxeye false sunflower	9068606	UNI,IARV,IAT,ICIA	N	1995
<i>Koeleria macrantha</i>	Central Iowa	prairie Junegrass	9068621	UNI, IARV, IAT, ICIA	N	2003
<i>Koeleria macrantha</i>	Northern Iowa	prairie Junegrass	9068620	UNI, IARV, IAT, ICIA	N	2003
<i>Lespedeza capitata</i> Michx.	Northern Iowa	roundhead lespedeza	9062284	UNI, IARV, IAT, ICIA	N	2000
<i>Lespedeza capitata</i> Michx.	Southern Iowa	roundhead lespedeza	9062283	UNI, IARV, IAT, ICIA	N	1997
<i>Lespedeza capitata</i> Michx.	Central Iowa	roundhead lespedeza	9062282	UNI, IARV, IAT, ICIA	N	1996
<i>Liatris aspera</i> , Michx.	Southern Iowa	rough blazing star	9068686	UNI, IARV, IAT, ICIA	N	2003
<i>Liatris aspera</i> , Michx.	Central Iowa	rough blazing star	9068685	UNI, IARV, IAT, ICIA	N	2003
<i>Liatris aspera</i> , Michx.	Northern Iowa	rough blazing star	9068684	UNI, IARV, IAT, ICIA	N	2003
<i>Liatris pycnostachya</i> , Michx	Northern MO	prairie blazing star	9079020	MDC, NAS	N	2001
<i>Liatris pycnostachya</i> , Michx	Northern Iowa	prairie blazing star	9068626	UNI, IARV, IAT, ICIA	N	1999
<i>Liatris pycnostachya</i> , Michx	Southern Iowa	prairie blazing star	9068628	UNI, IARV, IAT, ICIA	N	2000
<i>Liatris pycnostachya</i> , Michx	Central Iowa	prairie blazing star	9068627	UNI, IARV, IAT, ICIA	N	1999
<i>Liatris pycnostachya</i> , Michx.	Western MO	prairie blazing star	9079021	MDC, NAS	N	2001
<i>Liriodendron tulipifera</i> L.	Union	tulip poplar	9055584		N	1997
<i>Monarda fistulosa</i> L.	Northern Iowa	wild bergamot	9068678	UNI, IARV, IAT, ICIA	N	2007

**-Studies/Projects at the Elsberry Plant Materials Center
Studies 1958 through 2010**

Study/Project Number System: Initially the numbers were assigned numerically plus the year the study/project was initiated. Later a different numbering system was adopted which involved the designated state number, a letter to denote the type of project/study and finally a numerical number.

Study/ Project Number	Title	Annual Technical Report/ Page Reference	
		ATR	Page
2-58	Quaker Comphrey Evaluation	1962	28
3-58	Comparison of Winter Annual Cover Crops	1962	30
6-62	Fertilizer Rate Study on Midland Bermudagrass, <i>Cynodon dactylon</i>	1963	47
10-59	Interseeding Cover Crops in Corn	1963	52
14-61	Evaluation of <i>Lotus corniculatus</i> L. Strains	1966	24
15-61	Evaluation of Bermudagrass Strains	1965	17
17-61	Black Locust, <i>Robinia pseudoacacia</i> L. Trials	1967	35
18-61	The Rate, Date and Method of Seeding <i>Lespedeza daurica schimadae</i>	1962	23
19-61	Living Fence Trials	1968	26
20-61	Plants for Bank stabilization	1962	10
21-62	Evaluation of Legumes for wildlife	1962	11
23-63	Evaluation of <i>Phalaris arundinacea</i> L. 'Ioreed' Reed Canarygrass Strains	1964	13
24-72	Method of Seeding Creeping Foxtail	1962	24
25-63	Advanced Evaluation of Plant Materials for Grass Waterways	1968	27
26-63	Evaluation of Japanese Pagodatree, <i>Sophora japonica</i> , for Posts	1962	16
27-63	Direct Seeding vs. Transplanting Sawtooth Oak, <i>Quercus acutissima</i> , Carruthers	1964	60
28-63	Effect of Cultural Methods on Crownvetch, <i>Coronilla varia</i> L., Seed Production	1964	64
31-63	<i>Lespedeza capitata</i> Michx. – Roundhead Lespedeza Ecotype Evaluation	1964	64
34-63	Cultural Methods for Seeding Grasses in woodland Pastures	1963	58
35-63	Effect of Cultural Methods on Seed Production of <i>Phalaris arundinacea</i> L., 'Ioreed' Reed Canarygrass	1964	13
37-63	Forage Yields and Season of Production for Several Grasses and Legumes Clipped Bi-Weekly at Three Inches and Six Inches	1964	78
38-64	Advanced Evaluation of Perennial Grasses for Summer Pasture	1968	28
42-65	Establishment of Crownvetch and Trefoil in Dead Litter Mulch	1967	41
44-65	Grasses and Legumes for Goose Browse on the Clarence Cannon Wildlife Refuge	1973 Part 1 Part 2	8 44
46-66	Method of Seeding Trials with 'Garrison' Creeping Foxtail	1966	26
49-69	Seed Yield of Three <i>Panicum virgatum</i> , Switchgrass Selections: Mich 381; 'Blackwell', M1-5714, and M1-5845, 'Cave-In-Rock'	1971 Part 1 Part 2	5 46
50-69	Seed Yield and Seed Retention of Four <i>Phalaris arundinacea</i> , Reed Canarygrass Selections – 'Ioreed', 'Rise', 'Frontier', and 'Auburn'	1976	12

Study/ Project Number	Title	Annual Technical Report/ Page Reference	
		ATR	Page
51-C-71	Herbicide tolerance of New Seeding of tall Fescue, Big Bluestem, Indiangrass, and Switchgrass	1979	55
29I052W	Growth Rate Study of European Alder on Deep Alluvial Soil	1980	4
53-72	Growth Rate Study of Poplar (Cottonwood) On A Deep Alluvial Soil	1972 Part 1 Part 2	7 53
54-72	Rhizome Development of Two tall Fescue, <i>Festuca arundinacea</i> , Selections: M1-6161 and M1-6162	1971 Part 1 Part 2	7 54
29A055	Evaluations of <i>Sorghastrum nutans</i> , Indiangrass (M1-7073), Poly-Cross Indiangrass for Leafiness, Disease-Free Characteristics and Seed Production	1981	81
56-71	Comparative Evaluation of New Lotus Accessions with Names and Used Varieties to Determine Potential as a Long Lived Legume in Three State Areas Served	1974 Part 1 Part 2	4 4
29I057-72	Growth Rate Study of Poplars (Cottonwood) on a Deep Alluvial Soil	1981	4
29A058-72	Evaluation For Naming and Releasing of Elsberry Developed Big Bluestem and Indiangrass	1981	83
59-72	Sorghum Evaluation as Wildlife Game Feed	1973 Part 1 Part 2	11 55
29I060-69	Replacement of the American Elm Tree	1979	80
61-72	Advanced evaluation of Meadow Foxtail, <i>Alopecurus pratensis</i> , PI-305495, as Waterway Grass as Compared to 'Garrison' Creeping Foxtail, <i>Alopecurus arundinaceus</i> , the Standard for Comparison	1973 Part 1 Part 2	12 56
29I062J	Trees and Shrubs for Use as Wildlife Food and Cover Plants	1979	11
29I063	Plants for Use in Critical Area Stabilization	1979	21
29I064W	Plants for Wood Products	1979	23
65-78	Plants for Use in Landscape and Beautification	1976	10
29I066W-72	Developing Winter Hardy Nut Bearing Trees and Shrubs for Planting in Parks, Wildlife Areas and Natural Areas	1979	27
29I067K	Trees for windbreaks	1979	29
29A068-72	Response of Yellow Poplar to thinning	1979	67
29A069-72	Black Cherry Demonstration	1979	70
29A070-73	<i>Desmodium</i> for Wildlife Food and Cover	1979	31
29A071-73	Evaluation for Naming and Releasing of Elsberry Developed Autumn Olive, M1-6369	1978	73
29A072-73	Evaluation of M1-4701, <i>Lonicera maackii</i> , Amur Honeysuckle for Naming and Releasing	1978	74
29A073G	Establishment of warm-Season Grasses with Herbicides for Weed Control. Herbicides are Not Tested or Have Label Clearance for Warm-Season Grasses	1979	72
29A139G	Field Evaluation of Establishment of Herbaceous Plant Materials on Sand Covered Flooded Areas in Missouri	1994-1998	149
29A074M	Cover Crops in Soybeans	1984	258
Misc. Study	NJ-927, <i>Eleagnus umbellata</i> , Autumn Olive for wildlife Food and Cover	1981	101

Study/ Project Number	Title	Annual Technical Report/ Page Reference	
		ATR	Page
29A075F	Plants for Shoreline and Wetland Stabilization	1990	64
29I076G-78	Establishment of Warm Season Grasses	1981	7
Misc. Study	Evaluation of Cold Hardy <i>Paspalum notatum</i> Selections	78	76
29I077P	Evaluation of Plants for Vegetating Salt Damaged Areas	1981	11
29I078D	Field Evaluation Planting to Evaluate Species of Plants for Use on Alkali Bearing Soils in Southern Illinois	1981	19
29I079D	Field Evaluation Planting to Evaluate Species of Plants for Use on Revegetating Acid Coal Mine Spoil in Illinois (Saline County SWCD and Peabody Coal Company)	1984	25
29I080D	Field Evaluation Planting to Evaluate Species of Plants for Use in Revegetating Acid Coal Mine Spoil in Iowa (VanBuren County SWCD)	1980	56
29I081D	Field Evaluation Planting to Evaluate Species of Plants for Use in Revegetating Acid Coal Mine Spoil in Iowa (Marion County SWCD)	1980	77
29I082D	Field Evaluation Planting to Evaluate Species of Plants for Use in Revegetating Acid Coal Mine Spoil in Illinois (Fulton County SWCD and Freeman United Coal Mine)	1984	117
29I083M	Legume Cover Crop for No-Till Corn Production	1984	160
29I084G	Legumes to Enhance Fescue Pastures	1986	6
29A085S	Debearding Fluffy Native Grass Seed (Big Bluestem and Indiangrass)	1981	92
29A086L	Use of an Absorbent Polymer in Coating Native Grass Seed	1982	106
29I087D	Plants with Increased tolerance to Aluminum and Manganese	1984	192
29A088W	Cooperative Screening Study of Native and Introduced Sources of Eastern Cottonwood	2000	129
29I089V	Multiple Use Legume Assembly and Evaluation	1988	4
29I090G	No-Till Establishment of Warm-Season Grasses in Cool Season Grass Sod	1984	219
29I091G	Weed Control Treatments for Warm Season Grass Establishment	1988	7
29I092G	Perennial Grasses as Cover Crops for Use in No-Till Systems	1988	12
29I093R	Miscellaneous Grass Evaluation	2010	8
29A094M	Cover Crops in Corn, Soybeans, and Milo	1987	5
29I096F	Streambank Stabilization	1988	14
29I097G	Assembly and Evaluation of Big Bluestem, <i>Andropogon gerardii</i> , Vitman.	2010	13
29I098M	'Tinga' Tangier Pea for Soil Protection	1987	7
29I099J	Assembly and Evaluation of Roughleaf Dogwood, <i>Cornus drummondii</i>	1994-1998	13
29I100J	Assembly and Evaluation of Blackhaw, <i>Viburnum prunifolium</i> L.	1999	17
29I101J	Assembly and Evaluation of Arrowwood, <i>Viburnum dentatum</i>	2006	21
29A102M	Evaluation of Perennial Grass as Cover Crops for No-Till Soybeans	1990	85
29A105M	Evaluation of Winter Annual Grass for Cover Crops in No-Till Soybeans	1993	34
29I107G	Assembly and Evaluation of Eastern Gamagrass, <i>Tripsacum dactyloides</i> L.	2006	24

Study/ Project Number	Title	Annual Technical Report/ Page Reference	
		ATR	Page
29I108G	Assembly and Evaluation of Low Growing Rhizomatous Switchgrass, <i>Panicum virgatum</i> L., for Use in Waterways, Filter Strips and Other Conservation Uses	2010	23
29I109W	Direct Seeding Methods of <i>Quercus</i> sp., Oaks	1993	17
29I110J	Assembly and Evaluation of Chokecherry, <i>Prunus virginiana</i> L.	2007	27
29A111G	Field Evaluation of Selected Perennial Grasses for Pasture Wildlife Habitat and Erosion control (Varietal Study)	1994-1998	91
29I112J	Assembly and Evaluation of Nannyberry, <i>Viburnum lentago</i> L.	1993	21
29I113J	Assembly and Evaluation of Serviceberry, <i>Amelanchier arborea</i> (Michx.F.) Fern.	1993	22
29I114K	Field Evaluation of Woody Plant Materials in Cooperation with Mineral area College	1993	22
29A116W	Field Evaluation of Woody Plant Materials in Cooperation with Mineral Area College	2010	31
29A117H	Intercenter Strain Trial of <i>Tripsacum dactyloides</i> L., Eastern gamagrass	1993	46
29A118G	Field Evaluation of Selected Perennial Grasses for Pasture, Wildlife Habitat and Erosion Control (Varietal Study)	1994-1998	91
29A121W	Conifer Evaluation for Windbreak Plantings	2000	137
29A122G	Evaluation of Perennial Warm-Season Grasses as Windbarriers in Southeast Missouri	1994-1998	125
29A123M	Winter Cover Crop Study for No-Till Soybeans	1993	54
29A124G	Fertility and Harvest Management of Eastern Gamagrass for Forage Production	2010	35
29I126W	Woody Columnar Collection	1993	30
29A127G	Field Evaluation of Selected Perennial Grasses for Pasture, Wildlife Habitat and Erosion Control	1994-1998	91
29A128J	<i>Cornus florida</i> L., Flowering Dogwood, Interagency Study Between Department of Interior, National Parks Service, National Capital Region and the Department of Agriculture	2006	52
29A131O	Treatment of Animal wastewaters by Constructed Wetlands	1993	66
29I132O	Miscellaneous Wetland Plant evaluation	2003	49
29I133J	Assembly and Evaluation of Gray Dogwood, <i>Cornus racemosa</i>	work plan not developed	
29I134J	Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginiana</i> L.	2002	55
29I135J	Assembly and Evaluation of Hazelnut, <i>Corylus Americana</i> , Marsh	2007	43
29I136J	Assembly and Evaluation of Wild Plum, <i>Prunus Americana</i> , Marsh.	2006	54
29A137O	Wetland Riparian Propagation, Establishment and Demonstration	2010	41
29I138G	Residue Decomposition Trial	1994-1998	68
29A139G	Field Evaluation of Establishment of Herbaceous Plants	1994-1998	149
29A140W	Yellow Poplar Evaluation	1994-1998	159
29I141G	Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i> , Michx.	2010	54
29I142G	Production of Native Missouri Ecotypes of Grasses, Legumes and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted	2010	74
29I143G	Seed Coating/Seed Rates	2002	129
29A144G	Biofuel Study of Different Strains/Varieties of Switchgrass	1999	147
29A145	Wear Tolerance Demonstration of Vegetation in High Traffic Areas	2000	154

Study/ Project Number	Title	Annual Technical Report/ Page Reference	
		ATR	Page
MOPMC-P-0001, WO, WL, WE	Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i> , Michx.	2010	80
MOPMC-P-0002, WE, WL	Assembly, Evaluation and Selection of False Indigo Bush, <i>Amorpha fruticosa</i> , L.	2006	127
MOPMC-P-003 PA,WL	Evaluation and Release of Eastern Gamagrass, <i>Tripsacum dactyloides</i> , L.	2006	136
MOPMC-T-0104	Native Plant Identification	2005	150
MOPMC-T-0105, PA	Compatibility Study Using Warm Season and Cool Season Native Grasses with Legumes and Forbs	2010	106
MOPMC-T-106, BU	Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips	2010	117
MOPMC-P-0107, PA, WL	Evaluation and Release of Big Bluestem, <i>Andropogon gerardii</i> L.	2002	176
MOPMC-T-0208, PA	Testing Warm Season Grasses for Forage Quality	2006	162
MOPMC-T-0209, PA, WL	Evaluation and Release of <i>Paspalum</i> Species 2003	2003	156
MOPMC-T-310-PA,WL	Incorporating Native Warm Season Grasses into Cool Season Pasture with Grazing Management	2007	132
MOPMC-T-0311, RI, BU	Control of Reed Canarygrass in Riparian Buffer Plantings	2006	175
MOPMC-T-0412, WE, WL, RI	Testing Selected Trees for Tolerance to the Herbicide Outrider	2006	185
MOPMC-P-0613-PA, WL	Evaluation and Release of a Shade Tolerant Big Bluestem, <i>Andropogon gerardii</i> , L. for Silvopasture	2010	131
MOPMC-P-0614-PA, WL, BF	Evaluation and Release of Switchgrass, <i>Panicum virgatum</i> L.	2007	138
MOPMS-T-0615-	Direct Seeding of Woody Shrubs for Establishing Shrub Cover for Wildlife Habitat of the Following Species. False Indigo Bush, <i>Amorpha fruticosa</i> ,(a native legume), American Plum, <i>Prunus americana</i> , Roughleaf Dogwood, <i>Cornus drummondii</i> , Fragrant Sumac, <i>Rhus aromatica</i> , Chokecherry <i>Prunus virginiana</i> , Arrowwood <i>Viburnum dentatum</i> , American Hazelnut, <i>Corylus Americana</i> , (all are native woody shrubs.)	2009	117

Study/ Project Number	Title	Annual Technical Report/ Page Reference	
		ATR	Page
MOPMC-T-0716-BF	In-Field Weathering Effects on Biomass Yield and Biofuel Quality of Warm Season Grasses	2010	136
MOPMC-P-0717-PA, WL	Evaluation and Release of a Shade Tolerant Little Bluestem for Silvopasture	2010	145
MOPMC-T-0718-WE, WL	Evaluation of Flood Tolerance of Planted Oak Seedlings Derived from Different Seed Origins	2010	148
MOPMS-T-0719-CR/RI07-016	Critical Area Roadside Vegetation Establishment	2009	144
MOPMC-P-0820-UR	Evaluation and Release of Native Plants for Urban Landscaping	2010	150
MOPMC-P-0821-BF	Evaluation and Release of an Iowa Source, stiff stemmed Indiangrass, <i>Sorghastrum nutans</i> , for Biofuel.	2010	152
MOPMC-P-0822-PA,WL	Inter Center Strain Trial – Yield and Persistence of 11 Big Bluestem Sources in Kansas, Missouri, Arkansas, and Mississippi	2010	163



Elsberry PMC Staff

(left – right) Jerry Kaiser, PM Specialist; Steve Bruckerhoff, PMC Manager; Dean Tapley, Lead Biological Science Technician; Nick Adams, Biological Science Technician; Ron Cordsiemon, Soil Conservationist; Bob Laird, Part-time/Volunteer

For more information about this and other conservation plants, contact your local NRCS field office or Conservation District, or browse the Web at <http://Plant-Materials.nrcs.usda.gov> (Plant Materials) or <http://plants.usda.gov> (PLANTS database).

“The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA’s [TARGET Center](#) at 202-720-2600 (voice and TDD).

“To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.”